

The most current laboratory reference ranges are included in the laboratory report from the LIS.
 Laboratory testing information is also available on The Ohio State Wexner Medical Center Labs Test Catalog Website. <https://theohiostatelabs.testcatalog.org/>

This document applies all laboratory testing at OSU/WMC Clinical Laboratories. This includes:

James Genetics Laboratories; 2001 Polaris Pkwy, Innovation Center, Suite 1500, Columbus OH 43240

Merchbouse Laboratory; 1st Fl Merchbouse Medical Plaza Tower, 2000 Kenny Rd, Columbus OH 43221

Speckman Laboratory; 1145 Olentangy River Rd Rm 2030, Columbus Oh 43212

Clinical Laboratories (LH); 410 West 10th Avenue, Columbus OH 43210

Clinical Laboratories (LHE); 181 Taylor Avenue, Columbus OH 43203

James West Campus Laboratory (JWC); 2121 Kenny Rd, Columbus, OH 43221

Test Name	Synonym	Reference Values		Interpretation	Method Description	Source of Reference Range	Technical Range / AMR	Reportable Range / CRR
		Female	Male					
Arterial Blood Gas (Full Panel)	PH	0-29 days: 7.29-7.45 30+ days: 7.35-7.45	PH 0-29 days: 7.29-7.45 30+ days: 7.35-7.45					
	PCO2	0-29 days: 27-40 mm Hg 30+ days: 32-48 mm Hg	PCO2 0-29 days: 27-40 mm Hg 30+ days: 32-48 mm Hg					
	PO2	83-108 mm Hg	PO2 83-108 mm Hg					
	HCO3	0-29 days: 17-24 mmol/L 30+ days: 22-28 mmol/L	HCO3 0-29 days: 17-24 mmol/L 30+ days: 22-28 mmol/L					
	OSAT	0-365 days: 40-90 % 1+ years: 94-99 %	OSAT 0-365 days: 40-90 % 1+ years: 94-99 %					
	BASE	-3.0-3.0 mmol/L	BASE -3.0-3.0 mmol/L					
	GLUC	0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL	GLUC 0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL					
	LACT	0-51.6 mmol/L	LACT 0-51.6 mmol/L					
	NA	0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L	NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L					
	K	0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-6.0 mmol/L 18+ years: 3.5-5.0 mmol/L	K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-6.0 mmol/L 18+ years: 3.5-5.0 mmol/L					
	ICA	4.00-5.30 mg/dL	ICA 4.00-5.30 mg/dL					
	BGHGB	0-7 days: 13.4-20.0 g/dL 8-14 days: 13.4-20.0 g/dL 15-30 days: 10.8-14.6 g/dL 31-60 days: 9.2-11.4 g/dL 61-179 days: 9.9-12.4 g/dL 180-2 years: 10.2-12.7 g/dL 2-5 years: 10.2-12.7 g/dL 6-11 years: 10.6-13.2 g/dL 12-17 years: 10.8-13.3 g/dL 18+ years: 11.4-15.2 g/dL	BGHGB 0-7 days: 13.9-19.1 g/dL 8-14 days: 13.9-19.1 g/dL 15-30 days: 10.0-15.2 g/dL 31-60 days: 8.9-12.2 g/dL 61-179 days: 9.6-12.4 g/dL 180-2 years: 10.1-12.5 g/dL 2-5 years: 10.2-12.7 g/dL 6-11 years: 10.7-13.4 g/dL 12-17 years: 11.0-14.5 g/dL 18+ years: 11.4-16.8 g/dL					
	BGHCT (Calculated) - Radiometer	0-14 days: 40-57 % 15-30 days: 32-45 % 31-60 days: 28-35 % 61-179 days: 30-37 % 180-2 years: 31-38 % 2-5 years: 31-38 % 6-11 years: 32-40 % 12-17 years: 33-40 % 18+ years: 34-46 %	BGHCT (Calculated) - Radiometer 0-14 days: 40-54 % 15-30 days: 31-45 % 31-60 days: 27-38 % 61-179 days: 29-37 % 180-2 years: 31-38 % 2-5 years: 31-38 % 6-11 years: 32-40 % 12-17 years: 34-44 % 18+ years: 40-50 %					
	BGHCT (Calculated) - GEM 5000/7000	0-15 days: 40-60 % 15-30 days: 32-44 % 30-60 days: 28-34 % 60-180 days: 30-37 % 180-2 years: 31-38 % 2-6 years: 31-38 % 6-12 years: 32-40 % 12-18 years: 32-40 % 18+ years: 34-46 %	BGHCT (Calculated) - GEM 5000/7000 0-15 days: 42-57 % 15-30 days: 30-46 % 30-60 days: 27-38 % 60-180 days: 29-37 % 180-2 years: 30-38 % 2-6 years: 31-38 % 6-12 years: 32-40 % 12-18 years: 33-44 % 18+ years: 40-50 %					
	OZHGB	94-98 %	OZHGB 94-98 %					
	COHGB	≤1.5 %	COHGB ≤1.5 %					
	MTHGB	≤1.5 %	MTHGB ≤1.5 %					

Arterial Blood Gas	GASS	<p>PH 0-29 days: 7.29-7.45 30+ days: 7.35-7.45</p> <p>PCO2 0-29 days: 27-40 mm Hg 30+ days: 32-48 mm Hg</p> <p>PO2 83-108 mm Hg</p> <p>HCO3 0-29 days: 17-24 mmol/L 30+ days: 22-28 mmol/L</p> <p>OSAT 0-365 days: 40-90 % 1+ years: 94-98 %</p> <p>BASE -3.0-3.0 mmol/L</p>		<p>pO2: Amperometry</p> <p>pH, pCO2: Potentiometry</p> <p>Base excess GEM: $HCO_3^- - 24.8 + 16.2 \times (pH - 7.4)$</p> <p>HCO3 GEM: $\text{Log}(HCO_3^-) - pH + \text{log}(pCO_2) - 7.608$ mmol/L</p> <p>sO2: Co-oximetry; $sO_2 = 100 \times O_2Hb\% / (O_2Hb\% + HHb\%)$</p>	See Arterial Blood Gas (Full Panel)	See Arterial Blood Gas (Full Panel)	See Arterial Blood Gas (Full Panel)	
Arterial Blood Gas Plus Serial Lactate	GASSL 2HRLACT	<p>PH 0-29 days: 7.29-7.45 30+ days: 7.35-7.45</p> <p>PCO2 0-29 days: 27-40 mm Hg 30+ days: 32-48 mm Hg</p> <p>PO2 83-108 mm Hg</p> <p>HCO3 0-29 days: 17-24 mmol/L 30+ days: 22-28 mmol/L</p> <p>OSAT 0-365 days: 40-90 % 1+ years: 94-98 %</p> <p>BASE -3.0-3.0 mmol/L</p> <p>LACT 0.5-1.6 mmol/L</p>		<p>pO2, Lactate: Amperometry</p> <p>pH, pCO2: Potentiometry</p> <p>Base excess GEM: $HCO_3^- - 24.8 + 16.2 \times (pH - 7.4)$</p> <p>HCO3 GEM: $\text{Log}(HCO_3^-) - pH + \text{log}(pCO_2) - 7.608$ mmol/L</p> <p>sO2: Co-oximetry; $sO_2 = 100 \times O_2Hb\% / (O_2Hb\% + HHb\%)$</p>	See Arterial Blood Gas (Full Panel)	See Arterial Blood Gas (Full Panel)	See Arterial Blood Gas (Full Panel)	
Arterial Blood Gas, Umbilical Cord	GASCA	<p>PH 7.257-7.33</p> <p>PCO2 41-58 mm Hg</p> <p>PO2 12-24 mm Hg</p> <p>HCO3 20-25 mmol/L</p> <p>OSAT 3-69 %</p> <p>BASE -3.0-3.0 mmol/L</p>		<p>pO2: Amperometry</p> <p>pH, pCO2: Potentiometry</p> <p>Base: Calculation of the expression that approximates the amount of acid or base required to titrate one liter of blood back to a normal pH of 7.40</p> <p>HCO3: Calculation</p> <p>sO2: Co-oximetry</p>	<p>pH, pCO2, pO2, HCO3: Clinical Guide to Laboratory Tests, Tietz, 3rd Edition, 1995.</p> <p>sO2: Brit Journ Obst Gyn 8:2000 Vol 107 pp 987-994. Cord BH O2 SAT in vigorous infants at birth: What is normal?</p> <p>Base: Emml, Cord Blood Gas, R.R. Physician Established Base Excess, 11/2020</p>	<p>PH: 6.80-8.00 PCO2: 5-100 mm Hg PO2: 0-700 mm Hg HCO3: 0-100 mmol/L OSAT: 5-100 % BASE: -30.0-30.0 mmol/L</p>	<p>PH: 6.80-8.00 PCO2: 5-100 mm Hg PO2: 0-700 mmol/L HCO3: 0-100 mmol/L OSAT: 5-100 % BASE: -30.0-30.0 mmol/L</p>	
Carboxyhemoglobin	COHGB	≤1.5 %	Carboxyhemoglobin is elevated in patients exposed to high exogenous sources of carbon monoxide such as cigarette smoke, vehicle exhaust, etc. or increased endogenous production such as hemolysis.	Co-oximetry; $COHb\% = 100 \times (COHb/THb) \%$	See Arterial Blood Gas (Full Panel)	<p>Radiometer ABL800: 0.0-50.0 %</p> <p>GEM 5000: 0.0-75.0%</p>	<p>Radiometer ABL800: 0.0-50.0 %</p> <p>GEM 5000: 0.0-75.0%</p>	
Coometry, Whole Blood	COOXB	<p>O2HGB 94-98 %</p> <p>MTHGB ≤1.5 %</p> <p>COHGB ≤1.5 %</p> <p>BGHGB 0-7 days: 13.4-20.0 g/dL 8-14 days: 13.4-20.0 g/dL 15-30 days: 10.8-14.6 g/dL 31-60 days: 9.2-11.4 g/dL 61-179 days: 9.9-12.4 g/dL 180-2 years: 10.2-12.7 g/dL 2-5 years: 10.2-12.7 g/dL 6-11 years: 10.6-13.2 g/dL 12-17 years: 10.8-13.3 g/dL 18+ years: 11.4-15.2 g/dL</p> <p>OSATA 0-365 days: 40-90 % 1+ years: 94-98 %</p> <p>OSATV 70-80 %</p> <p>OSATMV 60-80%</p>	<p>O2HGB 94-98 %</p> <p>MTHGB ≤1.5 %</p> <p>COHGB ≤1.5 %</p> <p>BGHGB 0-7 days: 13.9-19.1 g/dL 8-14 days: 13.9-19.1 g/dL 15-30 days: 10.0-15.3 g/dL 31-60 days: 8.9-12.7 g/dL 61-179 days: 9.6-12.4 g/dL 180-2 years: 10.1-12.5 g/dL 2-5 years: 10.2-12.7 g/dL 6-11 years: 10.7-13.4 g/dL 12-17 years: 11.0-14.5 g/dL 18+ years: 13.4-16.8 g/dL</p> <p>OSATA 0-365 days: 40-90 % 1+ years: 94-98 %</p> <p>OSATV 70-80 %</p> <p>OSATMV 60-80%</p>		<p>COHGB: Co-oximetry; $COHb\% = 100 \times (COHb/THb) \%$</p> <p>MTHGB: Co-oximetry; $MetHb\% = 100 \times (MetHb/THb) \%$</p> <p>O2HGB: Co-oximetry; $O_2Hb\% = 100 \times (O_2Hb/THb) \%$</p> <p>Total HGB: Co-oximetry; $THb = O_2Hb + COHb + MetHb + HHb$</p> <p>sO2: Co-oximetry; $sO_2 = 100 \times O_2Hb\% / (O_2Hb\% + HHb\%)$</p>	See Arterial Blood Gas (Full Panel)	<p>Radiometer ABL800: O2HGB: 0-100 % MTHGB: 0.0-30.0 % COHGB: 0.0-50.0 % BGHGB: 4.8-23.5 g/dL OSAT: 5-100 %</p> <p>GEM 5000: BGHGB: 5.0-20.0 g/dL O2HGB: 0-100 % COHGB: 0.0-75.0 % MTHGB: 0.0-30.0 % sO2: 0-100 %</p>	<p>Radiometer ABL800: O2HGB: 0-100 % MTHGB: 0.0-30.0 % COHGB: 0.0-50.0 % BGHGB: 4.8-23.5 g/dL OSAT: 5-100 %</p> <p>GEM 5000: BGHGB: 5.0-20.0 g/dL O2HGB: 0-100 % COHGB: 0.0-75.0 % MTHGB: 0.0-30.0 % sO2: 0-100 %</p>
Ionized Calcium (CRRT)	ICACT		Post filter CRRT. When using Citrate anticoagulant protocol, the preferred targeted range is 1-2 mg/dL.	Potentiometry	Email_ICA CRRT RR Physician Established 11-19-2020	1.00-13.00 mg/dL	1.00-13.00 mg/dL	
Ionized Calcium, Serum	ICASST	4.60-5.30 mg/dL		Potentiometry	Blood Gases and Critical Care Testing Physiology, Clinical Interpretations, and Laboratory Applications, 3rd Edition, 2021 (p.102)	1.00-13.00 mg/dL	1.00-13.00 mg/dL	
Ionized Calcium, Whole Blood	ICA	4.60-5.30 mg/dL		Potentiometry	See Arterial Blood Gas (Full Panel)	1.00-13.00 mg/dL	1.00-13.00 mg/dL	
Lactate, Whole Blood		0.5-1.6 mmol/L		Amperometry	See Arterial Blood Gas (Full Panel)	<p>Radiometer ABL800: 0.0-30.0 mmol/L</p> <p>GEM 5000: 0.3-17.0 mmol/L</p>	<p>Radiometer ABL800: 0.0-30.0 mmol/L</p> <p>GEM 5000: 0.3-17.0 mmol/L</p>	
Lactate, Whole Blood, Serial	BELACT 0HRLACT 2HRLACT	0.5-1.6 mmol/L		Amperometry		0.3-17.0 mmol/L	0.3-17.0 mmol/L	
Methemoglobin	MTHGB	≤1.5 %		Co-oximetry; $MetHb\% = 100 \times (MetHb/THb) \%$	See Arterial Blood Gas (Full Panel)	0.0-30.0 %	0.0-30.0 %	
pH, Pleural Fluid	PH			Potentiometry		6.80-8.00	6.80-8.00	

Potassium, Whole Blood		0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L	Potentiometry	See Arterial Blood Gas (Full Panel)	Radiometer ABL800: 1.0-14.0 mmol/L GEM 5000: 1.0-10.0 mmol/L	Radiometer ABL800: 1.0-14.0 mmol/L GEM 5000: 1.0-10.0 mmol/L	
Sodium, Whole Blood	HCZ	0-6 days: 131-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L	Potentiometry	See Arterial Blood Gas (Full Panel)	Radiometer ABL 800: 80-175 mmol/L GEM 5000: 100-180 mmol/L	Radiometer ABL 800: 80-175 mmol/L GEM 5000: 100-180 mmol/L	
Total Hemoglobin	HCZ/GB	0-7 days: 13.4-20.0 g/dL 8-14 days: 13.4-20.0 g/dL 15-30 days: 10.8-14.6 g/dL 31-60 days: 9.2-11.4 g/dL 61-179 days: 9.9-12.4 g/dL 180-2 years: 10.2-12.7 g/dL 2-5 years: 10.2-12.7 g/dL 6-11 years: 10.6-13.2 g/dL 12-17 years: 10.8-13.3 g/dL 18+ years: 11.4-15.2 g/dL	Co-oximetry; TIB = O2IB + COB + MetHb + HHb	See Arterial Blood Gas (Full Panel)	5.0-20.0 g/dL	5.0-20.0 g/dL	
Venous Blood Gas (Full Panel)		PH 7.32-7.43 PCO2 36-52 mm Hg HCO3 22-29 mmol/L OSAT 70-80 % BASE -3.0-3.0 mmol/L GLUC 0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL LACT 0.5-1.6 mmol/L NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L ICA 4.60-5.30 mg/dL BGHGB 0-7 days: 13.9-19.1 g/dL 8-14 days: 13.9-19.1 g/dL 15-30 days: 10.8-14.6 g/dL 31-60 days: 9.2-11.4 g/dL 61-179 days: 9.9-12.4 g/dL 180-2 years: 10.1-12.5 g/dL 2-5 years: 10.2-12.7 g/dL 6-11 years: 10.7-13.4 g/dL 12-17 years: 11.0-14.5 g/dL 18+ years: 13.4-16.8 g/dL BGHCT (Calculated) - Radiometer 0-14 days: 40-57 % 15-30 days: 32-45 % 31-60 days: 28-35 % 61-179 days: 30-37 % 180-2 years: 31-38 % 2-5 years: 31-38 % 6-11 years: 32-40 % 12-17 years: 33-40 % 18+ years: 34-46 % BGHCT (Calculated) - GEM 5000/7000 0-15 days: 40-60 % 15-30 days: 32-44 % 30-60 days: 28-34 % 60-180 days: 30-37 % 180-2 years: 31-38 % 2-6 years: 31-38 % 6-12 years: 32-40 % 12-18 years: 32-40 % 18+ years: 34-46 % O2HGB 94-98 % COHGB ≤1.5 % MTHGB ≤1.5 %	PH 7.32-7.43 PCO2 36-52 mm Hg HCO3 22-29 mmol/L OSAT 70-80 % BASE -3.0-3.0 mmol/L GLUC 0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL LACT 0.5-1.6 mmol/L NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L ICA 4.60-5.30 mg/dL BGHGB 0-7 days: 13.9-19.1 g/dL 8-14 days: 13.9-19.1 g/dL 15-30 days: 10.8-15.3 g/dL 31-60 days: 8.9-12.7 g/dL 61-179 days: 9.6-12.4 g/dL 180-2 years: 10.1-12.5 g/dL 2-5 years: 10.2-12.7 g/dL 6-11 years: 10.7-13.4 g/dL 12-17 years: 11.0-14.5 g/dL 18+ years: 13.4-16.8 g/dL BGHCT (Calculated) - Radiometer 0-14 days: 40-57 % 15-30 days: 32-45 % 31-60 days: 28-35 % 61-179 days: 30-37 % 180-2 years: 31-38 % 2-5 years: 31-38 % 6-11 years: 32-40 % 12-17 years: 33-40 % 18+ years: 34-46 % BGHCT (Calculated) - GEM 5000/7000 0-15 days: 42-57 % 15-30 days: 30-46 % 30-60 days: 27-38 % 60-180 days: 29-37 % 180-2 years: 30-38 % 2-6 years: 31-38 % 6-12 years: 32-40 % 12-18 years: 33-44 % 18+ years: 40-50 % O2HGB 94-98 % COMGB ≤1.5 % MTHGB ≤1.5 %	pO2, Glucose, Lactate: Amperometry pH, pCO2, K, Na, ICA: Potentiometry HCO3 GEM: Log(HCO3 (c)) - pH + log(pCO2) - 7.688 mmol/L HCT: 3.0 x tHb COHGB: Co-oximetry; COHb% = 100 x (COHb/TIb) % MTHGB: Co-oximetry; MetHb% = 100 x (MetHb/TIb) % O2HGB: Co-oximetry; O2Hb% = 100 x (O2Hb/TIb) % Total HGB: Co-oximetry; TIB = O2IB + COB + MetHb + HHb sO2: Co-oximetry; sO2 = 100 x O2Hb%/(O2Hb% + HbHb%) Base excess GEM: HCO3 - 24.8 + 16.2 x (pH - 7.4)	See Arterial Blood Gas (Full Panel)	pH: Clinical Guide to Laboratory Tests 3rd Edition, Tietz, 1995. HCO3 Venous: Clinical Guide to Laboratory Tests 3rd Edition, Tietz, 1995. sO2 Venous: Blood Gas O2Sat: Radiometer Bulletin No: 44 Compendium of reference intervals pO2 Venous: Respiratory, 2014 Feb;19(2):168-75, doi:10.1111/resp.12225. Pub 2014 Jun 3. © 2012 Radiometer Medical Aps. All rights reserved. 995-950, 2012068. pCO2 Venous: Respiratory, 2014 Feb;19(2):168-75, doi:10.1111/resp.12225. Pub 2014 Jun 3. Base excess: Contemporary Practice in Clinical Chemistry 3rd Edition 2016. Chapter 32, Table 32-1 p450. Glucose: Clinical Guide to Laboratory Tests, 3rd Edition Tietz, 1995; Pediatric Reference Ranges, Solfin, 1999. Lactate: ABL 800 Flex Reference Manual, 2008, verified by GEM Validation Study, 2021. Na: Blood Gases and Critical Care Testing Physiology, Clinical Interpretation, and Laboratory Applications, 3rd Edition, 2021 (p.165). Clinical Guide to Laboratory Tests Third Edition, Tietz, 1995. Lower limit adjusted and verified with GEM Validation Study, 2021. K: Clinical Guide to Laboratory Tests, Tietz 3rd Edition, 1995. Clinical Guide to Laboratory Tests, Tietz, 1995. Upper limit adjusted and verified with GEM Validation Study, 2021. ICA: Blood Gases and Critical Care Testing Physiology, Clinical Interpretation, and Laboratory Applications, 3rd Edition, 2021 (p.102). HCT: Derived from total hemoglobin reference interval. ABL 800 FLEX Reference Manual, 2008. Derived from total hemoglobin reference interval based on GEM calculation of HCT. HGB: OSU Internal Normal Range Study, October 2018. Verified with GEM Validation Study, 2021. O2HGB, COHGB, MTHGB: ABL 800 Flex Reference Manual 2008, verified by GEM Validation Study, 2021. Clinical Guide to Laboratory Tests, Third Edition, Tietz, 1995. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics Sixth Edition, 2018.	Radiometer ABL800: PH: 6.80-8.00 PCO2: 5-100 mm Hg PO2: 0-700 mm Hg HCO3: 0-100 mmol/L OSAT: 5-100 % FRO2: 40-100 % BASE: -30.0-30.0 mmol/L GLUC: 1-1,000 mg/dL LACT: 0.0-30.0 mmol/L NA: 80-175 mmol/L K: 1.0-14.0 mmol/L ICA: 1.00-13.00 mg/dL BGHGB: 4.8-23.5 g/dL HCT: 14-71% O2HGB: 0-100 % COHGB: 0.0-50.0 % MTHGB: 0.0-30.0 % GEM 5000: PH: 7.00-7.92 PCO2: 18-125 mm Hg PO2: 35-529 mm Hg HCO3: 0-100 mmol/L sO2: 0-100 % FRO2: 40-100 % BASE: -30.0-30.0 mmol/L GLUC: 16-685 mg/dL LACT: 0.3-17.0 mmol/L NA: 100-180 mmol/L K: 1.0-10.0 mmol/L LACT: 0.3-17.0 mmol/L ICA: 1.00-13.00 mg/dL BGHGB: 5.0-20.0 g/dL HCT: 15-60% O2HGB: 0-100 % COHGB: 0.0-75.0 % MTHGB: 0.0-30.0 % GEM 5000: PH: 6.80-8.00 PCO2: 5-100 mm Hg PO2: 0-700 mm Hg HCO3: 0-100 mmol/L OSAT: 5-100 % FRO2: 40-100 % BASE: -30.0-30.0 mmol/L GLUC: 1-1,000 mg/dL LACT: 0.0-30.0 mmol/L NA: 80-175 mmol/L K: 1.0-14.0 mmol/L ICA: 1.00-13.00 mg/dL BGHGB: 4.8-23.5 g/dL HCT: 14-71% O2HGB: 0-100 % COHGB: 0.0-50.0 % MTHGB: 0.0-30.0 %
	GASALL						

Venous Blood Gas	GASV4		PH 7.32-7.43 PCO2 36-52 mm Hg HCO3 22-29 mmol/L OSAT 70-80 % BASE -3.0-3.0 mmol/L		pO2: Amperometry pH, pCO2: Potentiometry Base excess GEM: $HCO_3^- - 24.8 + 16.2 \times (pH - 7.4)$ HCO3 GEM: $\text{Log}(HCO_3^-) - pH + \text{log}(pCO_2) - 7.608$ mmol/L sO2: Co-oximetry; $sO_2 = 100 \times O_2Hb\% / (O_2Hb\% + Hb\%)$	See Venous Blood Gas (Full Panel)	See Venous Blood Gas (Full Panel)	See Venous Blood Gas (Full Panel)	
Venous Blood Gas Plus Serial Lactate	GASV4 2HBLACT		PH 7.32-7.43 PCO2 36-52 mm Hg HCO3 22-29 mmol/L OSAT 70-80 % BASE -3.0-3.0 mmol/L LACT 0.5-1.6 mmol/L		pO2, Lactate: Amperometry pH, pCO2: Potentiometry Base excess GEM: $HCO_3^- - 24.8 + 16.2 \times (pH - 7.4)$ HCO3 GEM: $\text{Log}(HCO_3^-) - pH + \text{log}(pCO_2) - 7.608$ mmol/L sO2: Co-oximetry; $sO_2 = 100 \times O_2Hb\% / (O_2Hb\% + Hb\%)$	See Venous Blood Gas (Full Panel)	See Venous Blood Gas (Full Panel)	See Venous Blood Gas (Full Panel)	
Venous Blood Gas, Umbilical Cord	GASCV		PH 7.30-7.40 PCO2 33-44 mm Hg PO2 23-35 mm Hg HCO3 16-25 mmol/L OSAT 16-83 % BASE -2.0-2.0 mmol/L		pO2: Amperometry pH, pCO2: Potentiometry Base: Calculation of the expression that approximates the amount of acid or base required to titrate one liter of blood back to a normal pH of 7.40 HCO3: Calculation sO2: Coximetry	<p>pCO2, pH, pO2, HCO3: Clinical Guide to Laboratory Tests, Tietz, 3rd Edition, 1995.</p> <p>Base: Emal. Cord blood Gas RR Physician Established Base Excess, 11-2020.</p> <p>sO2: Brit Jours Obst Gyn 8:2000 Vol 107 pp 987.</p> <p>994 Cord BH O2 SAT in vigorous infants at birth: What is normal?</p>	PH: 6.80-8.00 PCO2: 5-100 mm Hg PO2: 0-700 mm Hg HCO3: 0-100 mmol/L OSAT: 5-100 % BASE: -30.0-30.0 mmol/L	PH: 6.80-8.00 PCO2: 5-100 mm Hg PO2: 0-700 mm Hg HCO3: 0-100 mmol/L OSAT: 5-100 % BASE: -30.0-30.0 mmol/L	
1 Hour Post Glucose	GLUG		70-134 mg/dL		Photometric rate with hexokinase	ADA Standards October 2012, Clinical Guide to Laboratory Tests, Tietz, 1995, Pediatric Reference Ranges, Siddin, 1992	10-800 mg/dL	10-2,400 mg/dL	
Acetaminophen Level	ACTM		≤150.0 mcg/mL		Enzyme immunoassay	Applied Pharmacokinetics: Principles of Therapeutic Drug Monitoring, 2nd Edition 2002 Applied Therapeutics, Inc. and Micromedex On OSU Intranet.	10.0-200.0 mcg/mL	10.0-600.0 mcg/mL	
Albumin-Adjusted Calcium	AAC	ALB 0-30 days: 2.7-4.3 g/dL 31-182 days: 2.9-4.2 g/dL 183-365 days: 3.3-4.8 g/dL 1-19 years: 2.9-4.2 g/dL 19+ years: 3.5-5.0 g/dL CA 0-30 days: 8.4-10.6 mg/dL 31-365 days: 8.9-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL	ALB 0-30 days: 2.6-4.1 g/dL 31-182 days: 2.8-4.0 g/dL 183-365 days: 2.8-4.8 g/dL 1-19 years: 3.2-4.7 g/dL 19+ years: 3.5-5.0 g/dL CA 0-30 days: 8.5-10.6 mg/dL 31-365 days: 8.7-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL		ALB Colorimetric; Bromocresol green CA Photometric; arsenazo AAC Total calcium mg/dL + 0.8*(4-albumin g/dL)	See individual analytes	See individual analytes	See individual analytes	
Albumin	ALB	0-30 days: 2.7-4.3 g/dL 31-182 days: 2.9-4.2 g/dL 183-365 days: 3.3-4.8 g/dL 1-19 years: 2.9-4.2 g/dL 19+ years: 3.5-5.0 g/dL	0-30 days: 2.6-4.1 g/dL 31-182 days: 2.8-4.0 g/dL 183-365 days: 2.8-4.8 g/dL 1-19 years: 3.2-4.7 g/dL 19+ years: 3.5-5.0 g/dL		Colorimetric; Bromocresol green	Tietz 2nd Edition referenced by Beckman Coulter IFU for recumbent adult and verified by OSUWMC Reference Interval Study 2021.	1.5-6.0 g/dL	1.5-18.0 g/dL	
Albumin, CSF	CFMALB		10.0-30.0 mg/dL		Turbidimetry	CCIM Vol 54 issue 2 p285-292 Feb 2016.	1.0-45.0 mg/dL	1.0-450.0 mg/dL	
Albumin, Fluid	FALB			Pleural: Serum- pleural fluid albumin gradients of >1.2 g/dL are consistent with transudates. Peritoneal: Serum-asites albumin gradient (SAAG) of 1.1 g/dL or greater suggests portal hypertension. Pericardial: The reference range has not been established for this fluid specimen. The fluid results should be compared with the concentration in serum/plasma or integrated into the clinical context for interpretation.		Colorimetric; Bromocresol green	Pleural: Roth, B.J., et al. Chest, Vol 98, 546-549, 1990. Peritoneal: Runyon, B.A. Ann Intern Med. 1992;117:215-220.	0.5-6.0 g/dL	0.5-6.0 g/dL
Alcohol (Ethanol), Blood	ETOH		<10 mg/dL	10-50 mg/dL - effects of alcohol may or may not be apparent 30-120 mg/dL - mild euphoria and/or impairment 180-300 mg/dL - disorientation, confusion, dizziness 350-500 mg/dL - unconsciousness, coma, circulation and respiration impairment, possible death > 450 mg/dL - possible death due to respiration impairment Concentrations of blood alcohol causing impairment and/or intoxication may vary by person.		Enzymatic using alcohol dehydrogenase		10-600 mg/dL	10-600 mg/dL
Alk Phosphatase	ALP	0-30 days: 48-406 U/L 31-365 days: 124-341 U/L 1-3 years: 108-317 U/L 4-6 years: 96-297 U/L 7-9 years: 69-325 U/L 10-12 years: 51-332 U/L 13-15 years: 50-162 U/L 16-18 years: 47-119 U/L 19+ years: 32-126 U/L	0-30 days: 75-316 U/L 31-365 days: 82-383 U/L 1-3 years: 104-345 U/L 4-6 years: 93-309 U/L 7-9 years: 86-315 U/L 10-12 years: 42-362 U/L 13-15 years: 74-390 U/L 16-18 years: 52-171 U/L 19+ years: 32-126 U/L		Photometric rate with p-nitro-phenylphosphate (pNPP) in the presence of 2-amino-2-methyl-1-propanol (AMP) at pH 10.4.	OSUWMC Reference Range Study effective 12.31.2003. Verified by OSUWMC Reference Interval Study 2021. Pediatric Reference Ranges, Sokdin, 1999 Synchrotron Performance Verification Manual A22319	5-1,500 U/L	5-15,000 U/L	

ALP ALT AST	ENZ3	<p>ALP 0-30 days: 48-406 U/L 31-365 days: 124-341 U/L 1-3 years: 108-317 U/L 4-6 years: 96-297 U/L 7-9 years: 69-325 U/L 10-12 years: 51-332 U/L 13-15 years: 50-162 U/L 16-18 years: 47-119 U/L 19+ years: 32-126 U/L</p> <p>ALT 0-30 days: 8-25 U/L 31 days-4 years: 8-30 U/L 4-6 years: 8-25 U/L 7-9 years: 8-25 U/L 10-17 years: 8-20 U/L 18+ years: 9-48 U/L</p> <p>AST 0-30 days: 0-49 U/L 31-365 days: 0-80 U/L 1-3 years: 0-70 U/L 4-6 years: 0-60 U/L 7-9 years: 0-42 U/L 10-12 years: 0-38 U/L 13-15 years: 0-33 U/L 16-18 years: 0-31 U/L 19+ years: 10-39 U/L</p>	<p>ALP 0-30 days: 75-316 U/L 31-365 days: 82-383 U/L 1-3 years: 104-345 U/L 4-6 years: 93-309 U/L 7-9 years: 86-315 U/L 10-12 years: 42-362 U/L 13-15 years: 74-390 U/L 16-18 years: 52-171 U/L 19+ years: 32-126 U/L</p> <p>ALT 0-30 days: 8-25 U/L 31-365 days: 8-35 U/L 1-3 years: 8-30 U/L 4-6 years: 8-20 U/L 7-9 years: 8-25 U/L 10-17 years: 8-30 U/L 18+ years: 10-52 U/L</p> <p>AST 0-30 days: 0-52 U/L 31-365 days: 0-66 U/L 1-3 years: 0-57 U/L 4-6 years: 0-49 U/L 7-9 years: 0-43 U/L 10-12 years: 0-39 U/L 13-15 years: 0-40 U/L 16-18 years: 0-40 U/L 19+ years: 10-39 U/L</p>		<p>ALP Photometric rate with p-nitro-phenylphosphate (pNPP) in the presence of 2-amino-2-methyl-1-propanol (AMP) at pH 10.4.</p> <p>ALT Photometric rate with alanine to α-oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate.</p> <p>AST Photometric rate with aspartate and α-oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate.</p>	See individual analytes	See individual analytes	See individual analytes
ALP ALT AST LD	ENZ4	<p>ALP 0-30 days: 48-406 U/L 31-365 days: 124-341 U/L 1-3 years: 108-317 U/L 4-6 years: 96-297 U/L 7-9 years: 69-325 U/L 10-12 years: 51-332 U/L 13-15 years: 50-162 U/L 16-18 years: 47-119 U/L 19+ years: 32-126 U/L</p> <p>ALT 0-30 days: 8-25 U/L 31 days-4 years: 8-30 U/L 4-6 years: 8-25 U/L 7-9 years: 8-25 U/L 10-17 years: 8-20 U/L 18+ years: 9-48 U/L</p> <p>AST 0-30 days: 0-49 U/L 31-365 days: 0-80 U/L 1-3 years: 0-70 U/L 4-6 years: 0-60 U/L 7-9 years: 0-42 U/L 10-12 years: 0-38 U/L 13-15 years: 0-33 U/L 16-18 years: 0-31 U/L 19+ years: 10-39 U/L</p> <p>LD 0-30 days: 145-765 U/L 31-365 days: 190-420 U/L 1-3 years: 165-395 U/L 4-6 years: 155-345 U/L 7-9 years: 140-280 U/L 10-12 years: 120-260 U/L 13-15 years: 100-275 U/L 16-18 years: 105-230 U/L 19+ years: 100-190 U/L</p>	<p>ALP 0-30 days: 75-316 U/L 31-365 days: 82-383 U/L 1-3 years: 104-345 U/L 4-6 years: 93-309 U/L 7-9 years: 86-315 U/L 10-12 years: 42-362 U/L 13-15 years: 74-390 U/L 16-18 years: 52-171 U/L 19+ years: 32-126 U/L</p> <p>ALT 0-30 days: 8-25 U/L 31-365 days: 8-35 U/L 1-3 years: 8-30 U/L 4-6 years: 8-20 U/L 7-9 years: 8-25 U/L 10-17 years: 8-30 U/L 18+ years: 10-52 U/L</p> <p>AST 0-30 days: 0-52 U/L 31-365 days: 0-66 U/L 1-3 years: 0-57 U/L 4-6 years: 0-49 U/L 7-9 years: 0-43 U/L 10-12 years: 0-39 U/L 13-15 years: 0-40 U/L 16-18 years: 0-40 U/L 19+ years: 10-39 U/L</p> <p>LD 0-30 days: 125-735 U/L 31-365 days: 170-450 U/L 1-3 years: 155-345 U/L 4-6 years: 155-345 U/L 7-9 years: 145-300 U/L 10-12 years: 120-325 U/L 13-15 years: 120-290 U/L 16-18 years: 105-235 U/L 19+ years: 100-190 U/L</p>		<p>ALP Photometric rate with p-nitro-phenylphosphate (pNPP) in the presence of 2-amino-2-methyl-1-propanol (AMP) at pH 10.4.</p> <p>ALT Photometric rate with alanine to α-oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate.</p> <p>AST Photometric rate with aspartate and α-oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate.</p> <p>LD Photometric rate</p>	See individual analytes	See individual analytes	See individual analytes
	ENZ4	<p>ALP 0-30 days: 48-406 U/L 31-365 days: 124-341 U/L 1-3 years: 108-317 U/L 4-6 years: 96-297 U/L 7-9 years: 69-325 U/L 10-12 years: 51-332 U/L 13-15 years: 50-162 U/L 16-18 years: 47-119 U/L 19+ years: 32-126 U/L</p> <p>ALT 0-30 days: 8-25 U/L 31 days-4 years: 8-30 U/L 4-6 years: 8-25 U/L 7-9 years: 8-25 U/L 10-17 years: 8-20 U/L 18+ years: 9-48 U/L</p> <p>AST 0-30 days: 0-49 U/L 31-365 days: 0-80 U/L 1-3 years: 0-70 U/L 4-6 years: 0-60 U/L 7-9 years: 0-42 U/L 10-12 years: 0-38 U/L 13-15 years: 0-33 U/L</p>	<p>ALP 0-30 days: 75-316 U/L 31-365 days: 82-383 U/L 1-3 years: 104-345 U/L 4-6 years: 93-309 U/L 7-9 years: 86-315 U/L 10-12 years: 42-362 U/L 13-15 years: 74-390 U/L 16-18 years: 52-171 U/L 19+ years: 32-126 U/L</p> <p>ALT 0-30 days: 8-25 U/L 31-365 days: 8-35 U/L 1-3 years: 8-30 U/L 4-6 years: 8-20 U/L 7-9 years: 8-25 U/L 10-17 years: 8-30 U/L 18+ years: 10-52 U/L</p> <p>AST 0-30 days: 0-52 U/L 31-365 days: 0-66 U/L 1-3 years: 0-57 U/L 4-6 years: 0-49 U/L 7-9 years: 0-43 U/L 10-12 years: 0-39 U/L 13-15 years: 0-40 U/L</p>		<p>ALP Photometric rate with p-nitro-phenylphosphate (pNPP) in the presence of 2-amino-2-methyl-1-propanol (AMP) at pH 10.4.</p> <p>ALT Photometric rate with alanine to α-oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate.</p> <p>AST Photometric rate with aspartate and α-oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate.</p>			

ALP ALT AST LD GGT BILI		<p>16-18 years: 0-31 U/L 19+ years: 10-39 U/L</p> <p>LD 0-30 days: 145-765 U/L 31-365 days: 190-420 U/L 1-3 years: 165-395 U/L 4-6 years: 135-345 U/L 7-9 years: 140-280 U/L 10-12 years: 120-260 U/L 13-15 years: 100-275 U/L 16-18 years: 105-230 U/L 19+ years: 100-190 U/L</p> <p>GGT 0-182 days: 15-132 U/L 183-365 days: 8-39 U/L 1-12 years: 8-22 U/L 13-18 years: 8-24 U/L 19+ years: 8-64 U/L</p> <p>TBIL 0-1 day: 1.4-8.7 mg/dL 1-2 days: 3.4-11.5 mg/dL 3-4 days: 1.5-12.0 mg/dL 5-365 days: 0.3-1.2 mg/dL 1+ years: <1.5 mg/dL</p> <p>DBIL AR: <0.3 mg/dL</p>	<p>16-18 years: 0-40 U/L 19+ years: 10-39 U/L</p> <p>LD 0-30 days: 125-735 U/L 31-365 days: 170-450 U/L 1-3 years: 155-345 U/L 4-6 years: 155-345 U/L 7-9 years: 145-300 U/L 10-12 years: 120-325 U/L 13-15 years: 120-290 U/L 16-18 years: 105-235 U/L 19+ years: 100-190 U/L</p> <p>GGT 0-182 days: 12-122 U/L 183-365 days: 8-39 U/L 1-12 years: 8-22 U/L 13-18 years: 8-42 U/L 19+ years: 8-64 U/L</p> <p>TBIL 0-1 day: 1.4-8.7 mg/dL 1-2 days: 3.4-11.5 mg/dL 3-4 days: 1.5-12.0 mg/dL 5-365 days: 0.3-1.2 mg/dL 1+ years: <1.5 mg/dL</p> <p>DBIL AR: <0.3 mg/dL</p>	<p>phosphate:</p> <p>LD Photometric rate</p> <p>GGT Photometric rate</p> <p>DBIL Photometric with diazonium salt, 3,5-dichloroaniline (DPD).</p> <p>TBIL Photometric with 3,5-dichlorophenyldiazonium tetrafluoroborate (DPD), and caffeine and a surfactant as accelerators.</p>	See individual analytes	See individual analytes	See individual analytes
Alpha 1 Antitrypsin	ALAT		84-218 mg/dL	Turbidimetry	Package Insert. Verified by OSUWMC Reference Interval Study 2021.	10-500 mg/dL	30-5,000 mg/dL
ALT		<p>0-30 days: 8-25 U/L 31 days-4 years: 8-30 U/L 4-6 years: 8-25 U/L 7-9 years: 8-25 U/L 10-17 years: 8-20 U/L 18+ years: 8-48 U/L</p>	<p>0-30 days: 8-25 U/L 31-365 days: 8-35 U/L 1-3 years: 8-30 U/L 4-6 years: 8-20 U/L 7-9 years: 8-25 U/L 10-17 years: 8-30 U/L 18+ years: 10-52 U/L</p>	Photometric rate with alanine to α -oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate.	OSUWMC Reference Range Study effective 12/1/2013; verified by OSUWMC Reference Interval Study 2021. Pediatric Reference Ranges, Soltin, 1999 (Lower end of reference range modified to agree with the linear limits.)	3-500 U/L	3-25,000 U/L
Ammonia	NH3		6-47 μ mol/L	Photometric	Package insert	10-600 μ mol/L	10-3,000 μ mol/L
Ammonia, Arterial	NH3A		6-47 μ mol/L	Photometric	Package insert	10-600 μ mol/L	10-3,000 μ mol/L
AMY, LPA	AMY/LPA		0-30 days: 0-6 U/L 31-182 days: 1-17 U/L	Photometric rate	Prior study verified by OSUWMC Reference Interval Study 2021.	10-2,000 U/L	10-10,000 U/L
Amylase	AMY						
Amylase, Body Fluid				Photometric rate	Pleural: State of the art. The pleura Salu SA Am Rev Respir Dis. 1988;138(1):184. Pancreatic cyst. Eha GH, et al. Am J Gastroenterol. 208;113:464-479.	10-2,000 U/L	10-10,000 U/L
Amylase, Urine Random	FAMY			Photometric rate	The reference range has not been established for random urine specimens. The test result should be integrated into the clinical context for interpretation.		
Amylase, 24 Hour Urine	UAMYR		0-400 U/24 hrs	Photometric rate	Prior study verified by OSUWMC Reference Interval Study 2021.	10-1,500 U/L	10-75,000 U/L
Angiotensin Converting Enzyme	ACE		19-123 U/L	FurylcrokolylPhenylalanylGlycylglycine	Calculated reference range based on correlation study with Mayo Clinic.	14-145 U/L	14-1,450 U/L
Anti Streptolysin O	ASO		<250 IU/mL	Turbidimetry	Package Insert. Verified by OSUWMC Reference Interval Study 2021.	100-1,000 IU/mL	100-10,000 IU/mL
AST		<p>0-30 days: 0-49 U/L 31-365 days: 0-40 U/L 1-3 years: 0-70 U/L 4-6 years: 0-60 U/L 7-9 years: 0-42 U/L 10-12 years: 0-38 U/L 13-15 years: 0-33 U/L 16-18 years: 0-31 U/L 19+ years: 10-39 U/L</p>	<p>0-30 days: 0-52 U/L 31-365 days: 0-60 U/L 1-3 years: 0-57 U/L 4-6 years: 0-49 U/L 7-9 years: 0-43 U/L 10-12 years: 0-39 U/L 13-15 years: 0-40 U/L 16-18 years: 0-40 U/L 19+ years: 10-39 U/L</p>	Photometric rate with aspartate and α -oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate.	Verified by OSUWMC Reference Interval Study 2021.	3-1,000 U/L	3-50,000 U/L
Basic Metabolic Panel		<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>GLUC 0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL</p>	<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>GLUC 0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL</p>	<p>NA/K/CL Indirect ion-selective electrode</p> <p>CO2 Photometric</p> <p>GLUC Photometric rate with hexokinase</p> <p>BUN Photometric rate</p> <p>CREA Kinetic IR</p>	See individual analytes	See individual analytes	See individual analytes

		<p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-0.90 mg/dL 31-365 days: 0.40-0.60 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.50-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.70-1.10 mg/dL 16-18 years: 0.80-1.20 mg/dL 19+ years: 0.50-1.20 mg/dL</p> <p>CA 0-30 days: 8.4-10.6 mg/dL 31-365 days: 8.9-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL</p> <p>OSMO CALC 278-305 mOsm/kg</p> <p>ANION GAP 7-17 mmol/L</p> <p>eGFR ≥60 mL/min/1.73m²</p>	<p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-1.20 mg/dL 31-365 days: 0.40-0.70 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.60-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.60-1.20 mg/dL 16-18 years: 0.80-1.40 mg/dL 19+ years: 0.70-1.30 mg/dL</p> <p>CA 0-30 days: 8.5-10.6 mg/dL 31-365 days: 8.7-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL</p> <p>OSMO CALC 278-305 mOsm/kg</p> <p>ANION GAP 7-17 mmol/L</p> <p>eGFR ≥60 mL/min/1.73m²</p>		<p>CA Photometric, arsenazo</p> <p>ANION GAP (Na + K) - (Cl + CO₂)</p> <p>eGFR CKD-EPI 2021 refn using creatinine, age, and sex</p> <p>OSMO CALC (1.86 (Na + K) + 1.15 (Glucose/B) + (Urea/2.8) + 14 where Na and K are in mmol/L, Glucose and Urea are in mg/dL.</p>			
Beta hCG, Qual, Blood	BMP			Negative	Lateral-flow test using a monoclonal antibody specific to the beta subunit of hCG.	Package insert		Negative, Positive
Beta hCG, Quant, Blood	SPREG			<p>Non-pregnant: <10 mIU/mL Postmenopausal: <10 mIU/mL Male: <10 mIU/mL</p> <p>FEMALE GESTATIONAL AGE 2-4 Weeks: 39-14,388 mIU/mL 5-6 Weeks: 861-88,769 mIU/mL 6-8 Weeks: 8,656-216,085 mIU/mL 8-10 Weeks: 18,700-244,467 mIU/mL 10-12 Weeks: 23,143-181,999 mIU/mL 12-27 Weeks: 6,203-97,171 mIU/mL 24-40 Weeks: 4,360-74,883 mIU/mL</p> <p>Test results cannot be interpreted in absolute evidence for the presence or absence of malignant disease.</p>	Two-site sandwich immunoassay chemiluminescent.	Advia Centaur hCG Package Insert 10634917_EN Rev. F, 2011-04	2.6-1,000.0 mIU/mL	2.6-128,000,000.0 mIU/mL
Beta-Hydroxybutyrate, Serum	QHCGB		<0.27 mmol/L		Photometric	Stabio Package Insert and verified by OSUWMC Reference Interval Study 2021.	0.10-8.00 mmol/L	0.10-24.00 mmol/L
Bicarbonate, Fluid	HIBH			<p>Stool: The reference range has not been established for this fluid specimen. The fluid results should be compared with the concentration in serum/plasma or integrated into the clinical context for interpretation.</p> <p>PEN: Measurement of secretin stimulated electrolytes and enzymes in pancreatic fluid may correlate with the degree of pancreas function. Peak bicarbonate values > 75 mmol/L for screening (time 15 minutes) and >80 mmol/L for confirmatory (time 60 minutes) tests suggest normal pancreas function.</p>	Photometric		5-45 mmol/L	5-90 mmol/L
Bilirubin - Baby	DBIL		<p>DBIL <0.3 mg/dL</p> <p>TBL 0-1 day: 1.4-8.7 mg/dL 1-2 days: 3.4-11.5 mg/dL 3-4 days: 1.5-12.0 mg/dL 5-365 days: 0.3-1.2 mg/dL 1+ years: <1.5 mg/dL</p>		<p>DBIL Photometric with diazonium salt, 3,5-dichloroaniline (DPD).</p> <p>TBL Photometric with 3,5-dichlorophenyl diazotium tetrafluoroborate (DPD), and caffeine and a surfactant as accelerators.</p>	See individual analytes	See individual analytes	See individual analytes
Bilirubin Direct	HBL		<0.3 mg/dL		Photometric with diazonium salt, 3,5-dichloroaniline (DPD).	Clinical Guide to Laboratory Tests, Tietz 1995, verified by OSUWMC Reference Interval Study 2021.	>10.0 mg/dL	0.1-20.0 mg/dL
Bilirubin Total	TBL		<p>0-1 day: 1.4-8.7 mg/dL 1-2 days: 3.4-11.5 mg/dL 3-4 days: 1.5-12.0 mg/dL 5-365 days: 0.3-1.2 mg/dL 1+ years: <1.5 mg/dL</p>		Photometric with 3,5-dichlorophenyl diazotium tetrafluoroborate (DPD), and caffeine and a surfactant as accelerators.	Clinical Guide to Laboratory Tests, Tietz 1995, verified by OSUWMC Reference Interval Study 2021.	0.1-30.0 mg/dL	0.1-90.0 mg/dL
Bilirubin, Total and Direct	TDBL		<p>DBIL <0.3 mg/dL</p> <p>TBL 0-1 day: 1.4-8.7 mg/dL 1-2 days: 3.4-11.5 mg/dL 3-4 days: 1.5-12.0 mg/dL 5-365 days: 0.3-1.2 mg/dL 1+ years: <1.5 mg/dL</p>		<p>DBIL Photometric with diazonium salt, 3,5-dichloroaniline (DPD).</p> <p>TBL Photometric with 3,5-dichlorophenyl diazotium tetrafluoroborate (DPD), and caffeine and a surfactant as accelerators.</p>	See individual analytes	See individual analytes	See individual analytes
Bilirubin, Total, Fluid	FBLT			<p>Peritoneal: Peritoneal bilirubin concentrations greater than that of serum/plasma may suggest bile within the abdomen.</p> <p>Drainage: Drain fluid bilirubin concentration-to-serum/plasma bilirubin concentration ratios exceeding 5 indicates bile leakage.</p>	Photometric with 3,5-dichlorophenyl diazotium tetrafluoroborate (DPD), and caffeine and a surfactant as accelerators.	Peritoneal: Ranyon BA J Clin Gastroenterol. 1987;9(5):543. Drain: Darwin. Gastrointestinal Endosc. 2010 Jan;7(1):99-104.	0.1-30.0 mg/dL	0.1-90.0 mg/dL

BMP without Glucose		<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-0.90 mg/dL 31-365 days: 0.40-0.60 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.50-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.70-1.10 mg/dL 16-18 years: 0.80-1.20 mg/dL 19+ years: 0.50-1.20 mg/dL</p> <p>CA 0-30 days: 8.4-10.6 mg/dL 31-365 days: 8.9-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL</p> <p>ANION GAP 7-17 mmol/L</p> <p>eGFR ≥60 mL/min/1.73m²</p>	<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-1.20 mg/dL 31-365 days: 0.40-0.70 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.60-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.60-1.20 mg/dL 16-18 years: 0.80-1.40 mg/dL 19+ years: 0.70-1.30 mg/dL</p> <p>CA 0-30 days: 8.5-10.6 mg/dL 31-365 days: 8.7-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL</p> <p>ANION GAP 7-17 mmol/L</p> <p>eGFR ≥60 mL/min/1.73m²</p>		<p>NAK/CL Indirect ion-selective electrode</p> <p>CO2 Photometric</p> <p>BUN Photometric rate</p> <p>CREA Kinetic Jaffe</p> <p>CA Photometric, arsenazo</p> <p>ANION GAP (Na + K) - (Cl + CO2)</p> <p>eGFR CKD-EPI 2012 refit using creatinine, age, and sex</p>	See individual analytes	See individual analytes	See individual analytes
B-Type Natriuretic Peptide (Brain)	BNP		0-100 pg/mL		Two site sandwich immunoassay using direct chemiluminescent technology which uses constant amounts of two monoclonal antibodies.	Atellica IM BNP Package Insert 11202199_EN Rev. 05-2020-11	2-4,500 pg/mL	2-4,500 pg/mL
BUN	BUN		7-25 mg/dL		Photometric rate	Beckman Coulter IU for serum verified by OSUWMC Reference Interval Study 2021.	2-130 mg/dL	2-650 mg/dL
BUN CREA	BUN	<p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-0.90 mg/dL 31-365 days: 0.40-0.60 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.50-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.70-1.10 mg/dL 16-18 years: 0.80-1.20 mg/dL 19+ years: 0.50-1.20 mg/dL</p>	<p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-1.20 mg/dL 31-365 days: 0.40-0.70 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.60-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.60-1.20 mg/dL 16-18 years: 0.80-1.40 mg/dL 19+ years: 0.70-1.30 mg/dL</p>		<p>BUN Photometric rate</p> <p>CREA Kinetic Jaffe</p>	See individual analytes	See individual analytes	See individual analytes
C Reactive Protein	CRP		<10.00 mg/L		Measurement of the rate of decrease in light intensity transmitted (increase in absorbance) through particles suspended in solution is the result of complexes formed during the immunological reaction between the CRP of the patient serum and rabbit anti-CRP-antibodies coated on latex particles.	Clinical Guide to Laboratory Tests, Tietz, 2005; Verified by OSUWMC Reference Interval Study 2021.	0.20-80.00 mg/L	0.20-480.00 mg/L
C Reactive Protein for Cardiac Risk	CRP		<2.00 mg/dL		Measurement of the rate of decrease in light intensity transmitted (increase in absorbance) through particles suspended in solution is the result of complexes formed during the immunological reaction between the CRP of the patient serum and rabbit anti-CRP-antibodies coated on latex particles.	Clinical Guide to Laboratory Tests, Tietz, 2005; Verified by OSUWMC Reference Interval Study 2021.	0.20-80.00 mg/L	0.20-480.00 mg/L
C3 Complement	C3		87-200 mg/dL		Turbidimetry	Package Insert. Verified by OSUWMC Reference Interval Study 2021.	15-500 mg/dL	15-1,500 mg/dL
C3, C4			<p>C3 87-200 mg/dL</p> <p>C4 8-52 mg/dL</p>		Turbidimetry	See individual analytes	See individual analytes	See individual analytes
C4 Complement	COMP		8-52 mg/dL		Turbidimetry	Historic Reference Range. Verified by OSUWMC Reference Interval Study 2021.	8-150 mg/dL	8-450 mg/dL
CA 125	CA125		≤30 U/mL		Two-site sandwich immunoassay using direct chemiluminescent technology.	Advia Centaur CA 125H Package Insert 128516 Rev. 04, 2020-03	3-400 U/mL	3-360,000 U/mL
CA 15-3N	CA153	0.0-32.4 U/mL		0.0-32.4 U/mL (see note defined)	Two-site sandwich immunoassay chemiluminescent.	Atellica IM CA 15-3 Package Insert 11206285_EN Rev. 04, 2020-03	3.0-200.0 U/mL	3.0-200,000.0 U/mL
CA 19-9	CA199		≤37.00 U/mL		Two-site sandwich immunoassay using direct chemiluminescent technology.	Clinical Guide to Laboratory Tests, Tietz, 1995; see Source link for Additional Reference Range information	15.00-700.00 U/mL	15.00-33,600,000.00 U/mL
Calcium	CA	0-30 days: 8.4-10.6 mg/dL 31-365 days: 8.9-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL	<p>0-30 days: 8.5-10.6 mg/dL 31-365 days: 8.7-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL</p>		Photometric; arsenazo	Established by OSUWMC Reference Interval Study 2013, verified by OSUWMC Reference Interval study 2021.	4.0-18.0 mg/dL	4.0-18.0 mg/dL
Calcium, Urine 24HR	UCA		100.0-360.0 mg/24 hrs		Photometric; arsenazo			
Calcium/Creat Ratio, Random Urine	CA/CR		<p>0-209 days: <0.86 Ca mg/Crea mg 210-569 days: <0.60 Ca mg/Crea mg 570 days-2 years: <0.42 Ca mg/Crea mg 3+ years: <0.22 Ca mg/Crea mg</p>	Therapeutic Range: 4.0-12.0 mcg/mL	<p>CALCIUM Photometric; arsenazo</p> <p>CREA Kinetic Jaffe</p>	See individual analytes	See individual analytes	See individual analytes
Carbamazepine Total Level	CARB				Competition between drug in the sample and drug labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH) for antibody binding sites. Enzyme activity decreases upon binding to the antibody, so the drug concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts oxidized nicotinamide adenine dinucleotide (NAD) to NADH, resulting in an absorbance change.	Applied Clinical Pharmacokinetics, 2001 Micromedex, OSU Intranet	2.0-20.0 mcg/mL	2.0-100.0 mcg/mL
CEA	CEA		≤5.0 ng/mL		Two-site Sandwich Immunoassay Chemiluminescent	Clinical Guide to Laboratory Tests, Tietz, 1995. See source link for additional Reference Range information.	2.0-100.0 ng/mL	2.0-8,000,000.0 ng/mL
Ceruloplasmin	CERP		20-60 mg/dL		Turbidimetry	Verified by OSUWMC Reference Interval Study 2021.	6-200 mg/dL	6-4,000 mg/dL

Chem 6 (Urea, BUN, CREA)	CHM6	<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-0.90 mg/dL 31-365 days: 0.40-0.60 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.50-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.70-1.10 mg/dL 16-18 years: 0.80-1.20 mg/dL 19+ years: 0.50-1.20 mg/dL</p> <p>ANION GAP 7-17 mmol/L</p> <p>eGFR ≥50 mL/min/1.73m²</p>	<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-1.20 mg/dL 31-365 days: 0.40-0.70 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.60-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.60-1.20 mg/dL 16-18 years: 0.80-1.40 mg/dL 19+ years: 0.70-1.30 mg/dL</p> <p>ANION GAP 7-17 mmol/L</p> <p>eGFR ≥50 mL/min/1.73m²</p>	<p>NA/K/CL Indirect ion-selective electrode</p> <p>CO2 Photometric</p> <p>BUN Photometric rate</p> <p>CREA Kinetic Jaffe</p> <p>ANION GAP (Na + K) - (Cl + CO2)</p> <p>BUN/CREA RATIO BUN/Creatinine</p> <p>eGFR CKD-EPI 2021 refn using creatinine, age, and sex</p>	See individual analytes	See individual analytes	See individual analytes	
Chem 7 (Urea, BUN, CREA, GLUC)	CHM7	<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-0.90 mg/dL 31-365 days: 0.40-0.60 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.50-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.70-1.10 mg/dL 16-18 years: 0.80-1.20 mg/dL 19+ years: 0.50-1.20 mg/dL</p> <p>GLUC 0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL</p> <p>ANION GAP 7-17 mmol/L</p> <p>OSMO (CALC) 278-305 mOsm/kg</p> <p>eGFR ≥50 mL/min/1.73m²</p>	<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-1.20 mg/dL 31-365 days: 0.40-0.70 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.60-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.60-1.20 mg/dL 16-18 years: 0.80-1.40 mg/dL 19+ years: 0.70-1.30 mg/dL</p> <p>GLUC 0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL</p> <p>ANION GAP 7-17 mmol/L</p> <p>OSMO (CALC) 278-305 mOsm/kg</p> <p>eGFR ≥50 mL/min/1.73m²</p>	<p>NA/K/CL Indirect ion-selective electrode</p> <p>CO2 Photometric</p> <p>BUN Photometric rate</p> <p>CREA Kinetic Jaffe</p> <p>GLUC Photometric rate with hexokinase</p> <p>BUN/CREA RATIO BUN/Creatinine</p> <p>eGFR CKD-EPI 2021 refn using creatinine, age, and sex</p>	See individual analytes	See individual analytes	See individual analytes	
Chloride	CL		0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L	Indirect ion-selective electrode	Established by OSUWMC Reference Interval Study 2013, verified by OSUWMC Reference Interval study 2021.	50-200 mmol/L	50-200 mmol/L	
Chloride, 24 HR Urine	UCL		110-250 mmol/24 hrs	Indirect ion-selective electrode	Clinical Guide to Laboratory Tests, Tietz, 1995			
Chloride, Random Urine	UCLR			The reference range has not been established for random urine specimens. The test result should be integrated into the clinical context for interpretation.	Indirect ion-selective electrode	15-400 mmol/L	15-400 mmol/L	
Cholesterol, Body Fluid	FCHEOL			Pleural: Pleural fluid cholesterol concentrations > 200 mg/dL are associated with pseudochyloous effusions. Peritoneal: Peritoneal fluid cholesterol concentrations greater than 32-70 mg/dL may suggest malignant ascites.	Enzymatic colorimetric	Pleural: Hooper C, et al. Thorax. 2010 Aug;65(Suppl2):44-17. McGrath, et al. Int J Clin Pract. 2009 Nov;63(11):163-9 Peritoneal: Black, et al. Crit Rev Clin Lab Sci. 2013;50:107-124.	25-700 mg/dL	25-700 mg/dL
Cholesterol Total	CHOL	<p>0-30 days: 62-155 mg/dL 31-182 days: 62-141 mg/dL 183-365 days: 76-216 mg/dL 1-3 years: 108-193 mg/dL 4-6 years: 108-193 mg/dL 7-9 years: 104-210 mg/dL 10-12 years: 105-218 mg/dL 13-15 years: 108-205 mg/dL 16-18 years: 92-234 mg/dL 19+ years: <200 mg/dL</p>	<p>0-30 days: 54-151 mg/dL 31-182 days: 81-147 mg/dL 183-365 days: 76-179 mg/dL 1-3 years: 85-182 mg/dL 4-6 years: 110-217 mg/dL 7-9 years: 110-211 mg/dL 10-12 years: 105-223 mg/dL 13-15 years: 91-204 mg/dL 16-18 years: 82-192 mg/dL 19+ years: <200 mg/dL</p>	Enzymatic colorimetric	National Cholesterol Education Project (NCEP) Adult Treatment Protocol (ATP-III) (Circulation. 2002;106:3414-3421)	25-700 mg/dL	25-2,100 mg/dL	

Cholesterol/Triglyceride		<p>CHOL 0-30 days: 62-155 mg/dL 31-182 days: 62-141 mg/dL 183-365 days: 76-216 mg/dL 1-3 years: 108-193 mg/dL 4-6 years: 106-192 mg/dL 7-9 years: 108-210 mg/dL 10-12 years: 105-218 mg/dL 13-15 years: 108-205 mg/dL 16-18 years: 92-234 mg/dL 19+ years: <200 mg/dL</p> <p>TRIG 0-3 years: 27-125 mg/dL 4-6 years: 32-116 mg/dL 7-9 years: 28-129 mg/dL 10-11 years: 29-140 mg/dL 12-13 years: 37-130 mg/dL 14-15 years: 38-135 mg/dL 16-19 years: 37-140 mg/dL 20+ years: <150 mg/dL</p>	<p>CHOL 0-30 days: 54-151 mg/dL 31-182 days: 84-147 mg/dL 183-365 days: 76-179 mg/dL 1-3 years: 85-182 mg/dL 4-6 years: 110-217 mg/dL 7-9 years: 110-211 mg/dL 10-12 years: 105-223 mg/dL 13-15 years: 91-204 mg/dL 16-18 years: 82-192 mg/dL 19+ years: <200 mg/dL</p> <p>TRIG 0-3 years: 27-125 mg/dL 4-6 years: 32-116 mg/dL 7-9 years: 28-129 mg/dL 10-11 years: 24-137 mg/dL 12-13 years: 24-145 mg/dL 14-15 years: 34-165 mg/dL 16-19 years: 34-140 mg/dL 20+ years: <150 mg/dL</p>	Enzymatic colorimetric	See individual analytes	See individual analytes	See individual analytes
CK	CHERG	<p>0-30 days: 2-124 U/L 31-182 days: 2-146 U/L 183-365 days: 18-138 U/L 1-3 years: 2-134 U/L 4-6 years: 8-147 U/L 7-9 years: 26-145 U/L 10-12 years: 6-137 U/L 13-15 years: 2-143 U/L 16-18 years: 13-144 U/L 19+ years: 30-184 U/L</p>	<p>0-30 days: 2-183 U/L 31-182 days: 2-129 U/L 183-365 days: 2-143 U/L 1-3 years: 2-163 U/L 4-6 years: 18-158 U/L 7-9 years: 2-177 U/L 10-12 years: 6-217 U/L 13-15 years: 2-251 U/L 16-18 years: 2-238 U/L 19+ years: 30-220 U/L</p>	Photometric rate	Established by OSUWMC Reference Interval Study 2013, verified by OSUWMC Reference Interval study 2021.	10-2,000 U/L	10-200,000 U/L
Clozapine Level	CK		350-1,000 ng/mL	Turbidimetric immunoassay.	Established by OSUWMC Reference Interval Study 2023.		68-1,500 ng/mL
CMPN without Glucose	CKOZ	<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-0.90 mg/dL 31-365 days: 0.40-0.60 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.50-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.70-1.10 mg/dL 16-18 years: 0.80-1.20 mg/dL 19+ years: 0.50-1.20 mg/dL</p> <p>CA 0-30 days: 8.4-10.6 mg/dL 31-365 days: 8.9-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL</p> <p>ANION GAP 7-17 mmol/L</p> <p>eGFR 250 mL/min/1.73m²</p> <p>ALB 0-30 days: 2.7-4.3 g/dL 31-182 days: 2.9-4.2 g/dL 183-365 days: 3.3-4.8 g/dL 1-19 years: 2.9-4.2 g/dL 19+ years: 3.5-5.0 g/dL</p> <p>TBIL 0-1 day: 1.4-8.7 mg/dL 1-2 days: 3.4-11.5 mg/dL 3-4 days: 1.5-12.0 mg/dL 5-365 days: 0.3-1.2 mg/dL 1+ years: <1.5 mg/dL</p> <p>ALP 0-30 days: 48-406 U/L 31-365 days: 124-141 U/L 1-3 years: 108-317 U/L 4-6 years: 96-297 U/L 7-9 years: 69-252 U/L 10-12 years: 51-332 U/L 13-15 years: 50-162 U/L 16-18 years: 67-119 U/L 19+ years: 32-126 U/L</p> <p>ALT 0-30 days: 8-25 U/L</p>	<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-1.20 mg/dL 31-365 days: 0.40-0.70 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.60-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.60-1.20 mg/dL 16-18 years: 0.80-1.40 mg/dL 19+ years: 0.70-1.30 mg/dL</p> <p>CA 0-30 days: 8.5-10.6 mg/dL 31-365 days: 8.7-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL</p> <p>ANION GAP 7-17 mmol/L</p> <p>eGFR 0-4 days: 1,000-39,000 mL/min/1.73m²</p> <p>ALB 0-30 days: 2.6-4.1 g/dL 31-182 days: 2.8-4.6 g/dL 183-365 days: 2.8-4.6 g/dL 1-19 years: 3.2-4.7 g/dL 19+ years: 3.5-5.0 g/dL</p> <p>TBIL 0-1 day: 1.4-8.7 mg/dL 1-2 days: 3.4-11.5 mg/dL 3-4 days: 1.5-12.0 mg/dL 5-365 days: 0.3-1.2 mg/dL 1+ years: <1.5 mg/dL</p> <p>ALP 0-30 days: 75-316 U/L 31-365 days: 82-383 U/L 1-3 years: 104-345 U/L 4-6 years: 93-289 U/L 7-9 years: 86-315 U/L 10-12 years: 42-362 U/L 13-15 years: 74-390 U/L 16-18 years: 52-171 U/L 19+ years: 32-126 U/L</p> <p>ALT 0-30 days: 8-25 U/L 31-365 days: 8-35 U/L</p>	<p>NA/K/CL Indirect ion-selective electrode</p> <p>CO2 Photometric</p> <p>BUN Photometric rate</p> <p>CREA Kinetic Jaffe</p> <p>CA Photometric, arsenazo</p> <p>ANION GAP (Na + K) - (Cl + CO2)</p> <p>eGFR CKD-EPI 2021 refit using creatinine, age, and sex</p> <p>TBIL Photometric with 3,5-dichlorophenylidiazonium tetrafluoroborate (DPD), and caffeine and a surfactant as accelerators.</p> <p>ALP Photometric rate with p-nitro-phenylphosphate (pNPP) in the presence of 2-amino-2-methyl-1-propanol (AMP) at pH 10.4.</p> <p>ALT Photometric rate with alanine to α-oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate.</p> <p>AST Photometric rate with aspartate and α-oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate.</p> <p>TP Colorimetric with cupric ions in an alkaline solution.</p>	See individual analytes	See individual analytes	See individual analytes

		<p>31 days: 4 years: 8-30 U/L 4-6 years: 8-25 U/L 7-9 years: 8-25 U/L 10-17 years: 8-20 U/L 18+ years: 9-48 U/L</p> <p>AST 0-30 days: 0-49 U/L 31-365 days: 0-80 U/L 1-3 years: 0-70 U/L 4-6 years: 0-60 U/L 7-9 years: 0-42 U/L 10-12 years: 0-38 U/L 13-15 years: 0-33 U/L 16-18 years: 0-31 U/L 19+ years: 10-39 U/L</p> <p>TP 0-30 days: 4.2-6.2 g/dL 31-182 days: 4.4-6.6 g/dL 183-365 days: 5.6-7.9 g/dL 1-18 years: 5.7-8.0 g/dL 19+ years: 6.4-8.3 g/dL</p>	<p>1-3 years: 8-30 U/L 4-6 years: 8-20 U/L 7-9 years: 8-25 U/L 10-17 years: 8-30 U/L 18+ years: 10-52 U/L</p> <p>AST 0-30 days: 0-52 U/L 31-365 days: 0-66 U/L 1-3 years: 0-57 U/L 4-6 years: 0-49 U/L 7-9 years: 0-43 U/L 10-12 years: 0-39 U/L 13-15 years: 0-40 U/L 16-18 years: 0-40 U/L 19+ years: 10-39 U/L</p> <p>TP 0-30 days: 4.1-6.3 g/dL 31-182 days: 4.7-6.7 g/dL 183-365 days: 5.5-7.0 g/dL 1-18 years: 5.7-8.0 g/dL 19+ years: 6.4-8.3 g/dL</p>					
	CMPNG							
CO2 Total	CO2		0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L		Photometric	Beckman Coulter IFU verified by OSLWMC Reference Interval Study 2021	5-45 mmol/L	5-45 mmol/L
Comprehensive Metabolic Panel		<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>GLUC 0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL</p> <p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-0.90 mg/dL 31-365 days: 0.40-0.60 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.50-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.70-1.10 mg/dL 16-18 years: 0.80-1.20 mg/dL 19+ years: 0.50-1.20 mg/dL</p> <p>CA 0-30 days: 8.4-10.6 mg/dL 31-365 days: 8.9-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL</p> <p>OSMO 278-305 mOsm/kg</p> <p>ANION GAP 7-17 mmol/L</p> <p>eGFR ≥90 mL/min/1.73m²</p> <p>ALB 0-30 days: 2.7-4.3 g/dL 31-182 days: 2.9-4.2 g/dL 183-365 days: 3.3-4.8 g/dL 1-19 years: 2.9-4.2 g/dL 19+ years: 3.5-5.0 g/dL</p> <p>TBIL 0-1 day: 1.4-8.7 mg/dL 1-2 days: 3.4-11.5 mg/dL 3-4 days: 1.5-12.0 mg/dL 5-365 days: 0.3-1.2 mg/dL 1+ years: <1.5 mg/dL</p> <p>ALP 0-30 days: 48-400 U/L 31-365 days: 24-341 U/L 1-3 years: 108-317 U/L 4-6 years: 96-297 U/L 7-9 years: 69-325 U/L 10-12 years: 51-332 U/L 13-15 years: 50-162 U/L 16-18 years: 67-119 U/L 19+ years: 32-126 U/L</p>	<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>GLUC 0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL</p> <p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-1.20 mg/dL 31-365 days: 0.40-0.70 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.60-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.60-1.20 mg/dL 16-18 years: 0.80-1.40 mg/dL 19+ years: 0.70-1.30 mg/dL</p> <p>CA 0-30 days: 8.5-10.6 mg/dL 31-365 days: 8.7-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL</p> <p>OSMO 278-305 mOsm/kg</p> <p>ANION GAP 7-17 mmol/L</p> <p>eGFR 0-4 days: 1,000-39,000 mL/min</p> <p>ALB 0-30 days: 2.6-4.1 g/dL 31-182 days: 2.8-4.6 g/dL 183-365 days: 2.8-4.8 g/dL 1-19 years: 3.2-4.7 g/dL 19+ years: 3.5-5.0 g/dL</p> <p>TBIL 0-1 day: 1.8-8.7 mg/dL 1-2 days: 3.4-11.5 mg/dL 3-4 days: 1.5-12.0 mg/dL 5-365 days: 0.3-1.2 mg/dL 1+ years: <1.5 mg/dL</p> <p>ALP 0-30 days: 75-316 U/L 31-365 days: 82-383 U/L 1-3 years: 104-345 U/L 4-6 years: 93-289 U/L 7-9 years: 86-315 U/L 10-12 years: 42-362 U/L 13-15 years: 74-390 U/L 16-18 years: 52-171 U/L 19+ years: 32-126 U/L</p>	<p>NA/K/CL Indirect ion-selective electrode</p> <p>CO2 Photometric</p> <p>GLUC Photometric rate with hexokinase</p> <p>BUN Photometric rate</p> <p>CREA Kinetic Jaffe</p> <p>CA Photometric, arsenazo</p> <p>ANION GAP (Na + K) - (Cl + CO2)</p> <p>eGFR CKD-EPI 2021 refit using creatinine, age, and sex</p> <p>OSMO CALC (1.86 (Na + K) + 1.15 (Glucose/18) + (Urea/2.8) + 14 where Na and K are in mmol/L, Glucose and Urea are in mg/dL)</p> <p>TBIL Photometric with 3,5-dichlorophenylidiazonium tetrafluoroborate (DPD), and caffeine and a surfactant as accelerators.</p> <p>ALP Photometric rate with p-nitro-phenylphosphate (pNPP) in the presence of 2-amino-2-methyl-1-propanol (AMP) at pH 10.4.</p> <p>ALT Photometric rate with alanine to α-oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate.</p> <p>AST Photometric rate with aspartate and α-oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate.</p> <p>TP Colorimetric with cupric ions in an alkaline solution.</p>	See individual analytes	See individual analytes	See individual analytes	

		<p>ALT 0-30 days: 8-25 U/L 31-365 days: 8-30 U/L 4-6 years: 8-25 U/L 7-9 years: 8-25 U/L 10-17 years: 8-20 U/L 18+ years: 9-48 U/L</p> <p>AST 0-30 days: 0-49 U/L 31-365 days: 0-80 U/L 1-3 years: 0-70 U/L 4-6 years: 0-60 U/L 7-9 years: 0-42 U/L 10-12 years: 0-38 U/L 13-15 years: 0-33 U/L 16-18 years: 0-31 U/L 19+ years: 10-39 U/L</p> <p>TP 0-30 days: 4.2-6.2 g/dL 31-182 days: 4.4-6.6 g/dL 183-365 days: 5.6-7.9 g/dL 1-8 years: 5.7-8.0 g/dL 19+ years: 6.4-8.3 g/dL</p>	<p>ALT 0-30 days: 8-25 U/L 31-365 days: 8-35 U/L 1-3 years: 8-30 U/L 4-6 years: 8-20 U/L 7-9 years: 8-25 U/L 10-17 years: 8-30 U/L 18+ years: 10-52 U/L</p> <p>AST 0-30 days: 0-52 U/L 31-365 days: 0-66 U/L 1-3 years: 0-57 U/L 4-6 years: 0-49 U/L 7-9 years: 0-43 U/L 10-12 years: 0-39 U/L 13-15 years: 0-40 U/L 16-18 years: 0-40 U/L 19+ years: 10-39 U/L</p> <p>TP 0-30 days: 4.1-6.3 g/dL 31-182 days: 4.7-6.7 g/dL 183-365 days: 5.5-7.0 g/dL 1-18 years: 5.7-8.0 g/dL 19+ years: 6.4-8.3 g/dL</p>						
Cortisol	CORT		3.09-22.40 mcg/dL		Competitive immunoassay using direct chemiluminescent technology.	Atellica IM Cortisol Package Insert 11200390_EN Rev. 03-2020-04	0.50-75.00 mcg/dL	0.50-2,400.00 mcg/dL	
Cortisol (DST) Overnight	CORTDX		<1.80 mcg/dL		Competitive immunoassay using direct chemiluminescent technology.	Atellica IM Cortisol Package Insert 11200391_EN Rev. 03-2020-04	0.50-75.00 mcg/dL	0.50-2,400.00 mcg/dL	
Cortisol, Baseline	CORTB		3.09-22.40 mcg/dL		Competitive immunoassay using direct chemiluminescent technology.	Atellica IM Cortisol Package Insert 11200390_EN Rev. 03-2020-04	0.50-75.00 mcg/dL	0.50-2,400.00 mcg/dL	
Cortisol, 30 Minute	CORT30		≥18.00 mcg/dL		Competitive immunoassay using direct chemiluminescent technology.	Atellica IM Cortisol Package Insert 11200390_EN Rev. 03-2020-06	0.50-75.00 mcg/dL	0.50-2,400.00 mcg/dL	
Cortisol, 60 Minute	CORT60		≥18.00 mcg/dL		Competitive immunoassay using direct chemiluminescent technology.	Atellica IM Cortisol Package Insert 11200390_EN Rev. 03-2020-07	0.50-75.00 mcg/dL	0.50-2,400.00 mcg/dL	
Creatinine Body Fluid	FLCREA			Peritoneal and drainage: Fluid creatinine concentrations that are greater than serum/plasma creatinine concentrations may imply intraperitoneal leakage of urine outside of the urinary tract. Pleural: Pleural fluid creatinine to serum/plasma creatinine concentration ratio > 1 suggests urinothorax.	Kinetic Jaffe	Manahan KJ, et al. Obstet Gynecol. 1999 May;93(5 Pt 1):780-2 Pleural: Toebes, et al. J Thorac Dis. 2017;9(5):1209-1218.	0.20-25.00 mg/dL	0.20-25.00 mg/dL	
Creatinine Serum	CREA	<p>CREA 0-30 days: 0.50-0.90 mg/dL 31-365 days: 0.40-0.60 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.50-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.70-1.10 mg/dL 16-18 years: 0.80-1.20 mg/dL 19+ years: 0.50-1.20 mg/dL</p> <p>eGFR ≥60 mL/min/1.73m2</p>	<p>CREA 0-30 days: 0.50-1.20 mg/dL 31-365 days: 0.40-0.70 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.60-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.60-1.20 mg/dL 16-18 years: 0.80-1.40 mg/dL 19+ years: 0.70-1.30 mg/dL</p> <p>eGFR ≥60 mL/min/1.73m2</p>		Kinetic Jaffe	OSLWMC Reference Range Study effective 12.11.2013; verified by OSLWMC Reference Interval Study 2021. Pediatric Reference Ranges, Soldin, 1999.	CREA: 0.20-25.00 mg/dL eGFR: ≥90 mL/min/1.73m2	CREA: 0.20-25.00 mg/dL eGFR: ≥90 mL/min/1.73m2	
Creatinine, 8 HR Urine	8UR		0-2 years: 0.70-2.00 g/24 hrs 3-8 years: 0.11-0.68 g/24 hrs 9-12 years: 0.17-1.41 g/24 hrs 13-17 years: 0.29-1.87 g/24 hrs 18+ years: 0.60-1.80 g/24 hrs	0-2 years: 0.70-2.00 g/24 hrs 3-8 years: 0.11-0.68 g/24 hrs 9-12 years: 0.17-1.41 g/24 hrs 13-17 years: 0.29-1.87 g/24 hrs 18+ years: 0.80-2.00 g/24 hrs		Kinetic Jaffe	1.00-300.00 mg/dL	1.00-900.00 mg/dL	
Creatinine, 24 HR Urine	24UR		0-2 years: 0.70-2.00 g/24 hrs 3-8 years: 0.11-0.68 g/24 hrs 9-12 years: 0.17-1.41 g/24 hrs 13-17 years: 0.29-1.87 g/24 hrs 18+ years: 0.60-1.80 g/24 hrs	0-2 years: 0.70-2.00 g/24 hrs 3-8 years: 0.11-0.68 g/24 hrs 9-12 years: 0.17-1.41 g/24 hrs 13-17 years: 0.29-1.87 g/24 hrs 18+ years: 0.80-2.00 g/24 hrs		Kinetic Jaffe	NKDPE traceable Clinical Guide to Laboratory Tests, Tietz, 1995; Pediatric Reference Ranges, Soldin, 1999	1.00-300.00 mg/dL	1.00-900.00 mg/dL
Creatinine, Random Urine	UR			The reference range has not been established for random urine specimens. The test result should be integrated into the clinical context for interpretation.	Kinetic Jaffe		1.00-300.00 mg/dL	1.00-900.00 mg/dL	
Cystatin C and Creatinine with estimated GFR	CYSC	<p>CYSC 0.51-1.05 mg/L</p> <p>CREA 0-30 days: 0.50-0.90 mg/dL 31-365 days: 0.40-0.60 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.50-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.70-1.10 mg/dL 16-18 years: 0.80-1.20 mg/dL 19+ years: 0.50-1.20 mg/dL</p> <p>eGFR ≥60 mL/min/1.73m2</p>	<p>CYSC 0.51-1.05 mg/L</p> <p>CREA 0-30 days: 0.50-1.20 mg/dL 31-365 days: 0.40-0.70 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.60-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.60-1.20 mg/dL 16-18 years: 0.80-1.40 mg/dL 19+ years: 0.70-1.30 mg/dL</p> <p>eGFR ≥60 mL/min/1.73m2</p>		<p>CYSC Turkishometric CREA Kinetic Jaffe</p> <p>CKD-EPI 2021 Calculation: eGFR_{Scr-cys} = 135 x min(Scr/c, 1) x max(Scr/c, 1) x 0.54 x min(Scr/cy0.8, 1.0) x 2.3 x max(Scr/cy0.8, 1.0) x 0.778 x 0.9961 Age x 0.963 [if female] where c = 0.7 (female) or 0.9 (male) s = -0.219 (female) or -0.144 (male) Scr = serum creatinine in mg/dL Scr_c = serum cystatin C in mg/L Age (years)</p> <p>The "min (Scr/c, 1)" factor indicates the minimum of Scr/c or 1.0, "max(Scr/c, 1)" indicates the maximum of Scr/c or 1.0, "min(Scr/cy0.8, 1)" indicates the minimum of Scr/cy0.8 or 1.0, and "max(Scr/cy0.8, 1)" indicates the maximum of Scr/cy0.8 or 1.0.</p>	CREA: 0.20-25.00 mg/dL CYSC: 0.40-8.00 mg/L	CREA: 0.20-25.00 mg/dL CYSC: 0.40-30.00 mg/L		
Digoxin Level	DIG		Therapeutic Range: 0.5-1.0 ng/mL		Enzyme Immunoassay	Applied Clinical Pharmacokinetics, Bauer, 2001	0.3-5.0 ng/mL	0.3-10.0 ng/mL	
Estradiol, Enhanced	EE2B			<p>Male: 19+ years: <11.8-39.8</p> <p>Menstruating Females: Follicular phase: 19.5-144.2 Midcycle peak: 63.9-356.7 Luteal phase: 55.8-214.2</p> <p>Post-menopausal: <11.8-32.2</p> <p>This test is not recommended for patients receiving Fulvestrant (Faslodex) due to possible false elevations. EE2 levels vary widely throughout the menstrual cycle.</p>	Competitive assay format. The endogenous estradiol contained in a sample is released from its binding proteins by a releasing agent. Then, a sheep monoclonal anti-estradiol antibody labeled with acridinium ester is added to bind available estradiol. Finally, an oxidant derivative capture solid phase is added to the reaction to compete with estradiol for the binding of the acridinium-labeled antibody. After washing, acid and base are dispensed to initiate the chemiluminescent reaction.	Abria Centaur, Enhanced Estradiol (eE2) Package Insert 10491467 Rev. C, 2010-09; Pediatric Reference Ranges, Soldin, 1999	11.8-3,000.0 pg/mL	11.8-150,000.0 pg/mL	
ELS Pancreatic Lab (Fluid CEA, AMY)	FCAR			<p>CEA: Fluid CEA values >192 ng/mL may indicate mucinous cystic lesions of the pancreas. Results <192 ng/mL require clinical correlation with patient history and other imaging modalities. This chemiluminescent test was developed and its performance characteristics determined by the Critical Care Laboratory at The Ohio State University Wexner Medical Center. It has not been cleared or approved by the FDA. The laboratory is regulated under CLIA as qualified to perform high-complexity testing. This test is used for clinical purposes. It should not be regarded as investigational or for research.</p> <p>AMY: Very low pancreatic cyst fluid amylase concentrations (< 250 U/L) exclude a pseudocyst in the majority of cases.</p>	<p>AMY Photometric rate</p> <p>CEA Two-site Sandwich Immunoassay Chemiluminescent</p>	See individual analytes	See individual analytes	See individual analytes	
Ferritin	FERR	<p>0-365 days: 5.7-421.0 ng/mL 1-12 years: 12.8-88.7 ng/mL 13-18 years: 6.8-75.6 ng/mL 19+ years: 7.3-270.7 ng/mL</p>	<p>0-365 days: 5.7-421.0 ng/mL 1-12 years: 12.8-88.7 ng/mL 13-18 years: 10.9-135.0 ng/mL 19+ years: 10.5-307.3 ng/mL</p>		Two-site Sandwich Immunoassay Chemiluminescent.	Siemens Atellica IM Reference Interval Verification Study 2023 Siemens Atellica IM Reference Interval Verification Study Summary (Amendment)	0.9-1,650.0 ng/mL	0.9-1,650,000.0 ng/mL	

Folate, Serum	FOH.SB	0-1 years: 6.30-22.70 ng/mL 2-3 years: 1.70-15.70 ng/mL 4-6 years: 2.70-14.10 ng/mL 7-9 years: 2.40-13.40 ng/mL 10-12 years: 1.00-10.20 ng/mL 13-18 years: 1.20-7.20 ng/mL 19+ years: >5.38 ng/mL	0-1 years: 7.20-22.40 ng/mL 2-3 years: 2.50-15.00 ng/mL 4-6 years: 0.50-13.00 ng/mL 7-9 years: 2.30-11.90 ng/mL 10-12 years: 1.50-10.80 ng/mL 13-18 years: 1.20-8.80 ng/mL 19+ years: >5.38 ng/mL	Competitive immunoassay using direct chemiluminescent technology.	Aotlica IM Folate Package Insert 11200602_EN Rev. 04-2020-11; Pediatric Reference Ranges, Saldin, 1999	0.56-24.00 ng/mL	0.56-860.00 ng/mL		
FSH	FSH	0-365 days: 1.9-22.9 mIU/mL 1-8 years: 1.1-9.5 mIU/mL 9-11 years: 1.5-9.2 mIU/mL 12-18 years: 1.9-11.2 mIU/mL	0-365 days: 0.8-5.2 mIU/mL 1-8 years: 0.8-2.5 mIU/mL 9-11 years: 1.0-4.2 mIU/mL 12-18 years: 2.3-10.2 mIU/mL	Two-site sandwich immunoassay using direct chemiluminometric technology.	Aotlica IM FSH Package Insert 11200384_EN Rev. 06-2020-09	0.3-200.0 mIU/mL	0.3-4,400.0 mIU/mL		
GGT	GGT	0-182 days: 15-132 U/L 183-365 days: 8-99 U/L 1-12 years: 8-22 U/L 13-18 years: 8-24 U/L 19+ years: 8-64 U/L	0-182 days: 12-122 U/L 183-365 days: 8-99 U/L 1-12 years: 8-22 U/L 13-18 years: 8-42 U/L 19+ years: 8-64 U/L	Photometric rate	Beckman Coulter IFU verified by ORLWMC Reference Interval Study 2021 (lower end modified); Pediatric Reference Ranges, Saldin, 1999 (Lower end of reference range modified to agree with the linear limit)	3-1,200 U/L	3-6,000 U/L		
Gestational 3HR Glucose Tolerance Test	GTF GT1H GT2H GT3H			The upper limits of the reference intervals for each fasting, 1 hour, 2 hour, and 3 hour samples are based on the Carpenter-Coustan criteria. According to the ADA practice recommendations, the diagnosis of GDM is made when at least 2 of the four plasma glucose levels meet or exceed the Carpenter-Coustan criteria, although The American College of Obstetricians and Gynecologists notes that some clinicians choose to use just one elevated value.	Photometric rate with hexokinase		10-800 mg/dL	10-2,400 mg/dL	
Glucose	GLUC		0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL	The reference interval provided is for plasma glucose measurements from samples collected after an 8 hour fast (no caloric intake).	ADA Standards October 2012, Clinical Guide to Laboratory Tests, Tietz, 1995; Pediatric Reference Ranges, Saldin, 1999.	10-800 mg/dL	10-2,400 mg/dL		
Glucose Body Fluid	GLUC			Amniotic: Amniotic fluid glucose concentrations < 10 mg/dL are consistent with intra-amniotic inflammation in patients with prelabor rupture of membranes. Peritoneal: Peritoneal glucose concentrations >50 mg/dL (2.8 mmol/L) are consistent with spontaneous bacterial peritonitis and concentrations below this are consistent with secondary bacterial peritonitis due to gut perforation. Pericardial: Pericardial fluid glucose to serum/plasma glucose ratios are 1.0 in presumed normal patients. Pancreatic Body Cyst: Pancreatic Cyst Glucose measurements of <50 mg/dL are suggestive of a mucinous lesion. Pleural: Pleural fluid glucose concentrations are equivalent to serum/plasma glucose concentrations in the absence of pleural pathology. Pleural fluid glucose concentrations < 60 mg/dL may indicate parapneumonic or malignant effusion. Other less common effusions associated with low glucose concentrations include hemothorax, tuberculosis, rheumatoid pleuritis, Churg-Strauss syndrome, paragonimiasis, and lupus pleuritis. Synovial: Synovial fluid glucose concentrations lower than glucose in serum/plasma are associated with infection.	Photometric rate with hexokinase	Amniotic: Genzler-Bosquet, et al. J Matern Fetal Med. Jul-Aug 1999;8(4):155-8. Pancreatic cyst: Carr, et al. Surgery. 2018 Mar;163(3):690-695. Peritoneal: Ranjan BA, Hoefs JC. Hepatology. 1985;5(2):237. Pericardial: Ben-Haim S, et al. Am J Med 2005;118:636-40. Pleural: Toebes, et al. J Thorac Dis. 2017;9(5):1209-1218. Light. RW. N Engl J Med. 2002 Jun 20;346(25):1971-1977 Synovial: Margretem, et al. JAMA. 2007;297(13):1478-1488.	10-800 mg/dL	10-800 mg/dL	
Glucose CSF	GLUC		40-70 mg/dL		Clinical Guide to Laboratory Tests, Tietz, 1995	10-800 mg/dL	10-800 mg/dL		
Glucose Fasting	GLUF		0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL		ADA Standards October 2012, Clinical Guide to Laboratory Tests, Tietz, 1995; Pediatric Reference Ranges, Saldin, 1999.	10-800 mg/dL	10-2,400 mg/dL		
Haptoglobin	HAPTO		44-215 mg/dL		Package Insert. Verified by ORLWMC Reference Interval Study 2021.	30-400 mg/dL	30-1,200 mg/dL		
hCG Qualitative, Urine	hCG			Negative	Lateral-flow test using a monoclonal antibody specific to the beta subunit of hCG. Package insert		Negative, Positive		
hCG, Quant (Tumor Marker)	hCGTM		<10.0 mIU/mL		Two-site sandwich immunoassay chemiluminescent.	Advia Centaur hCG Package Insert 10634917_EN Rev. F, 2011.04.	2.6-1,000.0 mIU/mL	2.6-128,000,000.0 mIU/mL	
HDL, Cholesterol	HDL		0-9 years: 35-82 mg/dL 10-13 years: 36-84 mg/dL 14-19 years: 35-65 mg/dL 20+ years: >60 mg/dL		Enzymatic colorimetric	National Cholesterol Education Project (NCEP) Adult Treatment Protocol (ATP-III) (Circulation. 2002;106:3143-3421)	3-200 mg/dL	3-200 mg/dL	
		ALB 0-30 days: 2.7-4.3 g/dL 31-182 days: 2.9-4.2 g/dL 183-365 days: 3.3-4.8 g/dL 1-9 years: 2.9-4.2 g/dL 19+ years: 3.5-5.0 g/dL DBIL <0.3 mg/dL TBIL 0-1 day: 1.4-8.7 mg/dL 1-2 days: 3.4-11.5 mg/dL 3-4 days: 1.5-12.0 mg/dL 5-365 days: 0.3-1.2 mg/dL 1+ years: <1.5 mg/dL ALP 0-30 days: 48-406 U/L 31-365 days: 124-341 U/L 1-3 years: 108-317 U/L 4-6 years: 96-297 U/L 7-9 years: 69-328 U/L 10-12 years: 51-332 U/L 13-15 years: 50-162 U/L	ALB 0-30 days: 2.6-4.1 g/dL 31-182 days: 2.8-4.6 g/dL 183-365 days: 2.8-4.6 g/dL 1-9 years: 3.2-4.7 g/dL 19+ years: 3.5-5.0 g/dL DBIL <0.3 mg/dL TBIL 0-1 day: 1.4-8.7 mg/dL 1-2 days: 3.4-11.5 mg/dL 3-4 days: 1.5-12.0 mg/dL 5-365 days: 0.3-1.2 mg/dL 1+ years: <1.5 mg/dL ALP 0-30 days: 75-316 U/L 31-365 days: 82-383 U/L 1-3 years: 104-345 U/L 4-6 years: 93-309 U/L 7-9 years: 86-315 U/L 10-12 years: 42-362 U/L 13-15 years: 74-390 U/L			ALB Colorimetric; Bismoresol green. DBIL Photometric with diazonium salt, 3,5-dichloroaniline (DPD). TBIL Photometric with 3,5-dichlorophenyldiazonium tetrafluoroborate (DPD), and caffeine and a surfactant as accelerators. ALP			

Hepatic Function Panel		16-18 years: 47-119 U/L 19+ years: 32-126 U/L ALT 0-30 days: 8-25 U/L 31-365 days: 8-30 U/L 4-6 years: 8-25 U/L 7-9 years: 8-25 U/L 10-17 years: 8-20 U/L 18+ years: 9-48 U/L AST 0-30 days: 0-49 U/L 31-365 days: 0-80 U/L 1-3 years: 0-70 U/L 4-6 years: 0-60 U/L 7-9 years: 0-42 U/L 10-12 years: 0-38 U/L 13-15 years: 0-33 U/L 16-18 years: 0-31 U/L 19+ years: 10-39 U/L TP 0-30 days: 4.2-6.2 g/dL 31-182 days: 4.4-6.6 g/dL 183-365 days: 5.6-7.9 g/dL 1-18 years: 5.7-8.0 g/dL 19+ years: 6.4-8.3 g/dL	16-18 years: 5.0-11.1 U/L 19+ years: 32-126 U/L ALT 0-30 days: 8-25 U/L 31-365 days: 8-35 U/L 1-3 years: 8-30 U/L 4-6 years: 8-20 U/L 7-9 years: 8-25 U/L 10-17 years: 8-30 U/L 18+ years: 10-52 U/L AST 0-30 days: 0-52 U/L 31-365 days: 0-66 U/L 1-3 years: 0-57 U/L 4-6 years: 0-49 U/L 7-9 years: 0-43 U/L 10-12 years: 0-39 U/L 13-15 years: 0-40 U/L 16-18 years: 0-40 U/L 19+ years: 10-39 U/L TP 0-30 days: 4.1-6.3 g/dL 31-182 days: 4.7-6.7 g/dL 183-365 days: 5.5-7.0 g/dL 1-18 years: 5.7-8.0 g/dL 19+ years: 6.4-8.3 g/dL		Phenometric rate with p-nitro-phenylphosphate (PNPP) in the presence of 2-amino-2-methyl-1-propanol (AMP) at pH 10.4. ALT Phenometric rate with alanine to α -oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate. AST Phenometric rate with aspartate and α -oxoglutarate to form pyruvate and glutamate. This assay does not contain pyridoxyl-5-phosphate. TP Colorimetric with cupric ions in an alkaline solution.	See individual analytes	See individual analytes	See individual analytes
High Sensitivity Troponin I - Single Order	HST1LX	<34 ng/L	<53 ng/L	Three-site sandwich immunoassay using direct chemiluminescent technology.	Atellica IM Tahi Package Insert 11200498_EN Rev. 06, 2019-06.	3-25,000 ng/L	3-2,000,000 ng/L	
High Sensitivity Troponin I x2	HST1Hml HST1Hfr	<34 ng/L	<53 ng/L	Three-site sandwich immunoassay using direct chemiluminescent technology.	Atellica IM Tahi Package Insert 11200498_EN Rev. 06, 2019-06.	3-25,000 ng/L	3-2,000,000 ng/L	
Homocysteine	HCBCYS		3.7-13.9 μ mol/L	Competitive immunoassay using direct chemiluminescent technology.	Atellica IM Homocysteine Package Insert 1095362_EN Rev. 04-2021-03	0.5-65.0 μ mol/L	0.5-130.0 μ mol/L	
IgA	IgA	0-30 days: 0-10 mg/dL 31-182 days: \leq 42 mg/dL 183-365 days: 6-68 mg/dL 1-3 years: 15-111 mg/dL 4-6 years: 33-166 mg/dL 7-9 years: 28-180 mg/dL 10-12 years: 55-193 mg/dL 13-15 years: 62-241 mg/dL 16-18 years: 69-262 mg/dL 19-59 years: 66-433 mg/dL 60+ years: 90-410 mg/dL	0-30 days: \leq 11 mg/dL 31-182 days: \leq 40 mg/dL 183-365 days: 1-82 mg/dL 1-3 years: 9-137 mg/dL 4-6 years: 44-187 mg/dL 7-9 years: 38-204 mg/dL 10-12 years: 46-218 mg/dL 13-15 years: 29-251 mg/dL 16-18 years: 68-259 mg/dL 19-59 years: 66-433 mg/dL 60+ years: 90-410 mg/dL	Turbidimetry	Package Insert, Verified by OSUWMC Reference Interval Study 2021.	10-700 mg/dL	10-14,000 mg/dL	
IgG	IgG	0-30 days: 162-872 mg/dL 31-182 days: 311-664 mg/dL 183-365 days: 325-647 mg/dL 1-3 years: 451-1,202 mg/dL 4-6 years: 560-1,319 mg/dL 7-9 years: 485-1,473 mg/dL 10-12 years: 586-1,609 mg/dL 13-15 years: 540-1,640 mg/dL 16-18 years: 804-1,817 mg/dL 19-59 years: 600-1,714 mg/dL 60+ years: 600-1,560 mg/dL	0-30 days: 197-833 mg/dL 31-182 days: 140-533 mg/dL 183-365 days: 130-823 mg/dL 1-3 years: 413-1,112 mg/dL 4-6 years: 468-1,328 mg/dL 7-9 years: 582-1,441 mg/dL 10-12 years: 685-1,620 mg/dL 13-15 years: 590-1,600 mg/dL 16-18 years: 522-1,703 mg/dL 19-59 years: 600-1,714 mg/dL 60+ years: 600-1,560 mg/dL	Turbidimetry	OSUWMC Immoglobulin Reference Range Study, Verified by OSUWMC Reference Interval Study 2021.	75-3,000 mg/dL	75-60,000 mg/dL	
IgM	IgM	0-29 days: 1-57 mg/dL 30-182 days: <128 mg/dL 183-365 days: <131 mg/dL 1-3 years: 35-184 mg/dL 4-6 years: 42-184 mg/dL 7-9 years: 30-165 mg/dL 10-12 years: 62-211 mg/dL 13-15 years: 34-225 mg/dL 16-18 years: 45-224 mg/dL 19-59 years: 45-281 mg/dL 60+ years: 30-360 mg/dL	0-29 days: 0-65 mg/dL 30-182 days: 6-84 mg/dL 183-365 days: 15-117 mg/dL 1-3 years: 30-166 mg/dL 4-6 years: 31-151 mg/dL 7-9 years: 21-140 mg/dL 10-12 years: 27-151 mg/dL 13-15 years: 26-184 mg/dL 16-18 years: 28-179 mg/dL 19-59 years: 45-281 mg/dL 60+ years: 30-360 mg/dL	Turbidimetry	Package Insert, Verified by OSUWMC Reference Interval Study 2021.	20-500 mg/dL	20-50,000 mg/dL	
Immunoglobulins IgG IgA IgM	IgA IgG IgM	IgA 0-30 days: 0-10 mg/dL 31-182 days: \leq 42 mg/dL 183-365 days: 6-68 mg/dL 1-3 years: 15-111 mg/dL 4-6 years: 33-166 mg/dL 7-9 years: 28-180 mg/dL 10-12 years: 55-193 mg/dL 13-15 years: 62-241 mg/dL 16-18 years: 69-262 mg/dL 19-59 years: 66-433 mg/dL 60+ years: 90-410 mg/dL IgG 0-30 days: 162-872 mg/dL 31-182 days: 311-664 mg/dL 183-365 days: 325-647 mg/dL 1-3 years: 451-1,202 mg/dL 4-6 years: 560-1,319 mg/dL 7-9 years: 485-1,473 mg/dL 10-12 years: 586-1,609 mg/dL 13-15 years: 540-1,640 mg/dL 16-18 years: 804-1,817 mg/dL 19-59 years: 600-1,714 mg/dL 60+ years: 600-1,560 mg/dL IgM 0-29 days: 1-57 mg/dL 30-182 days: <128 mg/dL 183-365 days: <131 mg/dL 1-3 years: 35-184 mg/dL 4-6 years: 42-184 mg/dL 7-9 years: 30-165 mg/dL 10-12 years: 62-211 mg/dL 13-15 years: 34-225 mg/dL 16-18 years: 45-224 mg/dL 19-59 years: 45-281 mg/dL 60+ years: 30-360 mg/dL	IgA 0-30 days: \leq 11 mg/dL 31-182 days: \leq 40 mg/dL 183-365 days: 1-82 mg/dL 1-3 years: 9-137 mg/dL 4-6 years: 44-187 mg/dL 7-9 years: 38-204 mg/dL 10-12 years: 46-218 mg/dL 13-15 years: 29-251 mg/dL 16-18 years: 68-259 mg/dL 19-59 years: 66-433 mg/dL 60+ years: 90-410 mg/dL IgG 0-30 days: 197-833 mg/dL 31-182 days: 140-533 mg/dL 183-365 days: 130-823 mg/dL 1-3 years: 413-1,112 mg/dL 4-6 years: 468-1,328 mg/dL 7-9 years: 582-1,441 mg/dL 10-12 years: 685-1,620 mg/dL 13-15 years: 590-1,600 mg/dL 16-18 years: 522-1,703 mg/dL 19-59 years: 600-1,714 mg/dL 60+ years: 600-1,560 mg/dL IgM 0-29 days: 0-65 mg/dL 30-182 days: 6-84 mg/dL 183-365 days: 15-117 mg/dL 1-3 years: 30-166 mg/dL 4-6 years: 31-151 mg/dL 7-9 years: 21-140 mg/dL 10-12 years: 27-151 mg/dL 13-15 years: 26-184 mg/dL 16-18 years: 28-179 mg/dL 19-59 years: 45-281 mg/dL 60+ years: 30-360 mg/dL	Turbidimetry	See individual analytes	See individual analytes	See individual analytes	
Iron	IRON	0-30 days: 29-127 mcg/dL 31-365 days: 25-126 mcg/dL 1-3 years: 25-101 mcg/dL 4-6 years: 28-93 mcg/dL 7-9 years: 30-104 mcg/dL 10-12 years: 32-104 mcg/dL 13-15 years: 30-109 mcg/dL 16-18 years: 33-102 mcg/dL 19+ years: 40-174 mcg/dL	0-30 days: 32-112 mcg/dL 31-365 days: 27-109 mcg/dL 1-3 years: 28-91 mcg/dL 4-6 years: 25-115 mcg/dL 7-9 years: 27-96 mcg/dL 10-12 years: 26-112 mcg/dL 13-15 years: 26-110 mcg/dL 16-18 years: 27-138 mcg/dL 19+ years: 40-174 mcg/dL	Colorimetric	Established by OSUWMC Reference Interval Study 2013, verified by OSUWMC Reference Interval study 2021.	10-1,000 mcg/dL	10-2,000 mcg/dL	

Iron/Iron Binding/Transferin		<p>IRON 0-30 days: 29-127 mcg/dL 31-365 days: 25-126 mcg/dL 1-3 years: 25-101 mcg/dL 4-6 years: 28-93 mcg/dL 7-9 years: 30-104 mcg/dL 10-12 years: 32-104 mcg/dL 13-15 years: 30-109 mcg/dL 16-18 years: 33-102 mcg/dL 19+ years: 40-174 mcg/dL</p> <p>TRAN 200-400 mg/dL</p> <p>TIBC 0-30 days: 94-226 mcg/dL 31-182 days: 89-311 mcg/dL 183-365 days: 138-365 mcg/dL 1-3 years: 188-377 mcg/dL 4-6 years: 162-352 mcg/dL 7-9 years: 167-336 mcg/dL 10-12 years: 198-383 mcg/dL 13-15 years: 169-358 mcg/dL 16-18 years: 194-372 mcg/dL 19+ years: 290-425 mcg/dL</p> <p>IRON SATURATION 20-55%</p>	<p>IRON 0-30 days: 32-112 mcg/dL 31-365 days: 27-109 mcg/dL 1-3 years: 29-91 mcg/dL 4-6 years: 25-115 mcg/dL 7-9 years: 27-96 mcg/dL 10-12 years: 28-112 mcg/dL 13-15 years: 26-110 mcg/dL 16-18 years: 27-138 mcg/dL 19+ years: 40-174 mcg/dL</p> <p>TRAN 200-400 mg/dL</p> <p>TIBC 0-30 days: 94-222 mcg/dL 31-182 days: 116-322 mcg/dL 183-365 days: 176-384 mcg/dL 1-3 years: 204-382 mcg/dL 4-6 years: 180-390 mcg/dL 7-9 years: 183-369 mcg/dL 10-12 years: 173-356 mcg/dL 13-15 years: 193-377 mcg/dL 16-18 years: 174-351 mcg/dL 19+ years: 250-425 mcg/dL</p> <p>IRON SATURATION 20-55%</p>	<p>IRON Colorimetric</p> <p>TRAN In the procedure, the measurement of the decrease in light transmitted (increase in absorbance) through particles suspended in solution as a result of complexes formed during the antigen-antibody reaction, is the basis of this assay.</p> <p>IRON SATURATION (Iron/TIBC)*100</p>	See individual analytes	See individual analytes	See individual analytes	
Lactate Dehydrogenase	LDH	<p>0-30 days: 145-765 U/L 31-365 days: 190-820 U/L 1-3 years: 165-395 U/L 4-6 years: 135-345 U/L 7-9 years: 140-280 U/L 10-12 years: 120-260 U/L 13-15 years: 100-275 U/L 16-18 years: 105-230 U/L 19+ years: 100-190 U/L</p>	<p>0-30 days: 125-735 U/L 31-365 days: 170-840 U/L 1-3 years: 155-345 U/L 4-6 years: 155-345 U/L 7-9 years: 145-300 U/L 10-12 years: 120-325 U/L 13-15 years: 120-290 U/L 16-18 years: 105-235 U/L 19+ years: 100-190 U/L</p>	Photometric rate	Clinical Guide to Laboratory Tests, Tietz, 1995, verified by OCE/IMAC Reference Interval Study 2021. Pediatric Reference Ranges, Sodin, 1999.	25-1,200 U/L	25-60,000 U/L	
Lactate Dehydrogenase Body Fluid	FLDH		<p>Pleural: A Pleural fluid LDH to serum/plasma LDH ratio > 0.6 or a pleural fluid LDH concentration > two-thirds the upper limit of the serum/plasma LDH reference interval suggest an exudate.</p> <p>Pericardial: Pericardial fluid LDH to serum/plasma LDH ratio > 0.6 or > 300 U/L suggests an exudate.</p> <p>Peritoneal/Ascites: Peritoneal fluid LDH to serum/plasma LDH ratio > 0.6 is consistent with an exudate.</p> <p>CSF: Elevated LDH in CSF specimens may indicate a non-specific immune process. CSF LDH measurements above 40 U/L may be associated with Jakob-Creutzfeldt Disease, Bacterial Meningitis, Neurosyphilis, or tumors of the central nervous system. Contamination of red blood cells can falsely increase LDH measurements.</p> <p>Amniotic Fluid: The reference range has not been established for this fluid type. The fluid results should be compared with the concentration in serum/plasma or integrated into the clinical context for interpretation.</p>	Photometric rate	<p>CSF: Clinical Utility of Biochemical Analysis of Cerebrospinal Fluid Clinical Chemistry 1995 Watson MA.</p> <p>Pericardial and peritoneal: Burgess. Clinical Chemical Acta 343 (2004) 61-84.</p> <p>Pleural Lqhs. RW. N Engl J Med. 2002 Jun 20;346(25):1971-1977.</p>	25-1,200 U/L	25-30,000 U/L	
Lactate, Blood	LACT	0.5-1.6 mmol/L		Enzymatic colorimetric		RADIOMETER: 0.0-30.0 mmol/L	RADIOMETER: 0.0-30.0 mmol/L	
Lactate, CSF	CSLACT	0.5-2.2 mmol/L		Enzymatic colorimetric		DXC: 0.2-10.0 mmol/L	DXC: 0.2-30.0 mmol/L	
Lactate, Fluid	FLACT	<2.8 mmol/L		Enzymatic colorimetric	Beckman Coulter literature which cites Clinical Guide to Laboratory Tests, Tietz, 1995	0.2-10.0 mmol/L	0.2-30.0 mmol/L	
LDL, Direct Measure	LDL	<100 mg/dL		Enzymatic colorimetric	The reference range has not been established for this fluid specimen. The fluid results should be compared with the concentration in serum/plasma or integrated into the clinical context for interpretation.	0.2-10.0 mmol/L	0.2-30.0 mmol/L	
LH	LH	<p>0.365 days: 0.01-1.00 mIU/mL 1-10 years: 0.00-0.10 mIU/mL 11-13 years: 0.37-6.23 mIU/mL 14-18 years: 0.37-17.30 mIU/mL</p>	<p>0.365 days: <3.24 mIU/mL 1-10 years: 0.00-0.10 mIU/mL 11-13 years: 0.08-2.58 mIU/mL 16-18 years: 0.97-6.37 mIU/mL</p>	<p>Male: 19-70 years: 1.5-9.3 mIU/mL >70 years: 3.1-6.6 mIU/mL</p> <p>19+ years Female: Follicular phase: 1.9-12.5 mIU/mL Midcycle peak: 8.7-76.3 mIU/mL Luteal phase: 0.5-16.9 mIU/mL Pregnant: <0.1-1.5 mIU/mL Postmenopausal: 15.9-54.0 mIU/mL Contraceptives: 0.7-5.6 mIU/mL</p>	Two-site sandwich immunoassay chemiluminescent	Atellica IM LH Package insert 11280385. EN Rev. 04-2020-06; Pediatric Reference Ranges, Sodin, 1999	0.07-200.00 mIU/mL	0.07-6,400.00 mIU/mL
Lipase	LIPA	<p>0-30 days: 6-55 U/L 31-182 days: 4-29 U/L 183-365 days: 4-23 U/L 1-3 years: 4-31 U/L</p>		Photometric rate		6-600 U/L	6-6,000 U/L	
Lipid Panel w Calculated LDL		<p>CHOL 0-30 days: 62-155 mg/dL 31-182 days: 62-141 mg/dL 183-365 days: 76-216 mg/dL 1-3 years: 108-193 mg/dL 4-6 years: 106-193 mg/dL 7-9 years: 104-210 mg/dL 10-12 years: 105-218 mg/dL 13-15 years: 108-205 mg/dL 16-18 years: 92-234 mg/dL 19+ years: <200 mg/dL</p> <p>TRIG 0-3 years: 27-125 mg/dL 4-6 years: 32-116 mg/dL 7-9 years: 28-129 mg/dL 10-11 years: 29-140 mg/dL 12-13 years: 37-130 mg/dL 14-15 years: 38-135 mg/dL 16-19 years: 37-140 mg/dL</p>	<p>CHOL 0-30 days: 54-151 mg/dL 31-182 days: 81-147 mg/dL 183-365 days: 76-179 mg/dL 1-3 years: 85-182 mg/dL 4-6 years: 110-217 mg/dL 7-9 years: 110-211 mg/dL 10-12 years: 105-223 mg/dL 13-15 years: 91-204 mg/dL 16-18 years: 82-192 mg/dL 19+ years: <200 mg/dL</p> <p>TRIG 0-3 years: 27-125 mg/dL 4-6 years: 32-116 mg/dL 7-9 years: 28-129 mg/dL 10-11 years: 24-137 mg/dL 12-13 years: 24-145 mg/dL 14-15 years: 34-165 mg/dL 16-19 years: 34-140 mg/dL</p>	<p>CHOL Enzymatic colorimetric</p> <p>TRIG Enzymatic colorimetric</p>	See individual analytes	See individual analytes	See individual analytes	

		<p>20+ years: <150 mg/dL</p> <p>HDL 0-9 years: 35-82 mg/dL 10-13 years: 36-84 mg/dL 14-19 years: 35-65 mg/dL 20+ years: ≥40 mg/dL</p> <p>LDL CALC 0-9 years: <130 mg/dL 60-365 days: 32-117 mg/dL 1-2 years: 38-140 mg/dL 2+ years: 0-99 mg/dL</p> <p>TOT CHOL/HDL <4.5</p> <p>NON HDL <130 mg/dL</p>	<p>20+ years: <150 mg/dL</p> <p>HDL 0-9 years: 35-82 mg/dL 10-13 years: 36-84 mg/dL 14-19 years: 35-65 mg/dL 20+ years: ≥40 mg/dL</p> <p>LDL CALC 0-9 years: <130 mg/dL 60-365 days: 32-117 mg/dL 1-2 years: 38-140 mg/dL 2+ years: 0-99 mg/dL</p> <p>TOT CHOL/HDL <4.5</p> <p>NON HDL <130 mg/dL</p>		<p>HDL Enzymatic colorimetric</p> <p>LDL CALC T Cholesterol - HDL - (Trig5)</p>			
	HDLT							
Lipid Panel with Reflex to Measured LDL		<p>CHOL 0-30 days: 62-155 mg/dL 31-182 days: 62-141 mg/dL 183-365 days: 76-216 mg/dL 1-3 years: 106-193 mg/dL 4-6 years: 106-193 mg/dL 7-9 years: 104-210 mg/dL 10-12 years: 105-218 mg/dL 13-15 years: 108-205 mg/dL 16-18 years: 92-234 mg/dL 19+ years: <200 mg/dL</p> <p>TRIG 0-3 years: 27-125 mg/dL 4-6 years: 32-116 mg/dL 7-9 years: 28-129 mg/dL 10-11 years: 29-140 mg/dL 12-13 years: 37-130 mg/dL 14-15 years: 28-135 mg/dL 16-19 years: 37-140 mg/dL 20+ years: <150 mg/dL</p> <p>HDL 0-9 years: 35-82 mg/dL 10-13 years: 36-84 mg/dL 14-19 years: 35-65 mg/dL 20+ years: ≥40 mg/dL</p> <p>LDL CALC 0-9 years: <130 mg/dL 60-365 days: 32-117 mg/dL 1-2 years: 38-140 mg/dL 2+ years: 0-99 mg/dL</p> <p>TOT CHOL/HDL <4.5</p> <p>NON HDL <130 mg/dL</p>	<p>CHOL 0-30 days: 54-151 mg/dL 31-182 days: 81-147 mg/dL 183-365 days: 76-179 mg/dL 1-3 years: 85-182 mg/dL 4-6 years: 110-217 mg/dL 7-9 years: 110-211 mg/dL 10-12 years: 105-223 mg/dL 13-15 years: 91-204 mg/dL 16-18 years: 82-192 mg/dL 19+ years: <200 mg/dL</p> <p>TRIG 0-3 years: 27-125 mg/dL 4-6 years: 32-116 mg/dL 7-9 years: 28-129 mg/dL 10-11 years: 24-137 mg/dL 12-13 years: 24-145 mg/dL 14-15 years: 34-165 mg/dL 16-19 years: 34-140 mg/dL 20+ years: <150 mg/dL</p> <p>HDL 0-9 years: 35-82 mg/dL 10-13 years: 36-84 mg/dL 14-19 years: 35-65 mg/dL 20+ years: ≥40 mg/dL</p> <p>LDL CALC 0-9 years: <130 mg/dL 60-365 days: 32-117 mg/dL 1-2 years: 38-140 mg/dL 2+ years: 0-99 mg/dL</p> <p>TOT CHOL/HDL <4.5</p> <p>NON HDL <130 mg/dL</p>		<p>CHOL Enzymatic colorimetric</p> <p>TRIG Enzymatic colorimetric</p> <p>HDL Enzymatic colorimetric</p> <p>LDL CALC T Cholesterol - HDL - (Trig5)</p> <p>LDL DIRECT Enzymatic colorimetric</p>	See individual analytes	See individual analytes	See individual analytes
	LIPID							
Lipids w Direct Measure LDL		<p>CHOL 0-30 days: 62-155 mg/dL 31-182 days: 62-141 mg/dL 183-365 days: 76-216 mg/dL 1-3 years: 106-193 mg/dL 4-6 years: 106-193 mg/dL 7-9 years: 104-210 mg/dL 10-12 years: 105-218 mg/dL 13-15 years: 108-205 mg/dL 16-18 years: 92-234 mg/dL 19+ years: <200 mg/dL</p> <p>TRIG 0-3 years: 27-125 mg/dL 4-6 years: 32-116 mg/dL 7-9 years: 28-129 mg/dL 10-11 years: 29-140 mg/dL 12-13 years: 37-130 mg/dL 14-15 years: 28-135 mg/dL 16-19 years: 37-140 mg/dL 20+ years: <150 mg/dL</p> <p>HDL 0-9 years: 35-82 mg/dL 10-13 years: 36-84 mg/dL 14-19 years: 35-65 mg/dL 20+ years: ≥40 mg/dL</p> <p>LDL DIRECT <100 mg/dL</p> <p>TOT CHOL/HDL <4.5</p> <p>NON HDL <130 mg/dL</p>	<p>CHOL 0-30 days: 54-151 mg/dL 31-182 days: 81-147 mg/dL 183-365 days: 76-179 mg/dL 1-3 years: 85-182 mg/dL 4-6 years: 110-217 mg/dL 7-9 years: 110-211 mg/dL 10-12 years: 105-223 mg/dL 13-15 years: 91-204 mg/dL 16-18 years: 82-192 mg/dL 19+ years: <200 mg/dL</p> <p>TRIG 0-3 years: 27-125 mg/dL 4-6 years: 32-116 mg/dL 7-9 years: 28-129 mg/dL 10-11 years: 24-137 mg/dL 12-13 years: 24-145 mg/dL 14-15 years: 34-165 mg/dL 16-19 years: 34-140 mg/dL 20+ years: <150 mg/dL</p> <p>HDL 0-9 years: 35-82 mg/dL 10-13 years: 36-84 mg/dL 14-19 years: 35-65 mg/dL 20+ years: ≥40 mg/dL</p> <p>LDL DIRECT <100 mg/dL</p> <p>TOT CHOL/HDL <4.5</p> <p>NON HDL <130 mg/dL</p>		<p>CHOL Enzymatic colorimetric</p> <p>TRIG Enzymatic colorimetric</p> <p>HDL Enzymatic colorimetric</p> <p>LDL DIRECT Enzymatic colorimetric</p>	See individual analytes	See individual analytes	See individual analytes
	HDL.D							
Lithium Level	LI		<p>Therapeutic Range: 0-9 years: 0.60-1.20 mmol/L 60+ years: 0.40-0.80 mmol/L</p>		<p>A spectrophotometric method which can be readily adapted to automated clinical chemistry analyzers. Lithium present in the sample reacts with a substituted porphyrin compound at an alkaline pH, resulting in a change in absorbance which is directly proportional to the concentration of Lithium in the sample.</p>	<p>Applied Clinical Pharmacokinetics, Bauer, 2001 Clinical Pharmacokinetics, Ebers, 1995:29-44:50</p> <p>Bipolar Disord. 2019 Mar;21(2):117-123.</p> <p>Bipolar Disord. 2019 May;21(3):190-191.</p>	0.10-5.00 mmol/L	0.10-5.00 mmol/L
	LI							
Lytes (Na, K, Cl) - Urine - Random	ULYTR			The reference range has not been established for random urine specimens. The test result should be integrated into the clinical context for interpretation.	NAK/CL Indirect ion-selective electrode	See individual analytes	See individual analytes	See individual analytes
Lytes (Na, K, Cl) - Urine 24HR	ULYT		<p>UNA 40-220 mmol/24 hrs</p> <p>UK 25-125 mmol/24 hrs</p> <p>UCL 110-250 mmol/24 hrs</p>		NAK/CL Indirect ion-selective electrode	See individual analytes		

Lytes (Na, K, Cl) Plus CO2	LYTES	<p>NA</p> <p>0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K</p> <p>0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1+ years: 3.5-6.0 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL</p> <p>0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2</p> <p>0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>ANION GAP</p> <p>7-17 mmol/L</p>			<p>NAK/CL Indirect ion-selective electrode</p> <p>CO2 Photometric</p>	See individual analytes	See individual analytes	See individual analytes	
Lytes (Na, K, Cl Creat), Urine, Random	ULVTR UCREB			The reference range has not been established for random urine specimens. The test result should be integrated into the clinical context for interpretation.	<p>NAK/CL Indirect ion-selective electrode</p> <p>CREA Kinetic Jaffe</p>	See individual analytes	See individual analytes	See individual analytes	
Magnesium	MG	<p>0-30 days: 1.7-2.3 mg/dL 31-365 days: 1.6-2.4 mg/dL 1-3 years: 1.7-2.4 mg/dL 4-6 years: 1.7-2.2 mg/dL 7-9 years: 1.6-2.3 mg/dL 10-12 years: 1.6-2.2 mg/dL 13-15 years: 1.6-2.3 mg/dL 16-18 years: 1.5-2.2 mg/dL 19+ years: 1.6-2.6 mg/dL</p>	<p>0-30 days: 1.7-2.4 mg/dL 31-365 days: 1.6-2.4 mg/dL 1-3 years: 1.7-2.4 mg/dL 4-6 years: 1.7-2.4 mg/dL 7-9 years: 1.7-2.3 mg/dL 10-12 years: 1.6-2.2 mg/dL 13-15 years: 1.6-2.3 mg/dL 16-18 years: 1.5-2.2 mg/dL 19+ years: 1.6-2.6 mg/dL</p>		Colorimetric with xylidylblue	Clinical Guide to Laboratory Tests, Tietz, 1995, verified by OSUWAC Reference Interval Study 2021. Pediatric Reference Ranges, Soldin, 1999.	0.5-8.0 mg/dL	0.5-24.0 mg/dL	
Magnesium, 24HR Urine	UMG		72.9-121.5 mg/24 hrs		Colorimetric with xylidylblue	Clinical Guide to Laboratory Tests, Tietz, 1995			
Magnesium, Urine, Random	UMGR			The reference range has not been established for random urine specimens. The test result should be integrated into the clinical context for interpretation.	Colorimetric with xylidylblue		0.5-10.0 mg/dL	0.5-100.0 mg/dL	
Mechanical Support Hemolysis	HAPTO LDRFX	<p>HAPTO 44-215 mg/dL</p> <p>LDRFX 0-30 days: 145-765 U/L 31-365 days: 190-420 U/L 1-3 years: 165-395 U/L 4-6 years: 155-345 U/L 7-9 years: 140-280 U/L 10-12 years: 120-260 U/L 13-15 years: 100-275 U/L 16-18 years: 105-230 U/L 19+ years: 100-190 U/L</p>	<p>HAPTO 44-215 mg/dL</p> <p>LDRFX 0-30 days: 125-735 U/L 31-365 days: 170-450 U/L 1-3 years: 155-345 U/L 4-6 years: 155-345 U/L 7-9 years: 145-300 U/L 10-12 years: 120-325 U/L 13-15 years: 120-290 U/L 16-18 years: 105-235 U/L 19+ years: 100-190 U/L</p>			See individual analytes	See individual analytes	See individual analytes	
Methotrexate Level	MTXU			Due to the many different MTX protocols utilized and the many individual differences in the way this drug is handled, please contact the primary attending physician.	Homogeneous enzyme immunoassay	OSU Pharmacy	0.04-1.20 µmol/L	0.04-1,200.00 µmol/L	
Microalbumin, 24HR Urine	MALEU		<30.0 mg/24 hrs		Turbidometric	Package insert			
Microalbumin, Random Urine	MALEB				Turbidometric	See individual analytes	See individual analytes	See individual analytes	
Mononucleosis Screen	MONO			Negative	Color immunochromatographic dipstick technology with bovine erythrocyte extract coated on the membrane. In the test procedure serum is mixed with the Diluent. Then the Test Strip is placed in the mixture and the mixture migrates along the membrane.	Sure Vue Signature Mono Package Insert		Negative, Positive	
Non-Gestational 2HR Glucose Tolerance Test	GTF GTT		<p>GTF 70-125 mg/dL</p> <p>GTT 70-109 mg/dL</p>	<p>According to the American Diabetes Association's clinical practice recommendations, the diagnosis of diabetes mellitus can be made if either the fasting plasma glucose or the 2hr post-glucose samples meet or exceed 126 mg/dL or 200 mg/dL, respectively. The recommendations also note that in the absence of unequivocal hyperglycemia, the diagnosis requires two abnormal test results from the same sample or in two separate test samples.</p> <p>The ADA criteria for prediabetes is a fasting plasma glucose of 100-125 mg/dL or 2hr post-glucose sample of 140-199 mg/dL.</p>	Photometric rate with hexokinase		10-800 mg/dL	10-2,400 mg/dL	
Osmolality	OSMO		278-305 mOsm/kg		Freezing point depression	OSU In House Reference Range Validation, 2017	50-2,000 mOsm/kg	50-2,000 mOsm/kg	
Osmolality, Stool	OSMSO		275-300 mOsm/kg		Freezing point depression	Advanced Instruments Model A20, June 2014.	50-2,000 mOsm/kg	50-2,000 mOsm/kg	
Osmolality, 24 HR Urine	UOSM		300-900 mOsm/kg		Freezing point depression	Clinical Guidelines for Laboratory Tests, Tietz, 1995	50-2,000 mOsm/kg	50-2,000 mOsm/kg	
Osmolality, Urine	UOSM		300-900 mOsm/kg		Freezing point depression	Clinical Guidelines for Laboratory Tests, Tietz, 1995	50-2,000 mOsm/kg	50-2,000 mOsm/kg	
Pancreatic Fluid CEA	PFCA			Fluid CEA values >192 ng/mL may indicate mucinous cystic lesions of the pancreas. Results <192 ng/mL require clinical correlations with patient history and other imaging modalities.		Clinical Guide to Laboratory Tests, Tietz, 1995; see Source link for additional Reference Range information	2.0-100.0 ng/mL	2.0-10,000.0 ng/mL	
Pancreatic Function Test	PFUNZ PFUN15 PFUN30 PFUN45 PFUN60			Bicarbonate > 30 mmol/L is considered a normal result.	<p>NAK/CL Indirect ion-selective electrode</p> <p>CO2 Photometric</p> <p>AMY Photometric</p> <p>LIPA Photometric</p>	See individual analytes	See individual analytes	See individual analytes	
Phenobarbital Level, Random	PHNOR		Therapeutic Range: 15.0-40.0 mcg/mL			The assay is based on competition between drug in the sample and drug labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH) for antibody binding sites. Enzyme activity decreases upon binding to the antibody, so the drug concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts nicotinamide adenine dinucleotide (NAD) to NADH, resulting in an absorbance change that is measured spectrophotometrically.	Applied Clinical Pharmacokinetics, 2001	5.0-80.0 mcg/mL	5.0-240.0 mcg/mL
Phenobarbital Level, Trough (Pre Drug Level)	PHNO		Therapeutic Range: 15.0-40.0 mcg/mL			The assay is based on competition between drug in the sample and drug labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH) for antibody binding sites. Enzyme activity decreases upon binding to the antibody, so the drug concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts nicotinamide adenine dinucleotide (NAD) to NADH, resulting in an absorbance change that is measured spectrophotometrically.	Applied Clinical Pharmacokinetics, 2001	5.0-80.0 mcg/mL	5.0-240.0 mcg/mL

Phenytoin Total Level			Therapeutic Range: 10.0-20.0 mcg/mL		The assay is based on competition between drug in the sample and drug labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH) for antibody binding sites. Enzyme activity decreases upon binding to the antibody, so the drug concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts oxidized nicotinamide adenine dinucleotide (NAD) to NADH, resulting in an absorbance change that is measured spectrophotometrically.	Applied Clinical Pharmacokinetics, 2001	2.5-40.0 mcg/mL	2.5-200.0 mcg/mL
	PTN							
Phosphate, Inorganic		0-30 days: 4.3-7.7 mg/dL 31-365 days: 3.7-6.5 mg/dL 1-3 years: 3.4-6.0 mg/dL 4-6 years: 3.2-5.5 mg/dL 7-9 years: 3.1-5.5 mg/dL 10-12 years: 3.3-5.3 mg/dL 13-15 years: 2.8-4.8 mg/dL 16-18 years: 2.5-4.8 mg/dL 19+ years: 2.2-4.6 mg/dL	0-30 days: 3.9-6.9 mg/dL 31-365 days: 3.5-6.6 mg/dL 1-3 years: 3.1-6.0 mg/dL 4-6 years: 3.3-5.6 mg/dL 7-9 years: 3.0-5.4 mg/dL 10-12 years: 3.2-5.7 mg/dL 13-15 years: 2.8-5.1 mg/dL 16-18 years: 2.7-4.9 mg/dL 19+ years: 2.2-4.6 mg/dL		Photometric	OSLWMC Reference Range Study effective 12/11/2013; verified by OSLWMC Reference Interval Study 2021. Pediatric Reference Ranges, Söldn, 1999.	1.0-20.0 mg/dL	1.0-60.0 mg/dL
Phosphorus, Random, Urine	JP			The reference range has not been established for random urine specimens. The test result should be integrated into the clinical context for interpretation.	Photometric		10.0-200.0 mg/dL	10.0-1,000.0 mg/dL
Phosphorus, 24HR	LPR		0.4-1.3 g/24 hrs		Photometric	Clinical Guide to Laboratory Tests, Tietz, 1995		
Potassium			0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1-17 years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L		Indirect ion-selective electrode	OSLWMC Reference Range Study effective 12/11/2013; verified by OSLWMC Reference Interval Study 2021. Pediatric Reference Ranges, Söldn, 1999.	1.0-10.0 mmol/L	1.0-10.0 mmol/L
Potassium Body Fluid	K				Indirect ion-selective electrode			
Potassium 24 HR Urine	IK		25-125 mmol/24 hrs	Stool: The reference range has not been established for this fluid specimen. The fluid results should be compared with the concentration in serum/plasma or integrated into the clinical context for interpretation.	Indirect ion-selective electrode	Clinical Guide to Laboratory Tests, Tietz, 1995		
Potassium, Random Urine	LKB			The reference range has not been established for random urine specimens. The test result should be integrated into the clinical context for interpretation.	Indirect ion-selective electrode	Clinical Guide to Laboratory Tests, Tietz, 1995	2.0-200.0 mmol/L	2.0-200.0 mmol/L
Prealbumin	PALB		17-34 mg/dL		Turbidimetry	Package Insert. Verified by OSLWMC Reference Interval Study 2021	3-80 mg/dL	3-1,600 mg/dL
Procalcitonin	PROCAL		<0.50 ng/mL		Two-site sandwich immunoassay chemiluminescent.	Aliccia IM Procalcitonin 11200767_EN Rev. 03, 2019-06	0.04-50.00 ng/mL	0.04-2,000.00 ng/mL
Progesterone	PROG			Male: 0.28-1.22 ng/mL Female: Follicular phase: 0.1-40 ng/mL Luteal phase: 3.34-25.50 ng/mL Midluteal phase: 4.44-28.03 ng/mL Postmenopausal: 0-0.71 ng/mL	Competitive immunoassay using direct chemiluminescent technology.	Aliccia IM Progesterone Package Insert 11200386_EN Rev. 04-2020-06	0.21-60.00 ng/mL	0.21-3,000.00 ng/mL
Prolactin	PROL			Male: 2.1-17.7 ng/mL Female: Nursing infant: 2.8-29.2 ng/mL Pregnant: 9.7-208.5 ng/mL Postmenopausal: 1.8-20.3 ng/mL <2 years: 3.3-14.7 ng/mL 2-5 years: 1.0-12.8 ng/mL 6-10 years: 1.2-11.4 ng/mL 11-17 years: 1.4-14.3 ng/mL	Two-site sandwich immunoassay chemiluminescent.	Abnii Contour Prolactin Package Insert 111746 Rev. N, 2008-09; Pediatric Reference Intervals, 5th ed Söldn, 2005	0.3-200.0 ng/mL	0.3-800,000.0 ng/mL
Protein & Glucose, CSF	CFPG CFP		GLUC/CSF 40-70 mg/dL CFP 0-3 days: 40-120 mg/dL 4-30 days: 20-80 mg/dL 31+ days: 15-45 mg/dL		GLUC/CSF Photometric rate with hexokinase CFP Colorimetric with Pyrogallol red	See individual analytes	See individual analytes	See individual analytes
Protein Total	TP	0-30 days: 4.2-6.2 g/dL 31-182 days: 4.4-6.6 g/dL 183-365 days: 5.6-7.9 g/dL 1-18 years: 5.7-8.0 g/dL 19+ years: 6.4-8.3 g/dL	0-30 days: 4.1-6.3 g/dL 31-182 days: 4.7-6.7 g/dL 183-365 days: 5.5-7.9 g/dL 1-18 years: 5.7-8.0 g/dL 19+ years: 6.4-8.3 g/dL		Colorimetric with cupric ions in an alkaline solution	Clinical Guide to Laboratory Tests, Tietz, 1995; verified by OSLWMC Reference Interval Study 2021. Pediatric Reference Ranges, Söldn, 1999	3.0-12.0 g/dL	3.0-24.0 g/dL
Protein, CSF	CFP		0-3 days: 40-120 mg/dL 4-30 days: 20-80 mg/dL		Colorimetric with Pyrogallol red	Clinical Guide to Laboratory Tests, Tietz, 1995; Pediatric Reference Ranges, Söldn, 1999	4-200 mg/dL	4-5,000 mg/dL
Protein, Fluid	FLP			Pleural: Pleural fluid protein to serum/plasma protein ratio > 0.5 are consistent with exudates. Pericardial: The reference range has not been established for this fluid specimen. The fluid results should be compared with the concentration in serum/plasma or integrated into the clinical context for interpretation. Peritoneal/Ascites: The reference range has not been established for this fluid specimen. The fluid results should be compared with the concentration in serum/plasma or integrated into the clinical context for interpretation.	Colorimetric with cupric ions in an alkaline solution	4th Edition of Tietz	0.5-12.0 g/dL	0.5-24.0 g/dL
Protein, 24 HR Urine	LPRO		40-225 mg/24 hrs		Colorimetric with Pyrogallol red	Clinical Guide to Laboratory Tests, Tietz, 1995		
Protein, Random Urine	LPROR			The reference range has not been established for random urine specimens. The test result should be integrated into the clinical context for interpretation.	Colorimetric with Pyrogallol red		4-200 mg/dL	4-5,000 mg/dL
PSA - Diagnostic/Tumor Marker	PSATM		≤400 ng/mL		Two-site sandwich immunoassay chemiluminescent.	Aliccia IM PSA Package Insert 10997799_EN Rev. 03-2019-09	0.04-100.00 ng/mL	0.04-4,000,000.00 ng/mL
PSA, Reflex to Free and Total PSA	PSAR		≤400 ng/mL			Aliccia IM PSA Package Insert 10997799_EN Rev. 03-2019-09	0.04-100.00 ng/mL	0.04-4,000,000.00 ng/mL
PSA, Screening	PSA		≤400 ng/mL		Two-site sandwich immunoassay chemiluminescent.	Aliccia IM PSA Package Insert 10997799_EN Rev. 03-2019-09	0.04-100.00 ng/mL	0.04-4,000,000.00 ng/mL

Renal Panel	RENAL	<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1+ years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>GLUC 0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL</p> <p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-0.90 mg/dL 31-365 days: 0.40-0.60 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.50-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.70-1.10 mg/dL 16-18 years: 0.80-1.20 mg/dL 19+ years: 0.50-1.20 mg/dL</p> <p>CA 0-30 days: 8.4-10.6 mg/dL 31-365 days: 8.9-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL</p> <p>OSMO 278-305 mOsm/kg</p> <p>ANION GAP 7-17 mmol/L</p> <p>eGFR ≥60 mL/min/1.73m²</p> <p>ALB 0-30 days: 2.7-4.3 g/dL 31-182 days: 2.9-4.2 g/dL 183-365 days: 3.3-4.8 g/dL 1-19 years: 2.9-4.2 g/dL 19+ years: 3.5-5.0 g/dL</p> <p>IP 0-30 days: 4.3-7.7 mg/dL 31-365 days: 3.7-6.5 mg/dL 1-3 years: 3.4-6.0 mg/dL 4-6 years: 3.2-5.5 mg/dL 7-9 years: 3.1-5.5 mg/dL 10-12 years: 3.3-5.3 mg/dL 13-15 years: 2.8-4.8 mg/dL 16-18 years: 2.5-4.8 mg/dL 19+ years: 2.2-4.6 mg/dL</p>	<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1+ years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p> <p>CO2 0-2 years: 13-29 mmol/L 3+ years: 21-31 mmol/L</p> <p>GLUC 0-30 days: 55-115 mg/dL 31-182 days: 57-117 mg/dL 183-365 days: 70-126 mg/dL 1+ years: 70-99 mg/dL</p> <p>BUN 7-25 mg/dL</p> <p>CREA 0-30 days: 0.50-1.20 mg/dL 31-365 days: 0.40-0.70 mg/dL 1-3 years: 0.40-0.70 mg/dL 4-6 years: 0.50-0.80 mg/dL 7-9 years: 0.60-0.90 mg/dL 10-12 years: 0.60-1.00 mg/dL 13-15 years: 0.60-1.20 mg/dL 16-18 years: 0.80-1.40 mg/dL 19+ years: 0.70-1.30 mg/dL</p> <p>CA 0-30 days: 8.5-10.6 mg/dL 31-365 days: 8.7-10.5 mg/dL 1+ years: 8.6-10.5 mg/dL</p> <p>OSMO 278-305 mOsm/kg</p> <p>ANION GAP 7-17 mmol/L</p> <p>eGFR ≥60 mL/min/1.73m²</p> <p>ALB 0-30 days: 2.6-4.1 g/dL 31-182 days: 2.8-4.9 g/dL 183-365 days: 2.8-4.8 g/dL 1-19 years: 3.2-4.7 g/dL 19+ years: 3.5-5.0 g/dL</p> <p>IP 0-30 days: 3.9-6.9 mg/dL 31-365 days: 3.5-6.6 mg/dL 1-3 years: 3.1-6.0 mg/dL 4-6 years: 3.3-6.0 mg/dL 7-9 years: 3.0-5.4 mg/dL 10-12 years: 3.2-5.7 mg/dL 13-15 years: 2.8-5.1 mg/dL 16-18 years: 2.7-4.9 mg/dL 19+ years: 2.2-4.6 mg/dL</p>				
Rheumatoid Factor	RF		≤14 IU/mL	Turbidimetry	Package Insert. Verified by OSUWMC Reference Interval Study 2021.	10-120 IU/mL	10-3,000 IU/mL
Salicylate Level	SAL		Therapeutic Range: 20.0-30.0 mg/dL	Serum is mixed with Reagent 1, which contains antibodies to salicylic acid and the coenzyme nicotinamide adenine dinucleotide (NAD). Subsequently, Reagent 2, which contains salicylic acid labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH), is added. Salicylic acid in the sample and salicylic acid-labeled G6PDH compete for antibody binding sites. Enzyme activity decreases upon binding to the antibody, so the salicylic acid concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts oxidized NAD to NADH, resulting in an absorbance change that is measured spectrophotometrically.	Applied Pharmacokinetics: Principles of Therapeutic Drug Monitoring, 2nd Edition 2002 Applied Therapeutics, Inc. and Micromedex. On OSU Intranet	5.0-80.0 mg/dL	5.0-240.0 mg/dL
Sodium	NA		0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L	Indirect ion-selective electrode	Verified by OSUWMC Reference Interval Study 2021.	50-200 mmol/L	50-200 mmol/L
Sodium Body Fluid	FNA			Indirect ion-selective electrode		50-200 mmol/L	50-200 mmol/L
Sodium, Potassium, Chloride	NA, K, CL		<p>NA 0-6 days: 133-146 mmol/L 7-30 days: 134-144 mmol/L 31-182 days: 134-142 mmol/L 183-365 days: 133-142 mmol/L 1+ years: 135-145 mmol/L</p> <p>K 0-7 days: 3.2-5.5 mmol/L 8-30 days: 3.4-6.0 mmol/L 31-182 days: 3.5-5.6 mmol/L 183-365 days: 3.5-6.1 mmol/L 1+ years: 3.3-4.6 mmol/L 18+ years: 3.5-5.0 mmol/L</p> <p>CL 0-365 days: 98-113 mmol/L 1-17 years: 102-112 mmol/L 18+ years: 98-108 mmol/L</p>	Indirect ion-selective electrode	See individual analytes	See individual analytes	See individual analytes
Sodium, 24 HR Urine	UNA		40-220 mmol/24 hrs	Indirect ion-selective electrode	Clinical Guide to Laboratory Tests, Tietz, 1995		

Sodium, Random Urine	LNAR			The reference range has not been established for random urine specimens. The test result should be integrated into the clinical context for interpretation.	Indirect ion-selective electrode		10-400 mmol/L	10-400 mmol/L
T3 Free	FT3	0-3 days: 1.4-5.4 pg/mL 4-29 days: 1.5-5.0 pg/mL 30-365 days: 2.5-5.5 pg/mL 1-5 years: 3.0-6.0 pg/mL 6-10 years: 2.7-6.2 pg/mL 11-15 years: 2.6-5.7 pg/mL 16-18 years: 2.8-5.2 pg/mL 19+ years: 2.3-4.2 pg/mL	0-3 days: 1.4-4.8 pg/mL 4-29 days: 1.4-5.5 pg/mL 30-365 days: 2.0-6.0 pg/mL 1-5 years: 2.4-6.7 pg/mL 6-10 years: 2.9-6.0 pg/mL 11-15 years: 3.1-5.9 pg/mL 16-18 years: 3.5-5.7 pg/mL 19+ years: 2.3-4.2 pg/mL		Competitive immunoassay using direct chemiluminescent technology.	Atellica IM Free T3 Package Insert 10995347_EN Rev. 03-2020-06	0.2-20.0 pg/mL	0.2-20.0 pg/mL
T3 Total (Triiodothyronine)	T3RIA	0-29 days: 0.4-1.77 ng/mL 30-365 days: 0.73-2.21 ng/mL 1-5 years: 1.26-2.16 ng/mL 6-10 years: 1.01-1.95 ng/mL 11-15 years: 1.04-1.84 ng/mL 16-18 years: 1.01-1.51 ng/mL 19+ years: 0.60-1.31 ng/mL	0-29 days: 0.51-1.84 ng/mL 30-365 days: 1.03-2.29 ng/mL 1-5 years: 0.93-2.13 ng/mL 6-10 years: 1.04-1.96 ng/mL 11-15 years: 0.88-1.76 ng/mL 16-18 years: 0.86-1.78 ng/mL 19+ years: 0.63-1.31 ng/mL		Competitive immunoassay using direct chemiluminescent technology.	Atellica IM Total T3 Package Insert 10995424_EN Rev. 03-2020-06	0.10-8.00 ng/mL	0.10-8.00 ng/mL
T4	T4	0-29 days: 3.4-11.8 mcg/dL 30-365 days: 4.7-11.8 mcg/dL 1-5 years: 6.0-11.3 mcg/dL 6-10 years: 5.2-9.7 mcg/dL 11-15 years: 4.9-9.0 mcg/dL 16-18 years: 5.1-9.0 mcg/dL 19+ years: 4.5-10.9 mcg/dL	0-29 days: 3.4-12.6 mcg/dL 30-365 days: 5.4-14.1 mcg/dL 1-5 years: 5.3-10.2 mcg/dL 6-10 years: 5.3-9.5 mcg/dL 11-15 years: 4.6-9.2 mcg/dL 16-18 years: 4.9-8.1 mcg/dL 19+ years: 4.5-10.9 mcg/dL		Competitive immunoassay using direct chemiluminescent technology.	Atellica IM Total T4 Package Insert 10995425_EN Rev. 03-2020-06; Pediatric Reference Ranges, Sodin, 1999	0.4-30.0 mcg/dL	0.4-300.0 mcg/dL
T4 Free	FT4		0-18 years: 1.06-1.64 ng/dL 19+ years: 0.89-1.76 ng/dL		Competitive immunoassay using direct chemiluminescent technology.	Atellica IM Free T4 Package Insert 10995348_EN Rev. 06-2020-11; Pediatric Reference Ranges, Sodin, 1999	0.10-12.00 ng/dL	0.10-12.00 ng/dL
Testosterone	TESTOS	0-6 months: 20-80 ng/dL 6 months to 9 years: <20 ng/dL 10-11 years: <44 ng/dL 12-16 years: <75 ng/dL 17-18 years: 20-75 ng/dL 19+ years: 8-60 ng/dL	0-6 months: 75-400 ng/dL 6 months to 9 years: <20 ng/dL 10-11 years: <130 ng/dL 12-13 years: <800 ng/dL 14-15 years: <1,200 ng/dL 16-18 years: 100-1,200 ng/dL 17-18 years: 300-1,200 ng/dL 19+ years: 240-950 ng/dL		Competitive immunoassay using direct chemiluminescent technology.	Siemens Atellica IM Reference Interval Verification Study 2023	7-1,500 ng/dL	7-3,000 ng/dL
Testosterone Total and Free, Includes SHBG	TESTOS SHBG	TESTOS 0-6 months: 20-80 ng/dL 6 months to 9 years: <20 ng/dL 10-11 years: <44 ng/dL 12-16 years: <75 ng/dL 17-18 years: 20-75 ng/dL 19+ years: 8-60 ng/dL SHBG 18.00-144.00 nmol/L FREE TESTOS 0-365 days: <0.25 ng/dL 1-6 years: <0.13 ng/dL 7-8 years: <0.23 ng/dL 8-9 years: <0.14 ng/dL 9-10 years: <0.46 ng/dL 10-11 years: <0.59 ng/dL 11-12 years: <0.74 ng/dL 12-13 years: <0.84 ng/dL 13-14 years: <0.96 ng/dL 14-15 years: <1.06 ng/dL 15-18 years: <1.09 ng/dL 19+ years: 0.60-1.08 ng/dL	TESTOS 0-6 months: 75-400 ng/dL 6 months to 9 years: <20 ng/dL 10-11 years: <130 ng/dL 12-13 years: <800 ng/dL 14-15 years: <1,200 ng/dL 15-16 years: 100-1,200 ng/dL 17-18 years: 300-1,200 ng/dL 19+ years: 240-950 ng/dL SHBG 10.00-57.00 nmol/L FREE TESTOS 0-365 days: 0.20-3.10 ng/dL 1-8 years: <0.13 ng/dL 9-10 years: <0.45 ng/dL 10-11 years: <1.26 ng/dL 11-12 years: <0.52 ng/dL 12-13 years: <0.28 ng/dL 13-14 years: <0.60 ng/dL 14-15 years: 0.48-1.30 ng/dL 15-16 years: 1.62-1.70 ng/dL 16-17 years: 2.09-1.90 ng/dL 17-18 years: 4.28-20.90 ng/dL 18-19 years: 5.40-21.80 ng/dL 19+ years: 2.28-20.70 ng/dL		See individual analytes	See individual analytes	See individual analytes	See individual analytes
Theophylline Level	THEO		Therapeutic Range: 5.0-20.0 mcg/mL		Based on competition between drug in the sample and drug labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH) for antibody binding sites. Enzyme activity decreases upon binding to the antibody, so the drug concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts oxidized nicotinamide adenine dinucleotide (NAD) to NADH, resulting in an absorbance change that is measured spectrophotometrically.	Applied Clinical Pharmacokinetics, 2001	2.5-40.0 mcg/mL	2.5-200.0 mcg/mL
Tobramycin Level, Extended Interval	TOBREI		Peak: 10.0 - 15.0 mcg/mL, Trough: <1.0 mcg/mL		This assay is based on competition for antibody binding sites between drug in the sample and drug labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH). Enzyme activity decreases upon binding to the antibody, so the drug concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts oxidized nicotinamide adenine dinucleotide (NAD) to NADH, resulting in an absorbance change that is measured spectrophotometrically.	Antimicrobial Stewardship Program, 2013	0.6-10.0 mcg/mL	0.6-50.0 mcg/mL
Tobramycin Level, Peak (Post Drug Level)	TOBRPK		Therapeutic Range: 10.0-15.0 mcg/mL		This assay is based on competition for antibody binding sites between drug in the sample and drug labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH). Enzyme activity decreases upon binding to the antibody, so the drug concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts oxidized nicotinamide adenine dinucleotide (NAD) to NADH, resulting in an absorbance change that is measured spectrophotometrically.	Antimicrobial Stewardship Program, 2013	0.6-10.0 mcg/mL	0.6-50.0 mcg/mL
Tobramycin Level, Random	TOBR		Peak: 10.0 - 15.0 mcg/mL, Trough: <1.0 mcg/mL		This assay is based on competition for antibody binding sites between drug in the sample and drug labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH). Enzyme activity decreases upon binding to the antibody, so the drug concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts oxidized nicotinamide adenine dinucleotide (NAD) to NADH, resulting in an absorbance change that is measured spectrophotometrically.	Antimicrobial Stewardship Program, 2013	0.6-10.0 mcg/mL	0.6-50.0 mcg/mL
Tobramycin Level, Trough (Pre Drug Level)	TOBTR		Therapeutic Range: <1.0 mcg/mL		This assay is based on competition for antibody binding sites between drug in the sample and drug labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH). Enzyme activity decreases upon binding to the antibody, so the drug concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts oxidized nicotinamide adenine dinucleotide (NAD) to NADH, resulting in an absorbance change that is measured spectrophotometrically.	Antimicrobial Stewardship Program, 2013	0.6-10.0 mcg/mL	0.6-50.0 mcg/mL

Transferrin/Inon Binding		<p>TRAN 200-400 mg/dL</p> <p>TIBC 0-30 days: 94-236 mg/dL 31-182 days: 99-311 mg/dL 183-365 days: 138-365 mg/dL 1-3 years: 184-377 mg/dL 4-6 years: 162-352 mg/dL 7-9 years: 167-336 mg/dL 10-12 years: 198-383 mg/dL 13-15 years: 169-358 mg/dL 16-18 years: 194-372 mg/dL 19+ years: 250-425 mg/dL</p>	<p>TRAN 200-400 mg/dL</p> <p>TIBC 0-30 days: 94-232 mg/dL 31-182 days: 116-322 mg/dL 183-365 days: 176-384 mg/dL 1-3 years: 204-382 mg/dL 4-6 years: 180-390 mg/dL 7-9 years: 183-369 mg/dL 10-12 years: 173-356 mg/dL 13-15 years: 193-377 mg/dL 16-18 years: 174-351 mg/dL 19+ years: 250-425 mg/dL</p>			Clinical Guide to Laboratory Tests, Tietz, 1995; Pediatric Reference Ranges, Sodin, 1999	See individual analytes	See individual analytes	
Triglyceride	TRANB	<p>0-3 years: 27-125 mg/dL 4-6 years: 32-116 mg/dL 7-9 years: 28-129 mg/dL 10-11 years: 39-140 mg/dL 12-13 years: 27-130 mg/dL 14-15 years: 38-135 mg/dL 16-19 years: 37-140 mg/dL 20+ years: <150 mg/dL</p>	<p>0-3 years: 27-125 mg/dL 4-6 years: 32-116 mg/dL 7-9 years: 28-129 mg/dL 10-11 years: 24-137 mg/dL 12-13 years: 24-145 mg/dL 14-15 years: 34-165 mg/dL 16-19 years: 34-140 mg/dL 20+ years: <150 mg/dL</p>	Enzymatic colorimetric		National Cholesterol Education Project (NCEP) Adult Treatment Protocol (ATP-III) (Circulation, 2002;106:3143-3421)	10-1,000 mg/dL	10-10,000 mg/dL	
Triglyceride Body Fluid	TRBG			<p>Pleural: Pleural triglycerides <50 mg/dL, excludes a chylothorax. Pleural triglycerides >110mg/dL, supports a diagnosis of a chylothorax.</p> <p>Peritoneal/Ascites: Peritoneal fluid triglyceride values greater than 110 mg/dL, have been suggested for diagnosis of chylothorax. Measurement may also be useful in distinguishing cirrhotic versus malignant origins.</p> <p>Drainage: The reference range has not been established for this fluid specimen. The fluid results should be compared with the concentration in serum/plasma or integrated into the clinical context for interpretation.</p>	Enzymatic colorimetric	Pleural: Staats BA, et al. Mayo Clin Proc. 1980;55(11):700. Peritoneal: Jung D, et al. Hepatology. 1986;6(2):239.	10-1,000 mg/dL	10-10,000 mg/dL	
TSH	FTTBC		<p>0-18 years: 0.652-4.409 uIU/mL 19+ years: 0.550-4.780 uIU/mL</p>			The Atellica IM TSH-UL assay is a third-generation assay that employs anti-FITC monoclonal antibody covalently bound to paramagnetic particles, an FITC-labeled anti-TSH capture mouse monoclonal antibody, and a tracer consisting of a proprietary scintidium ester and an anti-TSH mouse monoclonal antibody conjugated to bovine serum albumin (BSA) for chemiluminescent detection.	Atellica IM TSH-UL Package Insert 11202198. EN Rev. 04-2021-03.	0.008-150.000 uIU/mL	0.008-150.000 uIU/mL
TSH w/ FT4 Reflex	TSH		<p>0-18 years: 0.652-4.409 uIU/mL 19+ years: 0.550-4.780 uIU/mL</p>			The Atellica IM TSH-UL assay is a third-generation assay that employs anti-FITC monoclonal antibody covalently bound to paramagnetic particles, an FITC-labeled anti-TSH capture mouse monoclonal antibody, and a tracer consisting of a proprietary scintidium ester and an anti-TSH mouse monoclonal antibody conjugated to bovine serum albumin (BSA) for chemiluminescent detection.	See individual analytes	See individual analytes	See individual analytes
Urea Nitrogen, 24 HR Urine	TSHOR		10.0-20.0 g/24 hrs		Photometric rate	Clinical Guide to Laboratory Tests, Tietz, 1995			
Uric Acid	URIFA	<p>0-30 days: 1.0-4.6 mg/dL 31-365 days: 1.1-5.4 mg/dL 1-3 years: 1.8-5.0 mg/dL 4-6 years: 2.0-5.1 mg/dL 7-9 years: 1.8-5.5 mg/dL 10-12 years: 2.5-5.9 mg/dL 13-15 years: 2.2-6.4 mg/dL 16-18 years: 2.4-6.6 mg/dL 19+ years: 2.8-6.0 mg/dL</p>	<p>0-30 days: 1.2-3.9 mg/dL 31-365 days: 1.2-5.6 mg/dL 1-3 years: 2.1-5.6 mg/dL 4-6 years: 1.8-5.5 mg/dL 7-9 years: 1.8-5.4 mg/dL 10-12 years: 2.2-5.8 mg/dL 13-15 years: 3.1-7.0 mg/dL 16-18 years: 2.1-7.6 mg/dL 19+ years: 3.5-7.0 mg/dL</p>	Enzymatic colorimetric		OSLW/MC Reference Range Study effective 12.11.2013; verified by OSLW/MC Reference Interval Study 2021. Pediatric Reference Ranges, Sodin, 1999.	1.5-30.0 mg/dL	1.5-60.0 mg/dL	
Uric Acid (Special Handling)	URIC	<p>0-30 days: 1.0-4.6 mg/dL 31-365 days: 1.1-5.4 mg/dL 1-3 years: 1.8-5.0 mg/dL 4-6 years: 2.0-5.1 mg/dL 7-9 years: 1.8-5.5 mg/dL 10-12 years: 2.5-5.9 mg/dL 13-15 years: 2.2-6.4 mg/dL 16-18 years: 2.4-6.6 mg/dL 19+ years: 2.8-6.0 mg/dL</p>	<p>0-30 days: 1.2-3.9 mg/dL 31-365 days: 1.2-5.6 mg/dL 1-3 years: 2.1-5.6 mg/dL 4-6 years: 1.8-5.5 mg/dL 7-9 years: 1.8-5.4 mg/dL 10-12 years: 2.2-5.8 mg/dL 13-15 years: 3.1-7.0 mg/dL 16-18 years: 2.1-7.6 mg/dL 19+ years: 3.5-7.0 mg/dL</p>	Enzymatic colorimetric		OSLW/MC Reference Range Study effective 12.11.2013; Pediatric Reference Ranges, Sodin, 1999.	1.5-30.0 mg/dL	1.5-60.0 mg/dL	
Uric Acid, 24HR	URICSH		0.3-0.8 g/24 hrs		Enzymatic colorimetric	Clinical Guide to Laboratory Tests, Tietz, 1995			
Uric Acid, Random, Urine	URICR			The reference range has not been established for random urine specimens. The test result should be integrated into the clinical context for interpretation.	Enzymatic colorimetric		1.0-100.0 mg/dL	1.0-300.0 mg/dL	
Urine Protein/Creat Ratio, Random	UPCR				<p>UCRE Kinetic Jaffe</p> <p>UPRO Colorimetric with Pyrogallol red</p>	See individual analytes	See individual analytes	See individual analytes	
Urine Protein/Creat Ratio, 24HR	UPCR24	<p>UCRE 0-2 years: 0.70-2.00 g/24 hrs 3-8 years: 0.11-0.68 g/24 hrs 9-12 years: 0.17-1.41 g/24 hrs 13-17 years: 0.29-1.87 g/24 hrs 18+ years: 0.60-1.00 g/24 hrs</p> <p>UPRO 60-225 mg/24 hrs</p>	<p>UCRE 0-2 years: 0.70-2.00 g/24 hrs 3-8 years: 0.11-0.68 g/24 hrs 9-12 years: 0.17-1.41 g/24 hrs 13-17 years: 0.29-1.87 g/24 hrs 18+ years: 0.80-2.00 g/24 hrs</p> <p>UPRO 60-225 mg/24 hrs</p>			<p>UCRE Kinetic Jaffe</p> <p>UPRO Colorimetric with Pyrogallol red</p>	See individual analytes	See individual analytes	See individual analytes
Urine Urea Nitrogen - Random	URUAR			The reference range has not been established for random urine specimens. The test result should be integrated into the clinical context for interpretation.	Photometric rate	Clinical Guide to Laboratory Tests, Tietz, 1995	20-1,300 mg/dL	20-13,000 mg/dL	
Vancomycin Level, Continuous Infusion	VANCI		20.0-25.0 mcg/mL			Serum or plasma is mixed with Reagent 1, which contains vancomycin labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH). Subsequently, Reagent 2, which contains antibodies to vancomycin and the coenzyme nicotinamide adenine dinucleotide (NAD), is added. Vancomycin in the sample and vancomycin-labeled G6PDH compete for antibody binding sites. Enzyme activity decreases upon binding to the antibody, so the vancomycin concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts oxidized NAD to NADH, resulting in an absorbance change that is measured spectrophotometrically.	American Journal of Health-System Pharmacy, Volume 77, Issue 11, 1 June 2020, Pages 835-864	2.0-50.0 mcg/mL	2.0-250.0 mcg/mL
Vancomycin Level, Random	VANCO		Peak: 20.0-40.0 mcg/mL Trough: 10.0-20.0 mcg/mL			Serum or plasma is mixed with Reagent 1, which contains vancomycin labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH). Subsequently, Reagent 2, which contains antibodies to vancomycin and the coenzyme nicotinamide adenine dinucleotide (NAD), is added. Vancomycin in the sample and vancomycin-labeled G6PDH compete for antibody binding sites. Enzyme activity decreases upon binding to the antibody, so the vancomycin concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts oxidized NAD to NADH, resulting in an absorbance change that is measured spectrophotometrically.	Applied Clinical Pharmacokinetics, 2001: Clinical Pharmacotherapy, 1995; 15:85-91	2.0-50.0 mcg/mL	2.0-250.0 mcg/mL

Vancomycin Level, Trough (Pre Drug Level)	VANCTR	Therapeutic Range: 10.0-20.0 mcg/mL	Serum or plasma is mixed with Reagent 1, which contains vancomycin labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH). Subsequently, Reagent 2, which contains antibodies to vancomycin and the coenzyme nicotinamide adenine dinucleotide (NAD), is added. Vancomycin in the sample and vancomycin-labeled G6PDH compete for antibody binding sites. Enzyme activity decreases upon binding to the antibody, so the vancomycin concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts oxidized NAD to NADH, resulting in an absorbance change that is measured spectrophotometrically.	Applied Clinical Pharmacokinetics, 2001. Clinical Pharmacotherapy, 1995; 15:85-91	2.0-50.0 mcg/mL	2.0-250.0 mcg/mL
Vancomycin, Peak (Post Drug Level)	VANCPK	Therapeutic Range: 20.0-40.0 mcg/mL	Serum or plasma is mixed with Reagent 1, which contains vancomycin labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH). Subsequently, Reagent 2, which contains antibodies to vancomycin and the coenzyme nicotinamide adenine dinucleotide (NAD), is added. Vancomycin in the sample and vancomycin-labeled G6PDH compete for antibody binding sites. Enzyme activity decreases upon binding to the antibody, so the vancomycin concentration in the sample can be measured in terms of enzyme activity. Active enzyme converts oxidized NAD to NADH, resulting in an absorbance change that is measured spectrophotometrically.	Applied Clinical Pharmacokinetics, 2001. Clinical Pharmacotherapy, 1995; 15:85-91	2.0-50.0 mcg/mL	2.0-250.0 mcg/mL
Vitamin B12	B12B	0-4 years: <213 pg/mL 5-9 years: 343-1,184 pg/mL 10-14 years: 298-955 pg/mL	Competitive immunoassay using direct chemiluminescent technology	Archives IM Vitamin B12 Package Insert 10995437, EN Rev. 02-2019-08; Pediatric Reference Ranges, Soldin, 1999	45-2,000 pg/mL	45-20,000 pg/mL
Acute Myeloid Leukemia (AML) (ASP, BX, FE)	PLACANTH IDMANP		Manual			
CBC, EDIF, Platelet	CBC, EDIF	See HEM3 and INPT209	The Sysmex XN performs hematology analysis according to the hydrodynamic focusing (DC detection), flow cytometry method (semiconductor laser) and sodium lauryl sulfate (SLS)-hemoglobin method.	OSU Internal Normal Range Study, October 2018 Soldin, Steven J. <i>Pediatric Reference Intervals</i> , 7th ed., AACCPress, 2011.	See HEM3 and INPT209	See HEM3 and INPT209
CBC, Platelets		<p>WBC</p> <p>0-14 days: 8.16-14.56 K/uL 15-30 days: 8.36-14.42 K/uL 31-60 days: 7.05-14.08 K/uL 61-179 days: 6.00-13.25 K/uL 180 days-2 years: 6.48-13.02 K/uL 2-5 years: 4.86-13.18 K/uL 6-11 years: 4.27-11.40 K/uL 12-17 years: 4.19-9.43 K/uL 18+ years: 3.99-11.19 K/uL</p> <p>RBC</p> <p>0-14 days: 4.12-5.74 M/uL 15-30 days: 3.32-4.80 M/uL 31-60 days: 2.93-3.87 M/uL 61-179 days: 3.45-4.75 M/uL 180 days-2 years: 3.97-5.01 M/uL 2-5 years: 3.84-4.92 M/uL 6-11 years: 3.90-4.96 M/uL 12-17 years: 3.93-4.90 M/uL 18+ years: 3.91-5.04 M/uL</p> <p>HGB</p> <p>0-7 days: 13.4-20.0 g/dL 8-14 days: 13.4-20.0 g/dL 15-30 days: 10.8-14.6 g/dL 31-60 days: 9.2-11.4 g/dL 61-179 days: 9.9-12.4 g/dL 180 days-2 years: 10.2-12.7 g/dL 2-5 years: 10.2-12.7 g/dL 6-11 years: 10.6-14.2 g/dL 12-17 years: 10.8-13.3 g/dL 18+ years: 11.4-15.2 g/dL</p> <p>HCT</p> <p>0-14 days: 39.6-57.2 % 15-30 days: 32.0-44.5 % 31-60 days: 27.7-35.1 % 61-179 days: 28.5-37.1 % 180 days-2 years: 30.9-37.9 % 2-5 years: 31.2-37.8 % 6-11 years: 32.4-39.5 % 12-17 years: 33.4-40.4 % 18+ years: 34.9-44.3 %</p> <p>MCV</p> <p>0-14 days: 92.7-106.4 fL 15-30 days: 90.1-103.0 fL 31-60 days: 83.4-96.4 fL 61-179 days: 74.8-88.3 fL 180 days-2 years: 71.3-82.6 fL 2-5 years: 72.3-85.0 fL 6-11 years: 75.8-87.6 fL 12-17 years: 76.9-90.6 fL 18+ years: 79.6-97.7 fL</p>	<p>WBC</p> <p>0-14 days: 8.04-15.40 K/uL 15-30 days: 7.80-15.91 K/uL 31-60 days: 8.14-14.99 K/uL 61-179 days: 6.51-13.32 K/uL 180 days-2 years: 5.98-13.51 K/uL 2-5 years: 5.14-13.38 K/uL 6-11 years: 4.31-11.00 K/uL 12-17 years: 3.84-9.84 K/uL 18+ years: 3.73-10.10 K/uL</p> <p>RBC</p> <p>0-14 days: 4.16-5.55 M/uL 15-30 days: 3.16-4.43 M/uL 31-60 days: 3.02-4.22 M/uL 61-179 days: 3.43-4.80 M/uL 180 days-2 years: 4.03-5.07 M/uL 2-5 years: 3.89-4.97 M/uL 6-11 years: 3.96-5.03 M/uL 12-17 years: 4.03-5.29 M/uL 18+ years: 4.38-5.83 M/uL</p> <p>HGB</p> <p>0-7 days: 13.9-19.1 g/dL 8-14 days: 13.9-19.1 g/dL 15-30 days: 10.0-15.3 g/dL 31-60 days: 8.9-12.7 g/dL 61-179 days: 9.6-12.4 g/dL 180 days-2 years: 10.1-12.5 g/dL 2-5 years: 10.2-12.7 g/dL 6-11 years: 10.7-13.4 g/dL 12-17 years: 11.0-14.5 g/dL 18+ years: 13.4-16.8 g/dL</p> <p>HCT</p> <p>0-14 days: 39.8-53.4 % 15-30 days: 30.5-45.0 % 31-60 days: 26.8-37.5 % 61-179 days: 28.6-37.2 % 180 days-2 years: 30.8-37.8 % 2-5 years: 31.0-37.7 % 6-11 years: 32.3-39.8 % 12-17 years: 33.9-43.5 % 18+ years: 39.6-48.8 %</p> <p>MCV</p> <p>0-14 days: 91.3-103.1 fL 15-30 days: 89.4-99.7 fL 31-60 days: 84.3-94.2 fL 61-179 days: 74.1-87.5 fL 180 days-2 years: 69.5-81.7 fL 2-5 years: 71.3-84.0 fL 6-11 years: 74.4-86.1 fL 12-17 years: 76.7-89.2 fL 18+ years: 79.0-94.5 fL</p>	OSU Internal Normal Range Study, October 2018	<p>WBC: 0.30-440.00 K/uL RBC: 0.05-8.60 M/uL HGB: See individual method</p>	<p>WBC: 0.30 K/uL-dilute to endpoint RBC: 0.05 M/uL-dilute to endpoint HGB: See individual method</p>

		<p>MCH 0-14 days: 31.1-35.9 pg 15-30 days: 30.4-35.3 pg 31-60 days: 30.0-32.5 pg 61-179 days: 24.4-29.5 pg 180 days-2 years: 23.2-27.5 pg 2-5 years: 23.7-28.6 pg 6-11 years: 24.8-29.5 pg 12-17 years: 24.8-30.2 pg 18+ years: 25.9-33.9 pg</p> <p>MCHC 0-14 days: 33.4-35.4 g/dL 15-30 days: 32.2-35.0 g/dL 31-60 days: 32.5-34.9 g/dL 61-179 days: 32.1-34.4 g/dL 180 days-2 years: 31.9-34.2 g/dL 2-5 years: 31.8-34.6 g/dL 6-11 years: 31.8-34.6 g/dL 12-17 years: 31.5-34.2 g/dL 18+ years: 31.4-35.9 g/dL</p> <p>RDW 0-14 days: 14.6-17.3 % 15-30 days: 14.4-16.2 % 31-60 days: 13.6-15.8 % 61-179 days: 12.2-14.3 % 180 days-2 years: 12.7-15.1 % 2-5 years: 12.4-14.9 % 6-11 years: 12.2-14.4 % 12-17 years: 12.3-14.6 % 18+ years: 10.8-14.9 %</p> <p>PLT 0-14 days: 144-449 K/uL 15-30 days: 279-571 K/uL 31-60 days: 331-597 K/uL 61-179 days: 247-580 K/uL 180 days-2 years: 214-459 K/uL 2-5 years: 189-394 K/uL 6-11 years: 199-367 K/uL 12-17 years: 194-345 K/uL 18+ years: 150-393 K/uL</p> <p>MPV 0-14 days: 10.4-12.0 fL 15-30 days: 10.0-12.2 fL 31-60 days: 9.4-11.1 fL 61-179 days: 9.0-10.9 fL 180 days-2 years: 8.8-10.6 fL 2-5 years: 8.9-11.0 fL 6-11 years: 9.3-11.3 fL 12-17 years: 9.6-11.7 fL 18+ years: 8.5-12.2 fL</p>	<p>MCH 0-14 days: 31.3-35.6 pg 15-30 days: 29.9-34.1 pg 31-60 days: 27.8-32.0 pg 61-179 days: 24.4-28.9 pg 180 days-2 years: 22.7-27.2 pg 2-5 years: 23.7-28.3 pg 6-11 years: 24.9-29.2 pg 12-17 years: 25.2-30.2 pg 18+ years: 26.1-33.3 pg</p> <p>MCHC 0-14 days: 33.0-35.7 g/dL 15-30 days: 32.7-35.1 g/dL 31-60 days: 32.3-34.4 g/dL 61-179 days: 31.9-34.4 g/dL 180 days-2 years: 31.6-34.4 g/dL 2-5 years: 32.0-34.7 g/dL 6-11 years: 32.2-34.9 g/dL 12-17 years: 31.8-34.8 g/dL 18+ years: 31.8-36.5 g/dL</p> <p>RDW 0-14 days: 14.8-17.0 % 15-30 days: 14.2-16.8 % 31-60 days: 13.8-16.1 % 61-179 days: 12.4-15.3 % 180 days-2 years: 12.9-15.6 % 2-5 years: 12.5-14.9 % 6-11 years: 12.3-14.1 % 12-17 years: 12.4-14.5 % 18+ years: 10.9-14.3 %</p> <p>PLT 0-14 days: 218-419 K/uL 15-30 days: 248-586 K/uL 31-60 days: 229-562 K/uL 61-179 days: 244-529 K/uL 180 days-2 years: 206-445 K/uL 2-5 years: 202-403 K/uL 6-11 years: 206-369 K/uL 12-17 years: 175-332 K/uL 18+ years: 146-337 K/uL</p> <p>MPV 0-14 days: 10.2-11.9 fL 15-30 days: 10.1-12.1 fL 31-60 days: 9.2-10.8 fL 61-179 days: 8.9-10.6 fL 180 days-2 years: 8.7-10.5 fL 2-5 years: 9.0-10.9 fL 6-11 years: 9.2-11.4 fL 12-17 years: 9.6-11.8 fL 18+ years: 8.7-12.3 fL</p>	(semiconductor laser), and sodium lauryl sulfate (SLS)-hemoglobin method.	Soldin, Steven J. <i>Pediatric Reference Intervals</i> . 7th ed., AACCPress, 2011.	<p>ICHT: See individual analyte PLT: See individual analyte</p>	<p>ICHT: See individual analyte PLT: See individual analyte</p>
Clamped Platelet, Citrated Draw	CBCP	<p>0-14 days: 144-449 K/uL 15-30 days: 279-571 K/uL 31-60 days: 331-597 K/uL 61-179 days: 247-580 K/uL 180 days-2 years: 214-459 K/uL 2-5 years: 189-394 K/uL 6-11 years: 199-367 K/uL 12-17 years: 194-345 K/uL 18+ years: 150-393 K/uL</p>	<p>0-14 days: 218-419 K/uL 15-30 days: 248-586 K/uL 31-60 days: 229-562 K/uL 61-179 days: 244-529 K/uL 180 days-2 years: 206-445 K/uL 2-5 years: 202-403 K/uL 6-11 years: 206-369 K/uL 12-17 years: 175-332 K/uL 18+ years: 146-337 K/uL</p>	Electronic resistance detection and flow cytometry.	OSU Internal Normal Range Study, October 2018	5-5,000 K/uL	≥5 K/uL
	CDPLT	<p>NRBC 0-3 days: 0.1-8.3 /100 WBC 4 days - 17 years: <0.0 /100 WBC 18+ years: <0.2 /100 WBC</p> <p>SEG-BANDS 0-14 days: 1.73-6.75 K/uL 15-30 days: 1.23-4.80 K/uL 31-60 days: 1.00-4.68 K/uL 61-179 days: 1.04-7.20 K/uL 180 days-2 years: 1.27-7.18 K/uL 2-5 years: 1.60-8.29 K/uL 6-11 years: 1.64-7.87 K/uL 12-17 years: 1.82-7.47 K/uL 18+ years: 1.64-7.28 K/uL</p> <p>LYMPHS 0-14 days: 1.75-8.00 K/uL 15-30 days: 2.42-8.20 K/uL 31-60 days: 2.29-9.14 K/uL 61-179 days: 2.14-8.99 K/uL 180 days-2 years: 1.52-8.09 K/uL 2-5 years: 1.25-5.77 K/uL 6-11 years: 1.16-4.28 K/uL 12-17 years: 1.16-3.33 K/uL 18+ years: 1.16-3.51 K/uL</p> <p>MONOABS 0-14 days: 0.27-1.72 K/uL 15-30 days: 0.42-1.21 K/uL 31-60 days: 0.28-1.21 K/uL 61-179 days: 0.24-1.17 K/uL 180 days-2 years: 0.26-1.08 K/uL 2-5 years: 0.24-0.92 K/uL 6-11 years: 0.19-0.81 K/uL 12-17 years: 0.19-0.72 K/uL 18+ years: 0.22-0.87 K/uL</p> <p>EOSABS 0-14 days: 0.09-0.64 K/uL 15-30 days: 0.06-0.75 K/uL 31-60 days: 0.04-0.63 K/uL 61-179 days: 0.02-0.74 K/uL 180 days-2 years: 0.02-0.58 K/uL 2-5 years: 0.03-0.46 K/uL 6-11 years: 0.03-0.47 K/uL 12-17 years: 0.02-0.32 K/uL 18+ years: 0.00-0.42 K/uL</p>	<p>NRBC 0-3 days: 0.1-8.3 /100 WBC 4 days - 17 years: <0.0 /100 WBC 18+ years: <0.2 /100 WBC</p> <p>SEG-BANDS 0-14 days: 1.60-6.06 K/uL 15-30 days: 1.18-5.45 K/uL 31-60 days: 0.82-4.23 K/uL 61-179 days: 0.97-5.45 K/uL 180 days-2 years: 1.19-7.21 K/uL 2-5 years: 1.54-7.92 K/uL 6-11 years: 1.63-7.55 K/uL 12-17 years: 1.54-7.04 K/uL 18+ years: 1.57-6.19 K/uL</p> <p>LYMPHS 0-14 days: 2.07-7.53 K/uL 15-30 days: 2.11-8.38 K/uL 31-60 days: 2.47-7.95 K/uL 61-179 days: 2.45-8.99 K/uL 180 days-2 years: 1.56-7.83 K/uL 2-5 years: 1.13-5.52 K/uL 6-11 years: 0.97-3.90 K/uL 12-17 years: 0.97-3.26 K/uL 18+ years: 0.83-3.57 K/uL</p> <p>MONOABS 0-14 days: 0.25-1.77 K/uL 15-30 days: 0.28-1.38 K/uL 31-60 days: 0.28-1.05 K/uL 61-179 days: 0.28-1.07 K/uL 180 days-2 years: 0.25-1.15 K/uL 2-5 years: 0.19-0.94 K/uL 6-11 years: 0.19-0.85 K/uL 12-17 years: 0.14-0.78 K/uL 18+ years: 0.24-0.93 K/uL</p> <p>EOSABS 0-14 days: 0.12-0.66 K/uL 15-30 days: 0.08-0.80 K/uL 31-60 days: 0.05-0.57 K/uL 61-179 days: 0.03-0.61 K/uL 180 days-2 years: 0.02-0.52 K/uL 2-5 years: 0.03-0.53 K/uL 6-11 years: 0.03-0.52 K/uL 12-17 years: 0.04-0.38 K/uL 18+ years: 0.00-0.48 K/uL</p>	Flow cytometry	OSU Internal Normal Range Study, October 2018	<p>SEG-BANDS%: 0.0-100.0 % LYMPHS%: 0.0-100.0 % MONO%: 0.0-100.0 % EOS%: 0.0-100.0 % BASO%: 0.0-100.0 % RRE%: 0.0-100.0 % NRBC: 0.0-600.0 /100 WBC SEG-BANDABS: 0.04-440.00 K/uL LYMPHSABS: 0.04-440.00 K/uL MONOABS: >400.00 K/uL EOSABS: >400.00 K/uL BASOABS: >400.00 K/uL</p>	<p>SEG-BANDS%: 0.0-100.0 % LYMPHS%: 0.0-100.0 % MONO%: 0.0-100.0 % EOS%: 0.0-100.0 % BASO%: 0.0-100.0 % RRE%: 0.0-100.0 % NRBC: 0.0-600.0 /100 WBC SEG-BANDABS: >400.00 K/uL LYMPHSABS: >400.00 K/uL MONOABS: >400.00 K/uL EOSABS: >400.00 K/uL BASOABS: >400.00 K/uL</p>

		<p>BASOABS 0-14 days: 0.02-0.07 K/uL 15-30 days: 0.01-0.06 K/uL 31-60 days: 0.01-0.05 K/uL 61-179 days: 0.01-0.07 K/uL 180 days-2 years: 0.01-0.06 K/uL 2-5 years: 0.01-0.06 K/uL 6-11 years: 0.01-0.05 K/uL 12-17 years: 0.01-0.05 K/uL 18+ years: 0.00-0.15 K/uL</p> <p>NEUTABS 0-1 days: <0.28 K/uL 2-13 days: <0.27 K/uL 14-30 days: <0.22 K/uL 31-90 days: <0.09 K/uL 91-180 days: <0.06 K/uL 181 days-2 years: <0.14 K/uL 2-5 years: <0.06 K/uL 6-11 years: <0.04 K/uL 12-17 years: <0.04 K/uL 18+ years: <0.08 K/uL</p>	<p>BASOABS 0-14 days: 0.02-0.11 K/uL 15-30 days: 0.01-0.07 K/uL 31-60 days: 0.01-0.07 K/uL 61-179 days: 0.01-0.06 K/uL 180 days-2 years: 0.01-0.06 K/uL 2-5 years: 0.01-0.06 K/uL 6-11 years: 0.01-0.06 K/uL 12-17 years: 0.01-0.05 K/uL 18+ years: 0.00-0.09 K/uL</p> <p>NEUTABS 0-1 days: <0.28 K/uL 2-13 days: <0.27 K/uL 14-30 days: <0.22 K/uL 31-90 days: <0.09 K/uL 91-180 days: <0.06 K/uL 181 days-2 years: <0.14 K/uL 2-5 years: <0.06 K/uL 6-11 years: <0.04 K/uL 12-17 years: <0.04 K/uL 18+ years: <0.07 K/uL</p>					
	CDC EDIFF EDIFF							
Extended Reticulocyte Panel		<p>RETIC 0-3 days: 3.47-5.40 % 4-30 days: 1.06-2.37 % 31-60 days: 2.12-3.47 % 61-179 days: 1.55-2.70 % 180 days-2 years: 0.99-1.82 % 2-5 years: 0.82-1.45 % 6-11 years: 0.86-1.04 % 12-17 years: 0.90-1.49 % 18+ years: 0.74-2.54 %</p> <p>RETICABS 0-3 days: 0.1475-0.2164 M/uL 4-30 days: 0.0513-0.1104 M/uL 31-60 days: 0.0518-0.0779 M/uL 61-179 days: 0.0482-0.0882 M/uL 180 days-2 years: 0.0425-0.1111 M/uL 2-5 years: 0.0364-0.0680 M/uL 6-11 years: 0.0424-0.0702 M/uL 12-17 years: 0.0416-0.0651 M/uL 18+ years: 0.0324-0.1142 M/uL</p> <p>IRETF 0-3 days: 30.5-35.1 % 4-30 days: 14.5-24.6 % 31-60 days: 19.1-28.9 % 61-179 days: 13.4-23.3 % 180 days-2 years: 11.4-25.8 % 2-5 years: 8.4-21.7 % 6-11 years: 8.9-24.1 % 12-17 years: 9.0-18.7 % 18+ years: 1.1-16.2 %</p> <p>RETIC 0-179 days: 29.2-37.5 pg 180 days-2 years: 30.1-35.7 pg 2-5 years: 29.3-37.3 pg 6-11 years: 30.4-39.7 pg 12-17 years: 29.9-38.4 pg 18+ years: 28.8-39.9 pg</p>	<p>RETIC 0-3 days: 3.47-5.40 % 4-30 days: 1.06-2.37 % 31-60 days: 2.12-3.47 % 61-179 days: 1.55-2.70 % 180 days-2 years: 0.99-1.82 % 2-5 years: 0.82-1.45 % 6-11 years: 0.86-1.04 % 12-17 years: 0.90-1.49 % 18+ years: 0.68-2.64 %</p> <p>RETICABS 0-3 days: 0.1475-0.2164 M/uL 4-30 days: 0.0513-0.1104 M/uL 31-60 days: 0.0518-0.0779 M/uL 61-179 days: 0.0482-0.0882 M/uL 180 days-2 years: 0.0425-0.1111 M/uL 2-5 years: 0.0364-0.0680 M/uL 6-11 years: 0.0424-0.0702 M/uL 12-17 years: 0.0416-0.0651 M/uL 18+ years: 0.0317-0.1377 M/uL</p> <p>IRETF 0-3 days: 30.5-35.1 % 4-30 days: 14.5-24.6 % 31-60 days: 19.1-28.9 % 61-179 days: 13.4-23.3 % 180 days-2 years: 11.4-25.8 % 2-5 years: 8.4-21.7 % 6-11 years: 8.9-24.1 % 12-17 years: 9.0-18.7 % 18+ years: 0.2-16.3 %</p> <p>RETIC 0-179 days: 27.6-38.7 pg 180 days-2 years: 28.7-35.7 pg 2-5 years: 27.7-37.8 pg 6-11 years: 32.4-37.6 pg 12-17 years: 30.3-40.4 pg 18+ years: 29.9-38.7 pg</p>	Flow cytometry / calculation	OSU Internal Normal Range Study, October 2018 Sollán, Steven J. <i>Pediatric Reference Intervals</i> . 7th ed., AACCPress, 2011.	<p>RETIC: 0.25-30.00% RETICABS: 0.0100-0.7200 XN-L: 0.0100-0.4576 Manual: 0.0100-2.5800 IRETF: 0.0-100.0 % RETIC: 0.0-100.0 %</p>	<p>RETIC: 0.25-30.00 % RETICABS: 0.0100-2.5800 Manual: 0.0100-2.5800* IRETF: 0.0-100.0 %</p> <p>*Instrument may prompt dilution but final results should not exceed 2.5800</p>	
	EXRETIC IRF RET:HE	See HEM3 and LAB256	See HEM3 and LAB256					
		<p>BLAST% <0.0 %</p> <p>LYMPHOMA% <0.0 %</p> <p>HABRY% <0.0 %</p> <p>PLASMA% <0.0 %</p> <p>PROLYMPH% <0.0 %</p> <p>OTHER% <0.0 %</p> <p>NRBC 0.0-0.2 /100 WBC</p> <p>SEG-BANDABS 0-14 days: 1.23-6.75 K/uL 15-30 days: 1.23-4.80 K/uL 31-60 days: 1.00-4.68 K/uL 61-179 days: 1.04-2.20 K/uL 180 days-2 years: 1.27-7.18 K/uL 2-5 years: 1.64-8.29 K/uL 6-11 years: 1.64-7.87 K/uL 12-17 years: 1.82-7.47 K/uL 18+ years: 1.64-7.28 K/uL</p> <p>LYMPHABS 0-14 days: 1.75-8.00 K/uL 15-30 days: 2.42-8.20 K/uL 31-60 days: 2.29-9.14 K/uL 61-179 days: 2.14-8.99 K/uL 180 days-2 years: 1.52-8.09 K/uL 2-5 years: 1.28-5.77 K/uL 6-11 years: 1.16-4.28 K/uL 12-17 years: 1.16-3.33 K/uL 18+ years: 1.16-3.31 K/uL</p> <p>MONOABS 0-14 days: 0.57-1.72 K/uL 15-30 days: 0.42-1.21 K/uL 31-60 days: 0.26-1.21 K/uL 61-179 days: 0.24-1.17 K/uL 180 days-2 years: 0.26-1.08 K/uL 2-5 years: 0.24-0.92 K/uL</p>	<p>BLAST% <0.0 %</p> <p>LYMPHOMA% <0.0 %</p> <p>HABRY% <0.0 %</p> <p>PLASMA% <0.0 %</p> <p>PROLYMPH% <0.0 %</p> <p>OTHER% <0.0 %</p> <p>NRBC 0.0-0.2 /100 WBC</p> <p>SEG-BANDABS 0-14 days: 1.69-6.06 K/uL 15-30 days: 1.18-5.45 K/uL 31-60 days: 0.83-4.21 K/uL 61-179 days: 0.97-5.45 K/uL 180 days-2 years: 1.19-7.21 K/uL 2-5 years: 1.54-7.92 K/uL 6-11 years: 1.63-7.53 K/uL 12-17 years: 1.54-7.04 K/uL 18+ years: 1.57-6.19 K/uL</p> <p>LYMPHABS 0-14 days: 2.07-7.53 K/uL 15-30 days: 2.11-8.38 K/uL 31-60 days: 2.47-7.95 K/uL 61-179 days: 2.45-8.89 K/uL 180 days-2 years: 1.56-7.83 K/uL 2-5 years: 1.13-5.52 K/uL 6-11 years: 0.97-3.96 K/uL 12-17 years: 0.97-3.26 K/uL 18+ years: 0.83-3.57 K/uL</p> <p>MONOABS 0-14 days: 0.52-1.77 K/uL 15-30 days: 0.28-1.38 K/uL 31-60 days: 0.28-1.05 K/uL 61-179 days: 0.28-1.07 K/uL 180 days-2 years: 0.25-1.15 K/uL 2-5 years: 0.19-0.94 K/uL</p>					
								See HEM3 and LAB256
								See HEM3 and LAB256
								<p>BAND%: 0.0-100.0 % SE%: 0.0-100.0 % LYMPH%: 0.0-100.0 % MONO%: 0.0-100.0 % EOS%: 0.0-100.0 % BASO%: 0.0-100.0 % MET%: 0.0-100.0 % MYELO%: 0.0-100.0 %</p>

Innate Platelet Fraction	IPF 0-179 days: 1.3-6.8 % 180 days-2 years: 1.4-4.5 % 2-5 years: 1.0-3.6 % 6-11 years: 1.0-4.7 % 12-17 years: 1.4-6.4 % 18+ years: 0.0-8.6 % PLTICPF 0-14 days: 144-449 K/uL 15-30 days: 279-571 K/uL 31-60 days: 331-597 K/uL 61-179 days: 247-580 K/uL 180 days-2 years: 214-459 K/uL 2-5 years: 189-394 K/uL 6-11 years: 199-367 K/uL 12-17 years: 194-345 K/uL 18+ years: 150-393 K/uL IPFPLT IPF	IPF 0-179 days: 2.0-6.8 % 180 days-2 years: 1.4-3.8 % 2-5 years: 1.1-3.9 % 6-11 years: 1.0-4.9 % 12-17 years: 1.6-6.1 % 18+ years: 0.0-9.0 % PLTICPF 0-14 days: 218-419 K/uL 15-30 days: 248-586 K/uL 31-60 days: 224-562 K/uL 61-179 days: 244-529 K/uL 180 days-2 years: 206-445 K/uL 2-5 years: 202-403 K/uL 6-11 years: 206-369 K/uL 12-17 years: 175-332 K/uL 18+ years: 146-337 K/uL	Calculation	OSU Internal Normal Range Study, October 2018 Saldin, Steven J. <i>Pediatric Reference Intervals</i> . 7th ed., AACCPress, 2011.	IPF: 0.0-100.0 % PLTICPF: 5-5,000 K/uL	IPF: 0.0-100.0 % PLTICPF: 5 K/uL-dilute to endpoint	
Malaria Prep	Parasitic Screen ID Blood MPB			Giemsa stain		No parasitic organisms seen, including <i>Plasmodium</i> organism <i>Positive for Plasmodium species</i>	
Platelet Count	PLT 0-14 days: 144-449 K/uL 15-30 days: 279-571 K/uL 31-60 days: 331-597 K/uL 61-179 days: 247-580 K/uL 180 days-2 years: 214-459 K/uL 2-5 years: 189-394 K/uL 6-11 years: 199-367 K/uL 12-17 years: 194-345 K/uL 18+ years: 150-393 K/uL MPV 0-14 days: 10.4-12.0 fL 15-30 days: 10.0-12.2 fL 31-60 days: 9.4-11.1 fL 61-179 days: 9.0-10.9 fL 180 days-2 years: 8.8-10.6 fL 2-5 years: 8.9-11.0 fL 6-11 years: 9.3-11.3 fL 12-17 years: 9.6-11.7 fL 18+ years: 8.5-12.2 fL PLAT	PLT 0-14 days: 218-419 K/uL 15-30 days: 248-586 K/uL 31-60 days: 224-562 K/uL 61-179 days: 244-529 K/uL 180 days-2 years: 206-445 K/uL 2-5 years: 202-403 K/uL 6-11 years: 206-369 K/uL 12-17 years: 175-332 K/uL 18+ years: 146-337 K/uL MPV 0-14 days: 10.2-11.9 fL 15-30 days: 10.1-12.1 fL 31-60 days: 9.2-10.8 fL 61-179 days: 8.9-10.6 fL 180 days-2 years: 8.7-10.5 fL 2-5 years: 9.0-10.9 fL 6-11 years: 9.2-11.4 fL 12-17 years: 9.6-11.8 fL 18+ years: 8.7-12.3 fL	Electronic resistance detection and flow cytometry.	OSU Internal Normal Range Study, October 2018 Saldin, Steven J. <i>Pediatric Reference Intervals</i> . 7th ed., AACCPress, 2011.	5-5,000 K/uL	5 K/uL-dilute to endpoint	
Reiculocytes	RETIC 0-3 days: 3.47-5.40 % 4-30 days: 1.06-2.37 % 31-60 days: 2.12-3.47 % 61-179 days: 1.55-2.70 % 180 days-2 years: 0.99-1.82 % 2-5 years: 0.82-1.45 % 6-11 years: 0.98-1.94 % 12-17 years: 0.90-1.49 % 18+ years: 0.74-2.34 % RETICABS 0-3 days: 0.175-0.2164 M/uL 4-30 days: 0.0513-0.1104 M/uL 31-60 days: 0.0518-0.0779 M/uL 61-179 days: 0.0462-0.0882 M/uL 180 days-2 years: 0.0435-0.1111 M/uL 2-5 years: 0.0364-0.0680 M/uL 6-11 years: 0.0424-0.0702 M/uL 12-17 years: 0.0416-0.0651 M/uL 18+ years: 0.0324-0.1142 M/uL	RETIC 0-3 days: 3.47-5.40 % 4-30 days: 1.06-2.37 % 31-60 days: 2.12-3.47 % 61-179 days: 1.55-2.70 % 180 days-2 years: 0.99-1.82 % 2-5 years: 0.82-1.45 % 6-11 years: 0.98-1.94 % 12-17 years: 0.90-1.49 % 18+ years: 0.68-2.64 % RETICABS 0-3 days: 0.1475-0.2164 M/uL 4-30 days: 0.0513-0.1104 M/uL 31-60 days: 0.0518-0.0779 M/uL 61-179 days: 0.0462-0.0882 M/uL 180 days-2 years: 0.0435-0.1111 M/uL 2-5 years: 0.0364-0.0680 M/uL 6-11 years: 0.0424-0.0702 M/uL 12-17 years: 0.0416-0.0651 M/uL 18+ years: 0.0317-0.1377 M/uL		Flow cytometry	OSU Internal Normal Range Study, October 2018 Saldin, Steven J. <i>Pediatric Reference Intervals</i> . 7th ed., AACCPress, 2011.	RETIC: 0.25-30.00% RETICABS: XN: 0.0100-0.7200 XN-L: 0.0100-0.4576 Manual: 0.0100-2.5800	RETIC: 0.25-30.00 % RETICABS: 0.0100-2.5000 Manual: 0.0100-2.5800* *Instrument may prompt dilution but final result should not exceed 2.5800
Schistocytes	PSCHISTO			Manual			
Sedimentation Rate, Automated	ESR	0-49 years: <20 mm/hr 50-84 years: <30 mm/hr 85+ years: <42 mm/hr	0-49 years: <15 mm/hr 50-84 years: <20 mm/hr 85+ years: <30 mm/hr	Westergren	McPherson, R.A., & Pincus, M.R. (2017). <i>Henry's Clinical Diagnosis and Management by Laboratory Methods</i> (23rd ed.). St. Louis, MO: Elsevier Inc. pg. 532	1-140 mm/hr	1-140 mm/hr
White Blood Count	WBC	0-14 days: 8.16-14.56 K/uL 15-30 days: 8.36-14.42 K/uL 31-60 days: 7.05-14.68 K/uL 61-179 days: 6.00-13.25 K/uL 180 days-2 years: 6.48-13.02 K/uL 2-5 years: 4.80-13.18 K/uL 6-11 years: 4.27-11.40 K/uL 12-17 years: 4.19-9.43 K/uL 18+ years: 3.99-11.19 K/uL	0-14 days: 8.04-15.40 K/uL 15-30 days: 7.80-15.91 K/uL 31-60 days: 8.14-14.99 K/uL 61-179 days: 6.51-13.32 K/uL 180 days-2 years: 5.98-13.51 K/uL 2-5 years: 5.14-13.33 K/uL 6-11 years: 4.31-11.00 K/uL 12-17 years: 3.84-9.84 K/uL 18+ years: 3.73-10.10 K/uL	Flow cytometry	OSU Internal Normal Range Study, October 2018 Saldin, Steven J. <i>Pediatric Reference Intervals</i> . 7th ed., AACCPress, 2011.	0.30-440.00 K/uL	0.30 K/uL-dilute to endpoint
Body Fluid Cell Count	FCNT			Hemocytometer counts and Iris instrument.		TNC: 3-2,500 /uL RBC: 3-50,000 /uL	TNC: ≥3 /uL RBC: ≥3 /uL
Body Fluid Cell Count with DfF	FCNT FPDfF (Body fluid differential)	SFBLAST 0% SFMALIG 0% SFLYMPHOM 0%		Hemocytometer counts, Iris instrument and manual differential.		TNC: 3-2,500 /uL RBC: 3-50,000 /uL FSEG: 0-100 % FLYME: 0-100 % FMOMAC: 0-100 % FMESO: 0-100 % FEOS: 0-100 % FRASO: 0-100 % FPLAS: 0-100 % FMALIG: 0-100 % FRLAS: 0-100 % FLPHOM: 0-100 % FSYMLC: 0-100 % FSYRBC: 0-100 % FOTHR: 0-100 %	TNC: ≥3 /uL RBC: ≥3 /uL FSEG: 0-100 % FLYME: 0-100 % FMOMAC: 0-100 % FMESO: 0-100 % FEOS: 0-100 % FRASO: 0-100 % FPLAS: 0-100 % FMALIG: 0-100 % FRLAS: 0-100 % FLPHOM: 0-100 % FSYMLC: 0-100 % FSYRBC: 0-100 % FOTHR: 0-100 %

Cell Count & Diff. CSF	CSFLC CSFDF Spinal fluid cell count and body fluid differential	SFWBC 0-365 days <31 /ul. 1-4 years <21 /ul. 5+ years <6 /ul. SFRBC <3 /ul. FSEEG 0-89 days <5 % 90+ days <6 % SFLYM 0-89 days 35-50% 90+ days 40-80 % SFMOMA 0-89 days 50-90% 90+ days 15-45 %		Hemocytometer counts, Iris instrument and manual differential.	Body Fluids 3rd ed. Kjeldsberg, Knight 1993	SFWBC: 3-2,500 /ul SFRBC: 3-50,000 /ul FSEEG: 0-100 % SFLYM: 0-100 % SFMOMA: 0-100 % SFEOS: 0-100 % SFBAS: 0-100 % SFPLAS: 0-100 % SFMALG: 0-100 % SFBLAS: 0-100 % SFLPHOM: 0-100 % SFNRBC: 0-100 % SFOTHR: 0-100 %	SGROS: Amber, Bloody, Brown, Coliforms, Green, Orange, Pink, Red, Red-cell fringed, White, Xanthochromic, Clear, Hazy, Opaque SSPN: Amber, Bloody, Brown, Coliforms, Green, Orange, Pink, Red, Red-cell fringed, White, Xanthochromic, Clear, Hazy, Opaque SFRBC: <3 /ul FSEEG: 0-100 % SFLYM: 0-100 % SFMOMA: 0-100 % SFEOS: 0-100 % SFBAS: 0-100 % SFPLAS: 0-100 % SFMALG: 0-100 % SFBLAS: 0-100 % SFLPHOM: 0-100 % SFNRBC: 0-100 % SFOTHR: 0-100 %
Crystals, Fluid	PCRV5			Negative	Unstained synovial fluid slides reviewed by polarized microscopy.		Negative, Positive
CSF Fluid Count Only	CSFLC Spinal fluid cell count	SFWBC 0-365 days <31 /ul. 1-4 years <21 /ul. 5+ years <6 /ul. SFRBC <3 /ul.		Hemocytometer counts and Iris instrument.	Body Fluids 3rd ed. Kjeldsberg, Knight 1993	SFWBC: 3-2,500 /ul SFRBC: 3-50,000 /ul	CSF TUBERCULOSIS: CSF Tube 1, CSF Tube 2, CSF Tube 3, CSF Tube 4, No tube number indicated SGROS: Amber, Bloody, Brown, Coliforms, Green, Orange, Pink, Red, Red-cell fringed, White, Xanthochromic, Clear, Hazy, Opaque
Differential, Fluid		SFBLAST 0% SFMALG 0% SFLYMPHOM 0%		Microscopic evaluation of body fluid cells is performed by reviewing a Wright-stained body fluid cytoplasm slide using a brightfield microscope.		FSEEG: 0-100 % FLYM: 0-100 % FMOMAC: 0-100 % FMESD: 0-100 % FEOS: 0-100 % FBIASO: 0-100 % EPLAS: 0-100 % FMALG: 0-100 % FBLAS: 0-100 % FLPHOM: 0-100 % FSYNLC: 0-100 % FSNRBC: 0-100 % FOTHR: 0-100 %	FSEEG: 0-100 % FLYM: 0-100 % FMOMAC: 0-100 % FMESD: 0-100 % FEOS: 0-100 % FBIASO: 0-100 % EPLAS: 0-100 % FMALG: 0-100 % FBLAS: 0-100 % FLPHOM: 0-100 % FSYNLC: 0-100 % FSNRBC: 0-100 % FOTHR: 0-100 %
Hematocrit, Fluid	HDHPT Fluid HCT Fluid PCV			Manual spun hematocrit.		5.0-60.0 %	5.0-60.0 %
Myoglobin Urine	MYO Urine myoglobin screen			Negative	The peroxidase-like activity of hemoglobin catalyzes the reaction of diaminopyrene dithiopyrosin and 3,3',5,5'-tetramethylbenzidine to produce a color from orange to green.	Urinanalysis and Body Fluid, Ringwald 1995	UBLD: Negative, Trace, Small, Moderate, Large MYO: Negative, Reflexed to Sendout
Ocult Blood, Fecal Immunological	FOB			Negative	Immunossay utilizing rabbit polyclonal antibodies to detect presence of hemoglobin in feces.		Negative, Positive
Ocult Blood, Gastric	OCBDG Gastrocult			Negative	Developing solution stabilized mixture of hydrogen peroxide and denatured alcohol creates a reaction between hemoglobin and guaic to produce a blue color.		Negative, Positive
Ocult Blood, Stool	OCBDF Ocult blood Fecal hemocult			Negative	Developing solution stabilized mixture of hydrogen peroxide and denatured alcohol creates a reaction between hemoglobin and guaic to produce a blue color.		Negative, Positive
pH, Urine		5.0-7.0			Double indicator principle to cover the range of urinary pH range. Colors range from orange through yellow and green to blue.	Urinanalysis and Body Fluid, Ringwald 1995	MMMP: 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, >8.5. Unable to analyze due to interfering substance. SSHC: 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, >8.5. Unable to analyze due to interfering substance. Unspecified: 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5, >9.0. Unable to analyze due to interfering substance.
Specific Gravity, Urine	UPH	1.001-1.035			Clinical Lab CCL: Fiber optic refractive index method. Clinical Lab University East: Fiber optic refractive index method. Clinical Lab James: Fiber optic refractive index method. All Performing Labs: pKa change of pretreated polyelectrolyte in relation to ionic concentration.	Urinanalysis and Body Fluid, Ringwald 1995	CCL: 1.001-1.045 RRL: 1.001-1.045 JAMES: 1.001-1.045 Unspecified: 1.005-1.030
Synovasure PII, Synovial Fluid	SYNOVA Synovasure Alpha Defensin Lateral Flow Test			Negative	Qualitative, visually read immunochromatographic assay for the detection of human host response proteins, Alpha Defensins 1-3, in the synovial fluid of adults with a total joint replacement who are being evaluated for revision surgery.		Negative, Positive
Urinanalysis							COLOR: Yellow, Orange, Red, See Comment APPEAR: Clear, Cloudy, Turbid, Unable to analyze due to interfering substance. GLUCOSE: Negative, 100 mg/dL, 250 mg/dL, 500 mg/dL, ≥1000 mg/dL, Unable to analyze due to interfering substance. KETONE: Negative, Trace, 15 mg/dL, Small, 40 mg/dL, Moderate, ≥80 mg/dL, Large, Unable to analyze due to interfering substance. SPECKLE GRAVITY: See individual analyte BLOOD: Negative, Trace, Small, Moderate, Large, Unable to analyze due to interfering substance. PH: See individual analyte PROTEIN: Negative, Trace, 30 mg/dL, 100 mg/dL, 300 mg/dL, ≥300 mg/dL, Unable to analyze due to interfering substance.

	URIN			<p>SQUAMOUS: 0-2/hpf, 3-5/hpf = 1+</p> <p>BACTERIA: Absent</p> <p>RENAL TUBULAR CELLS: 0-2/hpf</p> <p>WBC CASTS: Absent</p> <p>WBC/RENAL TUBULAR CASTS: Absent</p> <p>RBC CASTS: Absent</p> <p>YEAST: Absent</p> <p>TRICHOMONAS: Absent</p> <p>TRICHOMONAS: Absent</p> <p>TYROSINE: Absent</p> <p>CYSTINE: Absent</p> <p>TRIPLE PHOSPHATE: Absent</p>	Dry pad urine chemistry or optic refractive index method			<p>UROBILINOGEN: 0.2 E.U./dL, 1.0 E.U./dL, 2.0 E.U./dL, 4.0 E.U./dL, >8.0 E.U./dL. Unable to analyze due to interfering substance.</p> <p>NITRITE: Negative. Positive. Unable to analyze due to interfering substance.</p> <p>LEUKOCYTE ESTERASE: Negative. Trace, Small, Moderate, Large. Unable to analyze due to interfering substance.</p> <p>RBC: 0-2, 3-5, 6-10, 11-25, >25 HPF</p> <p>WBC: 0-5, 6-10, 11-20, >20 HPF</p> <p>SQUAMOUS: 0-2/hpf, 3-5/hpf = 1+, 6-10/hpf = 2+, 11-20/hpf = 3+, >20/hpf = 4+</p> <p>BACTERIA: Absent, Trace, Present</p> <p>RENAL TUBULAR CELLS: 0-2/hpf, 3-5/hpf = 1+, 6-10/hpf = 2+, 11-20/hpf = 3+, >20/hpf = 4+</p> <p>AMORPHOUS: <25% = Slight, 26-50% = Moderate, >51% = Heavy</p> <p>HYALINE CASTS: 0-2, 3-5, 6-10, 11-20, >20 LPF</p> <p>GRANULAR CASTS: 0-2, 3-5, 6-10, 11-</p>
Urinalysis Reflex to Culture	URIN1			<p>COLOR: Yellow</p> <p>APPEARANCE: Clear</p> <p>GLUCOSE: Negative</p> <p>KETONE: Negative</p> <p>SPECIFIC GRAVITY: 1.001-1.035</p> <p>BLOOD: Negative</p> <p>PH: 5.0, 5.5, 6.0, 6.5, 7.0</p> <p>PROTEIN: Negative</p> <p>UROBILINOGEN: 0.2 E.U./dL, 1.0 E.U./dL</p> <p>NITRITE: Negative</p> <p>LEUKOCYTE ESTERASE: Negative</p> <p>RBC: 0-2 HPF</p> <p>WBC: 0-5 HPF</p> <p>SQUAMOUS: 0-2/hpf, 3-5/hpf = 1+</p> <p>BACTERIA: Absent</p> <p>RENAL TUBULAR CELLS: 0-2/hpf</p> <p>WBC CASTS: Absent</p> <p>WBC/RENAL TUBULAR CASTS: Absent</p> <p>RBC CASTS: Absent</p> <p>YEAST: Absent</p> <p>TRICHOMONAS: Absent</p> <p>LEUCINE: Absent</p> <p>TYROSINE: Absent</p> <p>CYSTINE: Absent</p> <p>TRIPLE PHOSPHATE: Absent</p>	<p>DIPSTICK</p> <p>Dry pad urine chemistry</p> <p>SG</p> <p>Dry pad urine chemistry or optic refractive index method</p> <p>URINE MICROSCOPIC</p> <p>Automatic particle counter and/or manual microscopic</p>	Varies	See LAB347	
Urine Dipstick with Reflex to Microscopy	URIN1			<p>COLOR: Yellow</p> <p>APPEARANCE: Clear</p> <p>GLUCOSE: Negative</p> <p>KETONE: Negative</p> <p>SPECIFIC GRAVITY: 1.001-1.035</p> <p>BLOOD: Negative</p> <p>PH: 5.0, 5.5, 6.0, 6.5, 7.0</p> <p>PROTEIN: Negative</p> <p>UROBILINOGEN: 0.2 E.U./dL, 1.0 E.U./dL</p> <p>NITRITE: Negative</p> <p>LEUKOCYTE ESTERASE: Negative</p>	<p>DIPSTICK</p> <p>Dry pad urine chemistry</p> <p>SG</p> <p>Dry pad urine chemistry or optic refractive index method</p>	Varies	See URN11	
Urine Dipstick; Reflex Microscopy; Reflex Culture	UASR1			<p>COLOR: Yellow</p> <p>APPEARANCE: Clear</p> <p>GLUCOSE: Negative</p> <p>KETONE: Negative</p> <p>SPECIFIC GRAVITY: 1.001-1.035</p> <p>BLOOD: Negative</p> <p>PH: 5.0, 5.5, 6.0, 6.5, 7.0</p> <p>PROTEIN: Negative</p> <p>UROBILINOGEN: 0.2 E.U./dL, 1.0 E.U./dL</p> <p>NITRITE: Negative</p> <p>LEUKOCYTE ESTERASE: Negative</p>	<p>DIPSTICK</p> <p>Dry pad urine chemistry</p> <p>SG</p> <p>Dry pad urine chemistry or optic refractive index method</p>	National guidelines and recommendations	See URN11	

<p>Urine Screen</p> <p>UAS Using dipstick</p>				<p>COLOR: Yellow APPEARANCE: Clear GLUCOSE: Negative KETONE: Negative SPECIFIC GRAVITY: 1.001-1.035 BLOOD: Negative PH: 5.0, 5.5, 6.0, 6.5, 7.0 PROTEIN: Negative UROBILINOGEN: 0.2 E.U./dL, 1.0 E.U./dL NITRITE: Negative LEUKOCYTE ESTERASE: Negative</p>	<p>DIPSTICK Dry pad urine chemistry SG Dry pad urine chemistry or optic refractive index method</p>	<p>Varies</p>	<p>COLOR: Yellow, Orange, Red, See Comment APPEARANCE: Clear, Cloudy, Turbid, Unable to analyze due to interfering substance. GLUCOSE: Negative, 100 mg/dL, 250 mg/dL, 500 mg/dL, >1000 mg/dL, Unable to analyze due to interfering substance. KETONE: Negative, Trace, 15 mg/dL, Small, 40 mg/dL, Moderate, 250 mg/dL = Large, Unable to analyze due to interfering substance. SPECIFIC GRAVITY: See individual analyze BLOOD: Negative, Trace, Small, Moderate, Large, Unable to analyze due to interfering substance. PH: See individual analyze PROTEIN: Negative, Trace, 30 mg/dL, 100 mg/dL, 200 mg/dL, >300 mg/dL, Unable to analyze due to interfering substance. UROBILINOGEN: 0.2 E.U./dL, 1.0 E.U./dL, 2.0 E.U./dL, 4.0 E.U./dL, >8.0 E.U./dL, Unable to analyze due to interfering substance. NITRITE: Negative, Positive, Unable to analyze due to interfering substance. LEUKOCYTE ESTERASE: Negative, Trace, Small, Moderate, Large, Unable to analyze due to interfering substance.</p>																																																
<p>Anti Phospholipid Antibody</p>	<p>PLAB Anticardiolipin Antibodies IgG, IgM Antiphospholipid Antibodies IgG, IgM Cardiolipin Antibodies IgG, IgM Phospholipid Antibodies IgG, IgM</p>	<p>PLABG 0.0-20.0 CU PLABM 0.0-20.0 CU</p>		<p>Chemiluminescent two-step immunoassay.</p>	<p>Inova Quanta-Flash Package Insert</p>	<p>PLABG: 10.0-500.0 CU PLABM: 10.0-200.0 CU</p>	<p>PLABG: 10.0, 10.000.0 CU PLABM: 10.0, 4.000.0 CU</p>																																																
<p>Anti Xa DOAC (Apixaban)</p> <p>AXAPFX Elisum</p>			<p>Therapeutic reference ranges have not been established. At steady state - median (5th-95th percentile) peak and trough levels have been observed in clinical trial:</p> <table border="1"> <thead> <tr> <th>Peak Conc., ng/mL (a)</th> <th>Trough Conc., ng/mL (b)</th> </tr> </thead> <tbody> <tr> <td colspan="2">-----</td> </tr> <tr> <td colspan="2">VTE prevention after total hip replacement</td> </tr> <tr> <td>2.5 mg twice daily</td> <td>77 (41-146)</td> </tr> <tr> <td>5 mg twice daily</td> <td>51 (23-109)</td> </tr> </tbody> </table> <p>Stroke and systemic embolism prevention in patients with AF</p> <table border="1"> <tbody> <tr> <td>2.5 mg twice daily</td> <td>123 (69-221)</td> <td>79 (34-162)</td> </tr> <tr> <td>5 mg twice daily</td> <td>171 (91-321)</td> <td>103 (41-230)</td> </tr> </tbody> </table> <p>Prevention and treatment of DVT and PE</p> <table border="1"> <tbody> <tr> <td>2.5 mg twice daily</td> <td>67 (30-151)</td> <td>32 (11-90)</td> </tr> <tr> <td>5 mg twice daily</td> <td>132 (59-302)</td> <td>63 (22-77)</td> </tr> <tr> <td>10 mg twice daily</td> <td>251 (111-572)</td> <td>120 (41-335)</td> </tr> </tbody> </table> <p>(a) Defined as samples collected 2-4 hours after dosing (b) Defined as samples collected 10-12 hours after dosing</p> <p>AF-atrial fibrillation, CR-CL-creatinine clearance, DVT-deep vein thrombosis, VTE-venous thromboembolism</p>	Peak Conc., ng/mL (a)	Trough Conc., ng/mL (b)	-----		VTE prevention after total hip replacement		2.5 mg twice daily	77 (41-146)	5 mg twice daily	51 (23-109)	2.5 mg twice daily	123 (69-221)	79 (34-162)	5 mg twice daily	171 (91-321)	103 (41-230)	2.5 mg twice daily	67 (30-151)	32 (11-90)	5 mg twice daily	132 (59-302)	63 (22-77)	10 mg twice daily	251 (111-572)	120 (41-335)	<p>Chromogenic measurement at 405nm.</p>	<p>1. Package insert: Rivaroxaban: Diagnostics stage. Revised December 2014 2. Moeck W, Stampfuss J, Kubitz D, Becka M: Clinical pharmacokinetic and pharmacodynamic profile of rivaroxaban. Clinical Pharmacokinetics 2014; 53(1):1-16 doi: 10.1007/s40262-013-0100-7 3. Bayer Pharma AG: Xarelto (rivaroxaban) Summary of Product Characteristics; 2013. Available at: www.ema.europa.eu/doc/en_GB/document_library/EPAR_-_Product_Information/human/000944/WC50005710_8.pdf 4. EINSTEIN Investigators, Basenachs R, Berkowitz SD, et al: Oral rivaroxaban for symptomatic venous thromboembolism. N Engl J Med 2010; 363:2499-510 5. EINSTEIN-PE Investigators, Buller HR, Prins MH, et al: Oral rivaroxaban for the treatment of symptomatic pulmonary embolism. N Engl J Med 2012; 366:1287-1297 6. Patel MR, Mahaffey KW, Garg J, et al: Rivaroxaban versus warfarin in nonvalvular atrial fibrillation. N Engl J Med 2011; 365:883-891 7. Sigal DM, Carruste JT, Connolly SJ, et al: Andexanet alfa for reversal of factor Xa inhibitor</p>	<p>23-500 ng/mL</p>	<p>23-500 ng/mL</p>																							
Peak Conc., ng/mL (a)	Trough Conc., ng/mL (b)																																																						

VTE prevention after total hip replacement																																																							
2.5 mg twice daily	77 (41-146)																																																						
5 mg twice daily	51 (23-109)																																																						
2.5 mg twice daily	123 (69-221)	79 (34-162)																																																					
5 mg twice daily	171 (91-321)	103 (41-230)																																																					
2.5 mg twice daily	67 (30-151)	32 (11-90)																																																					
5 mg twice daily	132 (59-302)	63 (22-77)																																																					
10 mg twice daily	251 (111-572)	120 (41-335)																																																					
<p>Anti Xa DOAC (Rivaroxaban)</p> <p>AXRIVA Xarelto</p>			<p>Therapeutic reference ranges have not been established. At steady state - median (5th-95th percentile) peak and trough levels have been observed in clinical trial:</p> <table border="1"> <thead> <tr> <th>Dose</th> <th>Indication</th> <th>Peak Conc., ng/mL (a)</th> <th>Trough Conc., ng/mL (b)</th> </tr> </thead> <tbody> <tr> <td colspan="4">-----</td> </tr> <tr> <td>2.5 mg</td> <td>Acute coronary</td> <td>46 (28-70)</td> <td>17 (6-37)</td> </tr> <tr> <td></td> <td>twice daily syndrome</td> <td></td> <td></td> </tr> <tr> <td>10 mg</td> <td>VTE prevention</td> <td>125 (91-196)</td> <td>9 (1-38)</td> </tr> <tr> <td></td> <td>once daily after total hip replacement</td> <td></td> <td></td> </tr> <tr> <td>15 mg</td> <td>Stroke prevention</td> <td>229 (178-313)</td> <td>57 (18-136)</td> </tr> <tr> <td></td> <td>once daily in patients with AF (CR-CL 30-49 mL/min)</td> <td></td> <td></td> </tr> <tr> <td>20 mg</td> <td>DVT treatment</td> <td>270 (189-419)</td> <td>26 (6-87)</td> </tr> <tr> <td></td> <td>once daily</td> <td></td> <td></td> </tr> <tr> <td>20 mg</td> <td>Stroke prevention</td> <td>249 (184-343)</td> <td>44 (12-137)</td> </tr> <tr> <td></td> <td>once daily in patients with AF (CR-CL ≥50 mL/min)</td> <td></td> <td></td> </tr> </tbody> </table> <p>(a) Defined as samples collected 2-4 hours after dosing (b) Defined as samples collected 20-28 hours after dosing</p> <p>AF-atrial fibrillation, CR-CL-creatinine clearance, DVT-deep vein thrombosis, VTE-venous thromboembolism</p>	Dose	Indication	Peak Conc., ng/mL (a)	Trough Conc., ng/mL (b)	-----				2.5 mg	Acute coronary	46 (28-70)	17 (6-37)		twice daily syndrome			10 mg	VTE prevention	125 (91-196)	9 (1-38)		once daily after total hip replacement			15 mg	Stroke prevention	229 (178-313)	57 (18-136)		once daily in patients with AF (CR-CL 30-49 mL/min)			20 mg	DVT treatment	270 (189-419)	26 (6-87)		once daily			20 mg	Stroke prevention	249 (184-343)	44 (12-137)		once daily in patients with AF (CR-CL ≥50 mL/min)			<p>Chromogenic measurement at 405nm.</p>	<p>1. Package insert: Rivaroxaban: Diagnostics Stage. Revised December 2014 2. Moeck W, Stampfuss J, Kubitz D, Becka M: Clinical pharmacokinetic and pharmacodynamic profile of rivaroxaban. Clinical Pharmacokinetics 2014; 53(1):1-16 doi: 10.1007/s40262-013-0100-7 3. Bayer Pharma AG: Xarelto (rivaroxaban) Summary of Product Characteristics; 2013. Available at: www.ema.europa.eu/doc/en_GB/document_library/EPAR_-_Product_Information/human/000944/WC50005710_8.pdf 4. EINSTEIN Investigators, Basenachs R, Berkowitz SD, et al: Oral rivaroxaban for symptomatic venous thromboembolism. N Engl J Med 2010; 363:2499-510 5. EINSTEIN-PE Investigators, Buller HR, Prins MH, et al: Oral rivaroxaban for the treatment of symptomatic pulmonary embolism. N Engl J Med 2012; 366:1287-1297 6. Patel MR, Mahaffey KW, Garg J, et al: Rivaroxaban versus warfarin in nonvalvular atrial fibrillation. N Engl J Med 2011; 365:883-891 7. Sigal DM, Carruste JT, Connolly SJ, et al: Andexanet alfa for reversal of factor Xa inhibitor</p>	<p>25-500 ng/mL</p>	<p>25-500 ng/mL</p>
Dose	Indication	Peak Conc., ng/mL (a)	Trough Conc., ng/mL (b)																																																				

2.5 mg	Acute coronary	46 (28-70)	17 (6-37)																																																				
	twice daily syndrome																																																						
10 mg	VTE prevention	125 (91-196)	9 (1-38)																																																				
	once daily after total hip replacement																																																						
15 mg	Stroke prevention	229 (178-313)	57 (18-136)																																																				
	once daily in patients with AF (CR-CL 30-49 mL/min)																																																						
20 mg	DVT treatment	270 (189-419)	26 (6-87)																																																				
	once daily																																																						
20 mg	Stroke prevention	249 (184-343)	44 (12-137)																																																				
	once daily in patients with AF (CR-CL ≥50 mL/min)																																																						
<p>Anti Xa Heparin (Unfractionated)</p> <p>AXHEP UFH Heparin assay Unfractionated heparin</p>		<p>0.30-0.70 IU/mL</p>		<p>Chromogenic measurement at 405nm.</p>	<p>Chest, vol 119, issue 1, January 2001, pp. 64S-75S.</p>	<p>0.10-0.80 IU/mL</p>	<p>0.10-1.60 IU/mL</p>																																																
<p>Anti Xa LMWH (Enoxaparin) Random</p> <p>AXLMWHR Low molecular weight heparin LMWH Random</p>		<p>0.60-1.00 Anti-Xa IU/mL</p>		<p>Chromogenic measurement at 405nm.</p>	<p>Chest, vol 119, issue 1, January 2001, pp. 64S-75S.</p>	<p>0.10-1.60 Anti-Xa IU/mL</p>	<p>0.10-1.60 Anti-Xa IU/mL</p>																																																
<p>Anti Xa LMWH (Enoxaparin). *Exact Time Required* 4 Hr Post</p> <p>AXLMWHPK Low molecular weight heparin LMWH Post</p>		<p>0.60-1.00 Anti-Xa IU/mL</p>		<p>Chromogenic measurement at 405nm.</p>	<p>Chest, vol 119, issue 1, January 2001, pp. 64S-75S.</p>	<p>0.10-1.60 Anti-Xa IU/mL</p>	<p>0.10-1.60 Anti-Xa IU/mL</p>																																																
<p>Antithrombin III</p> <p>AT Antithrombin Activity Antithrombin Functional ATIII AT3</p>			<p>0-4 days: 39-87 % 5-29 days: 41-93 % 30-89 days: 48-108 % 90-179 days: 73-121 % 180-365 days: 84-124 % 1-5 years: 82-139 % 6-10 years: 90-131 % 11-16 years: 77-112 % 17+ years: 85-118 %</p>	<p>Chromogenic measurement at 405nm.</p>	<p>OSU/WMC Normal Range Study 02/2004; Blood, Vol 80, 1998-2005; Andrew, 1992; Amer. Jour. Ped. Hematol. Oncol, Vol 12, 95-104, Andrew, 1990</p>	<p>9-200 %</p>																																																	

Beta-2 Glycoprotein 1 Ab, IgG & IgM	BZGP1 BZGP1 IgG BZGP1 IgG BZGP1 IgM BZGP1 IgM		BZGP1G 0.0-20.0 CU BZGP1M 0.0-20.0 CU		Chemiluminescent two-step immunoassay.	Inova Quanta-Flash Package Insert Verified in House	BZGP1G: 10.0-500.0 CU BZGP1M: 10.0-200.0 CU	BZGP1G: 10.0-10,000.0 CU BZGP1M: 10.0-4,000.0 CU
D-Dimer, Quantitative	HSDD1 High sensitivity D-Dimer		<0.50 mcg/mL FEU		A suspension of latex microparticles, coated by covalent bonding with monoclonal antibodies specific for D-dimer, is mixed with the test plasma whose D-dimer level is to be assayed. Agglutination of the microparticles results via an antigen-antibody reaction, which induces an increase in turbidity of the reaction medium. This increase in turbidity is reflected by an increase in absorbance, the latter being measured photometrically at 540 nm. The increase in absorbance is a function of the D-dimer level present in the test sample.	OSLWMC Normal Range Study (08/2007)	0.27-4.00 mcg/mL FEU	0.27-20.00 mcg/mL FEU
DIC Workup	DIC,DCMXES Disseminated Intravascular Coagulation Workup Consumptive Coagulopathy Workup		PT 11.9-14.2 sec INR 0.9-1.1 PTT 24.0-34.3 sec TT 13.0-20.0 sec FIB 220-410 mg/dL D-DIMER <0.50 mcg/mL FEU		PT, PTT, Mixing Studies, FIB, TT: Mechanical Clot Detection D-Dimer: Optical measurement at 540 nm	See individual analytes	See individual analytes	See individual analytes
Factor II Activity	FA2 Factor 2 Activity Prothrombin Activity		0-4 days: 26-70 % Activity 5-29 days: 33-93 % Activity 30-89 days: 34-102 % Activity 90-179 days: 45-105 % Activity 180-365 days: 60-116 % Activity 1-5 years: 71-116 % Activity 6-10 years: 67-107 % Activity 11-16 years: 61-104 % Activity 17+ years: 60-150 % Activity		1 stage clotting assay.	Clinical Guide to Laboratory Tests, Tietz, 1995, Blood, Vol 80, 1998-2005, Andrew, 1992; Blood, Vol 70, 165-172, Andrew, 1987	3-500 % Activity	3-500 % Activity
Factor II Inhibitor	FA2IN Factor 2 Inhibitor Factor II Bethesda Titer Bethesda Titer		0.0 Bethesda units	Negative	Bethesda method using a 1 stage clotting assay.			≥0.0 Bethesda units Negative, Positive
Factor IX Activity	FA9 Factor 9 Activity Hemophilia B Christmas Factor		0-4 days: 15-91 % Activity 5-29 days: 15-91 % Activity 30-89 days: 21-81 % Activity 90-179 days: 21-113 % Activity 180-365 days: 36-136 % Activity 1-5 years: 67-108 % Activity 6-10 years: 63-89 % Activity 11-16 years: 59-122 % Activity 17+ years: 77-147 % Activity	<1% severe hemophilia 1-~5% moderate hemophilia 5-~40% mild hemophilia	1 stage clotting assay.	OSLWMC Normal Range Study 02/2004; Blood, Vol 80, 1998-2005, Andrew, 1992; Blood, Vol 70, 165-172, Andrew, 1987	1-500 % Activity	1-500 % Activity
Factor IX Inhibitor	FA9IN Factor 9 Inhibitor Factor IX Bethesda Titer Bethesda Titer		0.0 Bethesda units	Negative	Bethesda method using a 1 stage clotting assay.			≥0.0 Bethesda units Negative, Positive
Factor V Activity	FA5 Factor 5 Activity Labile Factor Proaccelerin		0-4 days: 34-108 % Activity 5-29 days: 45-145 % Activity 30-89 days: 62-134 % Activity 90-179 days: 40-132 % Activity 180-365 days: 55-127 % Activity 1-5 years: 79-127 % Activity 6-10 years: 63-116 % Activity 11-16 years: 55-99 % Activity 17+ years: 50-150 % Activity		1 stage clotting assay.	Clinical Guide to Laboratory Tests, Tietz, 1995, Blood, Vol 80, 1998-2005, Andrew, 1992; Blood, Vol 70, 165-172, Andrew, 1987	3-500 % Activity	3-500 % Activity
Factor V Inhibitor	FA5IN Factor 5 Inhibitor Factor V Bethesda Titer Bethesda Titer		0.0 Bethesda units	Negative	Bethesda method using a 1 stage clotting assay.			≥0.0 Bethesda units Negative, Positive
Factor VII Activity	FA7 Factor 7 Activity Stable Factor Proconvertin		0-4 days: 28-104 % Activity 5-29 days: 35-143 % Activity 30-89 days: 42-138 % Activity 90-179 days: 39-143 % Activity 180-365 days: 47-127 % Activity 1-5 years: 55-116 % Activity 6-10 years: 52-120 % Activity 11-16 years: 58-115 % Activity 17+ years: 65-135 % Activity		1 stage clotting assay.	Clinical Guide to Laboratory Tests, Tietz, 1995, Blood, Vol 80, 1998-2005, Andrew, 1992; Blood, Vol 70, 165-172, Andrew, 1987	3-1,000 % Activity	3-1,000 % Activity

Factor VII Inhibitor	FA7IN Factor 7 Inhibitor Factor VII Bethesda Titer Bethesda Titer	0.0 Bethesda units	Negative	Bethesda method using a 1 stage clotting assay.		≥0.0 Bethesda units Negative, Positive
Factor VIII Activity	FAS Factor 8 Activity Factor VIII Clotting Anti-Hemophilic Factor	0-4 days: 50-178 % Activity 5-29 days: 50-154 % Activity 30-89 days: 50-157 % Activity 90-179 days: 50-125 % Activity 180-365 days: 50-109 % Activity 1-5 years: 59-142 % Activity 6-10 years: 58-132 % Activity 11-16 years: 53-131 % Activity 17+ years: 50-200 % Activity	<1% severe hemophilia 1- <5% moderate hemophilia 5- <40% mild hemophilia	1 stage clotting assay.	Clinical Laboratory Reference Values. In: Laposata M. ed. <i>Laboratory Medicine: The Diagnosis of Disease in the Clinical Laboratory</i> . McGraw-Hill Education; 2014; Blood, Vol 80, 1998-2005, Andrew, 1992; Blood, Vol 70, 165-172, Andrew, 1987	1-500 % Activity 1-500 % Activity
Factor VIII Activity, Chromogenic	FASC Factor 8 Activity Factor VIII Clotting Factor VIII Chromogenic	0-4 days: 50-178 % Activity 5-29 days: 50-154 % Activity 30-89 days: 50-157 % Activity 90-179 days: 50-125 % Activity 180-365 days: 50-109 % Activity 1-5 years: 59-142 % Activity 6-10 years: 58-132 % Activity 11-16 years: 53-131 % Activity 17+ years: 50-200% % Activity	<1% severe hemophilia 1- <5% moderate hemophilia 5- <40% mild hemophilia	Two stage chromogenic assay, measurement at 405nm.	Clinical Laboratory Reference Values. In: Laposata M. ed. <i>Laboratory Medicine: The Diagnosis of Disease in the Clinical Laboratory</i> . McGraw-Hill Education; 2014; Blood, Vol 80, 1998-2005, Andrew, 1992; Blood, Vol 70, 165-172, Andrew, 1987	1-200 % Activity 1-200 % Activity
Factor VIII Inhibitor	FASIN Factor 8 Inhibitor Factor VIII Bethesda Titer Bethesda Titer	0.0 Bethesda units	Negative	Bethesda method using a 1 stage clotting assay.		≥0.0 Bethesda units Negative, Positive
Factor X Activity	FA10 Factor 10 Activity Stuart-Power Factor	0-4 days: 12-68 % Activity 5-29 days: 19-79 % Activity 30-89 days: 31-87 % Activity 90-179 days: 35-107 % Activity 180-365 days: 38-118 % Activity 1-5 years: 58-116 % Activity 6-10 years: 55-101 % Activity 11-16 years: 50-117 % Activity 17+ years: 60-139 % Activity	<1% severe hemophilia 1- <5% moderate hemophilia 5- <40% mild hemophilia	1 stage clotting assay.	Clinical Guide to Laboratory Tests, Tietz, 1995; Blood, Vol 80, 1998-2005, Andrew, 1992; Blood, Vol 70, 165-172, Andrew, 1987	3-500 % Activity 3-500 % Activity
Factor X Inhibitor	FA10IN Factor 10 Inhibitor Factor X Bethesda Titer Bethesda Titer	0.0 Bethesda units	Negative	Bethesda method using a 1 stage clotting assay.		≥0.0 Bethesda units Negative, Positive
Factor XI Activity	FA11 Factor 11 Activity Hemophilia C Plasma Thromboplastin Antecedent (PTA)	0-4 days: 10-66 % Activity 5-29 days: 23-87 % Activity 30-89 days: 27-79 % Activity 90-179 days: 41-97 % Activity 180-365 days: 49-134 % Activity 1-5 years: 56-150 % Activity 6-10 years: 52-120 % Activity 11-16 years: 50-97 % Activity 17+ years: 65-135 % Activity		1 stage clotting assay.	Clinical Guide to Laboratory Tests, Tietz, 1995; Blood, Vol 80, 1998-2005, Andrew, 1992; Blood, Vol 70, 165-172, Andrew, 1987	1-500 % Activity 1-500 % Activity
Factor XI Inhibitor	FA11IN Factor 11 Inhibitor Factor XI Bethesda Titer Bethesda Titer	0.0 Bethesda units	Negative	Bethesda method using a 1 stage clotting assay.		≥0.0 Bethesda units Negative, Positive
Factor XIII Activity	FA13 Factor 13 Screen Factor 13 Qualitative Factor XIII Qualitative Fibrin Stabilization Factor		Present	Clot solubility.		Absent, Present
Fibrinogen, Clotable	FIB Factor I Activity Fibrinogen Activity Clas Fibrinogen	220-410 mg/dL		Mechanical clot detection of dilute plasma in the presence of excess thrombin.	OSU Lab Normal Range Study (05/2003)	60-900 mg/dL 60-900 mg/dL
Fibrinogen, Obstetrical	FIB Fibrinogen, OB	OB 1st Trimester: 244-510 mg/dL OB 2nd Trimester: 291-538 mg/dL OB 3rd Trimester: 373-619 mg/dL	Fibrinogen levels may be altered by the normal physiologic changes of pregnancy and should be interpreted considering reference ranges specific to gestational age.	Mechanical clot detection of dilute plasma in the presence of excess thrombin.	Reference: Abbassi-Ghannoum M, Greer LG, Cunningham FG. Pregnancy and laboratory studies: a reference table for clinicians. <i>Obstet Gynecol</i> 2009; 114:1326.	60-900 mg/dL 60-900 mg/dL
Heparin Platelet Factor 4 (HIT Screen) with Reflex to (SRA) PF4R2P		PF4 IgG OD: <0.400 % Heparin Inhibition: <50		Enzyme-linked immunosorbent assay (ELISA), IgG	Immucor LIFE CODES® PF4 IgG Assay Package Insert	PF4 IgG OD: 0.000-3.000 PF4 IgG OD: 0.000-3.000

Lupus Anticoagulant	<p>PT 11.9-14.2 sec</p> <p>INR 0.9-1.1</p> <p>TT 13.0-20.0 sec</p> <p>PTT-LA ≤44.7 sec</p> <p>DRVVT SCREEN RATIO ≤1.28</p> <p>DRVVT NORMALIZED RATIO ≤1.25</p> <p>HEXAGONAL PL. NEUTRALIZATION ≤8.6 sec</p> <p>PT/TT PTT-LA/DRVVT LA Screen APS Screen Antiphospholipid Antibody Screen Siaclat LA DRVVT</p>			<p>Hexagonal PL. Neutralization: STA/LOT LA: Mechanical clot detection in the presence/absence of hexagonal ring phospholipid</p> <p>DRVVT: Mechanical clot detection in the presence of Diluted Russell's Viper Venom</p> <p>PTT-LA: Mechanical clot detection in the presence of cephalin and activator</p> <p>PT, TT, Mixing Studies: Mechanical clot detection</p>	OSUWMC Normal Range Study	<p>PT: See individual analyte</p> <p>TT: See individual analyte</p> <p>PTT-LA: 20.0-180.0 sec</p>	<p>PT: See individual analyte</p> <p>TT: See individual analyte</p> <p>PTT-LA: 20.0-180.0 sec</p>
Platelet Aggregation			<p>Aggregation w/ADP 5 umol/L: 67-92 % Aggregation</p> <p>Aggregation w/ADP 10.0 umol/L: 73-91 % Aggregation</p> <p>Aggregation w/Arach. Acid 0.5 mmol/L: 72-91 % Aggregation</p> <p>Aggregation w/Epinephrine 5 umol/L: 64-105 % Aggregation</p> <p>Aggregation w/Ristocetin 0.5 mg/mL: 0-6 % Aggregation</p> <p>Aggregation w/Ristocetin 1.25 mg/mL: 70-105 % Aggregation</p> <p>Aggregation w/Collagen 2 ug/mL: 80-96 % Aggregation</p> <p>Aggregation w/Collagen 5 ug/mL: 71-92 % Aggregation</p> <p>Aggregation w/TXA₂ 2.0 umol/L: 73-112 % Aggregation</p> <p>ATP Release w/ Collagen 2 ug/mL: 0.26-1.07 nmol</p> <p>ATP Release w/ Collagen 5 ug/mL: 0.49-1.32 nmol</p> <p>ATP Release w/Arach Acid 0.5 mmol/L: 0.37-0.90 nmol</p> <p>ATP Release w/ADP 5 umol/L: 0.28-0.93 nmol</p> <p>ATP Release w/ADP 10 umol/L: 0.44-1.19 nmol</p> <p>ATP Release w/Thrombin 1U: 0.36-0.99 nmol</p> <p>ATP Release w/Epinephrine 5 umol/L: 0.35-0.96 nmol</p>	<p>Born method of turbidimetric aggregation with simultaneous measurement of ATP release by platelet lumi-aggregometry.</p>	OSUWMC Normal Range Study (02-2023)		
Platelet Function Test	<p>PLATGG Pl. Agg. Pl. Aggregation</p>	<p>Collagen / Epi Closure Time: 73-172 sec</p> <p>Collagen / ADP Closure Time: 53-111 sec</p>		<p>Instrument PFA-100 closure time: the time measured from the start of the test until a platelet to close aperture after exposure to agonist.</p>	OSU Normal Range Study (07/2004)	<p>Collagen / Epi Closure Time: 31-300 sec</p> <p>Collagen / ADP Closure Time: 31-300 sec.</p>	<p>Collagen / Epi Closure Time: 31-300 sec</p> <p>Collagen / ADP Closure Time: 31-300 sec.</p>
Platelet P2Y12 Inhibition Test	<p>P2Y12 Clopidogrel Inhibition Ticagrelor Inhibition</p>	<p>194-418 PRU</p>	<p>Test results are reported in P2Y12 Reaction Units (PRU). The pre-ding Normal Reference Range is 194-418 PRU. PRU measures the extent of platelet aggregation in the presence of P2Y12 inhibitor drugs such as clopidogrel (Plavix), prasugrel (Effient), ticagrelor (Ticlid), and ticagrelor (Brilinta).</p>	<p>Whole blood platelet aggregometry based on the ability of activated platelets to bind fibrinogen. Fibrinogen-coated microplatelets aggregate in whole blood in proportion to the number of expressed platelet GP IIb/IIIa receptors.</p>	Accriva Diagnostics Verity Now P2Y12 Package Isent VN1009WEL	<p>0.999 PRU</p>	<p>0.999 PRU</p>
Protein C Activity	<p>PROTC Protein C Functional PC Functional PC Activity</p>	<p>0-4 days: 17-53 % Activity</p> <p>5-29 days: 20-64 % Activity</p> <p>30-89 days: 21-65 % Activity</p> <p>90-179 days: 28-80 % Activity</p> <p>180-365 days: 37-81 % Activity</p> <p>1-5 years: 56-150 % Activity</p> <p>6-10 years: 40-92 % Activity</p> <p>11-16 years: 55-111 % Activity</p> <p>17+ years: 72-220 % Activity</p>		<p>This clotting assay measures Protein C which is activated in the presence of the specific activator extracted from <i>Akfitnodon c. constrictus</i> venous. The resulting activated protein C inhibits the factors V and VIII, and thus prolongs the APTT of a system in which all the factors are present, constant and in excess (provided by the test system), except for the protein C which is derived from the sample being tested.</p>	OSUWMC Normal Range Study; Blood, Vol 80, 1998-2005, Andrew, 1992; Amer. Jour. Ped. Hematol. Oncol, Vol 12, 95-104, Andrew, 1990		<p>10-300 % Activity</p>
Protein S Activity	<p>PROTS Protein S Functional PS Functional PS Activity</p>	<p>0-4 days: 12-60 % Activity</p> <p>5-29 days: 22-78 % Activity</p> <p>30-89 days: 33-93 % Activity</p> <p>90-179 days: 54-118 % Activity</p> <p>180-365 days: 55-119 % Activity</p> <p>1-5 years: 53-173 % Activity</p> <p>6-10 years: 55-155 % Activity</p> <p>11-16 years: 65-138 % Activity</p> <p>17+ years: 50-168 % Activity</p>		<p>Clotting assay based on the cofactor activity of protein S which enhances the anticoagulant action of activated protein C. This enhancement is reflected by the prolongation of the clotting time of a system enriched with factor Va which is a physiological substrate for activated protein C.</p>	OSUWMC Normal Range Study; Blood, Vol 80, 1998-2005, Andrew, 1992; Amer. Jour. Ped. Hematol. Oncol, Vol 12, 95-104, Andrew, 1990		<p>10-300 % Activity</p>
Protime - INR	<p>PTI PT/INR Protime-INR</p>	<p>PT 11.9-14.2 sec</p> <p>INR 0.9-1.1</p>		<p>PT: Mechanical clot detection initiated by Calcium Thromboplastin</p> <p>INR: Calculation</p>	OSUWMC Normal Range Study	<p>PT: 7.0-109.0 sec</p> <p>INR: 0.5-15.2</p>	<p>PT: 7.0-109.0 sec</p> <p>INR: 0.5-15.2</p>
PT and PT Mixing Study	<p>PT PT Mixing PT Inhibitor Screen</p>	<p>PT 11.9-14.2 sec</p> <p>INR 0.9-1.1</p>		<p>PT: Mechanical clot detection initiated by Calcium Thromboplastin</p> <p>INR: Calculation</p> <p>PT Mixing Study: PT performed immediately subsequent to 1:1 mix with normal plasma pool.</p>		<p>PT: 7.0-109.0 sec</p> <p>INR: 0.5-15.2</p>	<p>PT: 7.0-109.0 sec</p> <p>INR: 0.5-15.2</p>
PT, INR, PTT	<p>PT/PTT PT/INR/PTT Protime-INR Activated Partial Thromboplastin Time</p>	<p>PT 11.9-14.2 sec</p> <p>INR 0.9-1.1</p>		<p>PT, PTT: Mechanical clot detection</p> <p>INR: Calculation</p>	See individual analytes	<p>See individual analytes</p>	<p>See individual analytes</p>
PT INR - Stroke	<p>PT/INR PT/INR - Stroke</p>	<p>PT 11.9-14.2 sec</p> <p>INR 0.9-1.1</p>		<p>PT, PTT: Mechanical clot detection</p> <p>INR: Calculation</p>	OSUWMC Normal Range Study	<p>PT: 7.0-109.0 sec</p> <p>INR: 0.5-15.2</p>	<p>PT: 7.0-109.0 sec</p> <p>INR: 0.5-15.2</p>
PTT	<p>PTT Partial Thromboplastin Time aPTT Activated Partial Thromboplastin Time</p>	<p>24.0-34.3 sec</p>		<p>Mechanical clot detection initiated by Calcium in the presence of cephalin (platelet substitute) and a particulate activator (alka).</p>	OSUWMC Normal Range Study	<p>20.0-180.0 sec</p>	<p>20.0-180.0 sec</p>
PTT with Mixing Study	<p>PTTMS PTT/Mixing aPTT/Mixing PTT Inhibitor Screen</p>	<p>24.0-34.3 sec</p>		<p>PTT: Mechanical clot detection initiated by Calcium in the presence of cephalin (platelet substitute) and a particulate activator (alka).</p> <p>PTT Mixing Study: PTT performed immediately subsequent to 1:1 mix with normal plasma pool.</p>		<p>20.0-180.0 sec</p>	<p>20.0-180.0 sec</p>
Ristocetin Cofactor	<p>RCF RCF Activity VonWillebrand Factor Activity</p>	<p>40-200 % Activity</p>		<p>Ability of patient plasma to agglutinate formalin-fixed normal platelets in the presence of ristocetin.</p>	OSUWMC Normal Range Study		<p>13-400 % Activity</p>

<p>ROTEM Main Lab: Heparin Panel</p>	<p>RTMHHP</p>	<p>INTEM CT: 122-208 sec INTEM CFT: 45-110 sec INTEM ALPHA: 70-81 degree INTEM A20: 51-72 mm INTEM MCF: 51-72 mm</p> <p>EXTEM CT: 43-82 sec EXTEM CFT: 48-127 sec EXTEM ALPHA: 65-80 degree EXTEM A20: 50-70 mm EXTEM MCF: 52-70 mm</p> <p>FBITEM A20: 7-21 mm FBITEM MCF: 7-21 mm</p> <p>Reference ranges are not available for all ROTEM components unless otherwise noted. Interpretation of ROTEM results must include all available parameters, patient clinical context and current therapy.</p> <p>Interpretation of the HEPTM results must include clinical correlation between the INTEM and the HEPTM data. All results should be interpreted carefully based on patient clinical context and current therapy.</p> <p>Interpretation of APTM results (fibrinolytic activity) must include correlation between the EXTEM and APTM data. All results should be interpreted carefully based on patient clinical context and current therapy.</p>	<p>Rotem Physician Advice</p>	<p>Rotational Viscoelastic Testing- Rotem Delta: The patented ROTEM technology is based on a fixed cylindrical cup and a permanently oscillating vertical axis.</p> <p>The axis is supported by a high precision ball bearing and oscillates to the left and to the right through an angle of 4.75°.</p>	<p>Normal Range Study 2013</p>		
<p>ROTEM Main Lab: Routine Panel</p>	<p>RTMRN</p>	<p>INTEM CT: 122-208 sec INTEM CFT: 45-110 sec INTEM ALPHA: 70-81 degree INTEM A20: 51-72 mm INTEM MCF: 51-72 mm</p> <p>INTEM ML: No reference range available; see comment INTEM LIR: No reference range available; see comment</p> <p>EXTEM CT: 43-82 sec EXTEM CFT: 48-127 sec EXTEM ALPHA: 65-80 degree EXTEM A20: 50-70 mm EXTEM MCF: 52-70 mm</p> <p>FBITEM A20: 7-21 mm FBITEM MCF: 7-21 mm</p> <p>Reference ranges are not available for all ROTEM components unless otherwise noted. Interpretation of ROTEM results must include all available parameters, patient clinical context and current therapy.</p> <p>Interpretation of APTM results (fibrinolytic activity) must include correlation between the EXTEM and APTM data. All results should be interpreted carefully based on patient clinical context and current therapy.</p>	<p>Rotem Physician Advice</p>	<p>Rotational Viscoelastic Testing- Rotem Delta: The patented ROTEM technology is based on a fixed cylindrical cup and a permanently oscillating vertical axis.</p> <p>The axis is supported by a high precision ball bearing and oscillates to the left and to the right through an angle of 4.75°.</p>	<p>Normal Range Study 2013</p>		
<p>ROTEM Main Lab: Trauma Panel</p>	<p>RTMTRN</p>	<p>EXTEM CT: 43-82sec EXTEM CFT: 48-127 sec EXTEM ALPHA: 65-80 degree EXTEM A20: 50-70 mm EXTEM MCF: 52-70 mm</p> <p>FBITEM A20: 7-21 mm FBITEM MCF: 7-21 mm</p> <p>Reference ranges are not available for all ROTEM components unless otherwise noted. Interpretation of ROTEM results must include all available parameters, patient clinical context and current therapy.</p> <p>Interpretation of APTM results (fibrinolytic activity) must include correlation between the EXTEM and APTM data. All results should be interpreted carefully based on patient clinical context and current therapy.</p>	<p>Rotem Physician Advice</p>	<p>Rotational Viscoelastic Testing- Rotem Delta: The patented ROTEM technology is based on a fixed cylindrical cup and a permanently oscillating vertical axis.</p> <p>The axis is supported by a high precision ball bearing and oscillates to the left and to the right through an angle of 4.75°.</p>	<p>Normal Range Study 2013</p>		
<p>Thrombin Time</p>	<p>TT</p>	<p>13.0-20.0 sec</p>		<p>Mechanical clot detection of undiluted plasma in the presence of a predetermined quantity of thrombin, resulting in a fibrin clot.</p>	<p>OSUWMC Normal Range Study</p>	<p>10.0-120.0 sec</p>	<p>10.0-120.0 sec</p>
<p>Thrombin Time w/ Mixing Studies</p>	<p>TTMS TT Mixing TT with Mixing</p>	<p>13.0-20.0 sec</p>		<p>TT: Mechanical clot detection of undiluted plasma in the presence of a predetermined quantity of thrombin, resulting in a fibrin clot.</p> <p>TT Mixing Study: Thrombin Time is performed subsequent to a 1:1 mix with normal plasma pool and also in the presence of protamine sulfate (if indicated).</p>	<p>OSUWMC Normal Range Study</p>	<p>10.0-120.0 sec</p>	<p>10.0-120.0 sec</p>
<p>Von Willebrand Battery Agn + Factor VIII</p>	<p>PTT (VW) F8 RCF: VWAG vWF Antigen and Factor VIII vWF Immunologic and Factor VIII vWF Antigen and Factor 9 vWF Immunologic and Factor 9</p>	<p>PTT 24.0-34.3 sec</p> <p>F8 0.4 days: 50-178 % Activity 5-29 days: 50-154 % Activity 30-89 days: 50-157 % Activity 90-179 days: 50-125 % Activity 180-365 days: 50-109 % Activity 1-5 years: 50-142 % Activity 6-10 years: 58-132 % Activity 11-16 years: 53-131 % Activity 17+ years: 50-200 % Activity</p> <p>RCF 40-200 % Activity</p> <p>VWFAG 0.4 days: 50-287 % 5-29 days: 50-254 % 30-89 days: 50-266 % 90-179 days: 50-206 % 180-365 days: 50-197 % 1-5 years: 60-129 % 6-10 years: 44-144 % 11-16 years: 46-153 % 17+ years: 50-180 %</p>		<p>PTT: Mechanical clot detection initiated by Calcium in the presence of cephalin (platelet substitute) and a particulate activator (silica).</p> <p>VWF Ag: Immuno-turbidimetric method (STA - Liatest VWF:Ag)</p> <p>Factor VIII: One Stage Clotting Assay</p> <p>Ristocetin Cofactor: Ability of patient plasma to agglutinate formalin-fixed normal platelets in the presence of ristocetin.</p>	<p>See individual analytes</p>	<p>See individual analytes</p>	<p>See individual analytes</p>

Von Willebrand Factor Ag	VWFAG vWF Antigen vWF Immunologic	0.4 days: 50-287 % 5-29 days: 50-254 % 30-89 days: 50-246 % 90-179 days: 50-206 % 180-365 days: 50-197 % 1-5 years: 60-120 % 6-10 years: 44-144 % 11-16 years: 46-153 % 17+ years: 50-180 %		Immuno-turbidimetric method (STA - Latex VWF-Ag)	OSU/WMC Normal Range Study; Blood, Vol 80, 1998-2005, Andrew, 1992; Blood, Vol 70, 165-172, Andrew, 1987	3-400 %	
Amikacin Level, Extended Interval	AMIKEL	Peak: 30-60 mcg/mL, Trough: <6 mcg/mL		Homogeneous particle-enhanced turbidimetric immunoassay.	OSU Pharmacy	3.0-50.0 mcg/mL	3.0-150.0 mcg/mL
Amikacin Level, Peak (Post Drug Level)	AMIKPK	Therapeutic Range: 30.0-60.0 mcg/mL		Homogeneous particle-enhanced turbidimetric immunoassay.	OSU Pharmacy	3.0-50.0 mcg/mL	3.0-150.0 mcg/mL
Amikacin Level, Random	AMIKR	Peak: 30-60 mcg/mL, Trough: <6 mcg/mL		Homogeneous particle-enhanced turbidimetric immunoassay.	OSU Pharmacy	3.0-50.0 mcg/mL	3.0-150.0 mcg/mL
Amikacin Level, Trough (Pre Drug Level)	AMIKTR	Therapeutic Range: <6.0 mcg/mL		Homogeneous particle-enhanced turbidimetric immunoassay.	OSU Pharmacy	3.0-50.0 mcg/mL	3.0-150.0 mcg/mL
Amphetamines, Urine, Confirmation	AMPFC Adderall Meth Speed	None Detected Cutoff concentrations by liquid chromatography-tandem mass spectrometry: Amphetamine: 25 ng/mL Methamphetamine: 25 ng/mL	Numeric results (positive) indicate the measured concentration of drug in the urine sample. Negative results may reflect the absence of drug or drug present below the assay's limit of quantitation. The detection of drugs in urine is dependent on several factors including the amount of drug ingested, time since exposure, individual metabolism, kidney and liver function, urine concentration, and the assay's limit of detection. Drug concentrations in urine cannot determine the dose, route of administration, or exact time of exposure.	Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)		CONFIRMATION (Amphetamine/Methamphetamine): 25-5,000 ng/mL	CONFIRMATION (Amphetamine/Methamphetamine): 25-25,000 ng/mL
Benzodiazepines, Urine, Confirmation		None Detected Cutoff concentrations by liquid chromatography-tandem mass spectrometry: 2-Hydroxyethylthiazepam: 50 ng/mL 7-Aminoclonazepam: 10 ng/mL 7-Aminothiazepam: 10 ng/mL Alpha-hydroxyalprazolam: 10 ng/mL Alpha-hydroxymidazolam: 10 ng/mL Alprazolam: 10 ng/mL Diazepam: 10 ng/mL Flurazepam: 10 ng/mL Midazolam: 10 ng/mL Nortriazepam: 10 ng/mL Temazepam: 10 ng/mL Clonazepam: 25 ng/mL Lorazepam: 25 ng/mL Oxazepam: 25 ng/mL	Numeric results (positive) indicate the measured concentration of drug in the urine sample. Negative results may reflect the absence of drug or drug present below the assay's limit of quantitation. The detection of drugs in urine is dependent on several factors including the amount of drug ingested, time since exposure, individual metabolism, kidney and liver function, urine concentration, and the assay's limit of detection. Drug concentrations in urine cannot determine the dose, route of administration, or exact time of exposure.	Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)		2-Hydroxyethylthiazepam: 50-1,750 ng/mL 7-Aminoclonazepam: 10-1,000 ng/mL 7-Aminothiazepam: 10-1,300 ng/mL Alpha-hydroxyalprazolam: 10-2,000 ng/mL Alpha-hydroxymidazolam: 10-1,750 ng/mL Alprazolam: 10-2,000 ng/mL Diazepam: 10-2,000 ng/mL Flurazepam: 10-1,000 ng/mL Midazolam: 10-2,000 ng/mL Nortriazepam: 10-2,000 ng/mL Temazepam: 10-2,000 ng/mL Clonazepam: 25-2,000 ng/mL Lorazepam: 25-2,000 ng/mL Oxazepam: 25-2,000 ng/mL	2-Hydroxyethylthiazepam: 50-1,750 ng/mL 7-Aminoclonazepam: 10-1,000 ng/mL 7-Aminothiazepam: 10-1,300 ng/mL Alpha-hydroxyalprazolam: 10-2,000 ng/mL Alpha-hydroxymidazolam: 10-1,750 ng/mL Alprazolam: 10-2,000 ng/mL Diazepam: 10-2,000 ng/mL Flurazepam: 10-1,000 ng/mL Midazolam: 10-2,000 ng/mL Nortriazepam: 10-2,000 ng/mL Temazepam: 10-2,000 ng/mL Clonazepam: 25-2,000 ng/mL Lorazepam: 25-2,000 ng/mL Oxazepam: 25-2,000 ng/mL
Buprenorphine and Norbuprenorphine, Urine, Confirmation	QBUPR Subutex Suboxone Buprenex	None Detected Cutoff concentrations by liquid chromatography-tandem mass spectrometry: Buprenorphine: 5.0 ng/mL Norbuprenorphine: 5.0 ng/mL	Numeric results (positive) indicate the measured concentration of drug in the urine sample. Negative results may reflect the absence of drug or drug present below the assay's limit of quantitation. The detection of drugs in urine is dependent on several factors including the amount of drug ingested, time since exposure, individual metabolism, kidney and liver function, urine concentration, and the assay's limit of detection. Drug concentrations in urine cannot determine the dose, route of administration, or exact time of exposure.	Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)		Buprenorphine: 5.0 -5,000.0 ng/mL Norbuprenorphine: 5.0 - 5,000.0 ng/mL	Buprenorphine: 5.0 -25,000.0 ng/mL Norbuprenorphine: 5.0 - 25,000.0 ng/mL
Carboxy THC, Urine Confirmation	THCCON Cannabinoids THC Marijuana Mrvy Jms	None Detected Cutoff concentrations by gas chromatography-tandem mass spectrometry: 9-Carboxy-11-Nor-Delta-THC: 5.0 ng/mL	Numeric results (positive) indicate the measured concentration of drug in the urine sample. Negative results may reflect the absence of drug or drug present below the assay's limit of quantitation. The detection of drugs in urine is dependent on several factors including the amount of drug ingested, time since exposure, individual metabolism, kidney and liver function, urine concentration, and the assay's limit of detection. Drug concentrations in urine cannot determine the dose, route of administration, or exact time of exposure.	Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)		5.0-500.0 ng/mL	5.0-500.0 ng/mL
Cocaine, Urine, Confirmation	COCCON Coke Crack	None Detected Cutoff concentrations by liquid chromatography-tandem mass spectrometry: Benzoylecgonine: 25 ng/mL Cocaine: 25 ng/mL	Numeric results (positive) indicate the measured concentration of drug in the urine sample. Negative results may reflect the absence of drug or drug present below the assay's limit of quantitation. The detection of drugs in urine is dependent on several factors including the amount of drug ingested, time since exposure, individual metabolism, kidney and liver function, urine concentration, and the assay's limit of detection. Drug concentrations in urine cannot determine the dose, route of administration, or exact time of exposure.	Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)		Benzoylecgonine: 25-5,000 ng/mL Cocaine: 25-5,000 ng/mL	Benzoylecgonine: 25-25,000 ng/mL Cocaine: 25-25,000 ng/mL
Cyclic Citrullinated Peptide Ab	ACCP Anti-CCP	<5.0 U/mL		Chemiluminescent microparticle immunoassay	Abbot	0.5-200.0 U/mL	0.5-1,200.0 U/mL
Cyclosporine Level, 2HR	CSANZ Sandimmune Neoral Gengraf	Therapeutic Range: 320-960 ng/mL			OSU Pharmacy	30-1,500 ng/mL	30-3,000 ng/mL
Cyclosporine Level, Random	CSANR Sandimmune Neoral Gengraf	Kidney/Pancreas Transplant: 0 to 3 months: 800-1000 ng/mL 3 to 12 months: 600-800 ng/mL >12 months: 400-600 ng/mL			OSU Pharmacy	30-1,500 ng/mL	30-3,000 ng/mL
Cyclosporine Level, Trough (Pre Drug Level)	CSANT Sandimmune Neoral Gengraf	Kidney/Pancreas Transplant: 0 to 3 months: 800-1000 ng/mL 3 to 12 months: 600-800 ng/mL >12 months: 400-600 ng/mL Aplastic anemia and stem cell transplant: 200-400			OSU Pharmacy	30-1,500 ng/mL	30-3,000 ng/mL
Ethanol (Alcohol), Urine	ALCOU	<10 mg/dL Negative	Detectable ethanol in urine indicates exposure to ethanol within the past 8-12 hours.	Beckman Coulter DxC700AU; EmP II Plus Ethyl Alcohol Assay		10-600 mg/dL	10-600 mg/dL
Ethylene Glycol, Blood, Screen with Reflex to Confirmation	EGU	Cutoff concentrations by gas chromatography: 10 mg/dL		Beckman Coulter DxC700AU; Catechem DiscrepPak Ethylene Glycol Reagent Kit Confirmation: Gas Chromatography Flame Ionization Detection (GC-FID)		CONFIRMATION: 10-250 mg/dL	SCREEN: Negative, Presumptive Positive, Confirmation to follow, CONFIRMATION: 10-250 mg/dL
Everolimus, Trough (Pre Drug Level)	EVERTR	Kidney/Pancreas Transplant: 0 to 3 months: 7-10 ng/mL 3 to 12 months: 5-8 ng/mL >12 months: 4-8 ng/mL		Particle enhanced turbidimetric immunoassay	OSU Pharmacy	2.0-20.0 ng/mL	2.0-40.0 ng/mL

Fentanyl, Urine, Confirmation		None Detected Cutoff concentrations by liquid chromatography-tandem mass spectrometry: Fentanyl: 2.5 ng/mL Norfentanyl: 2.5 ng/mL	Numeric results (positive) indicate the measured concentration of drug in the urine sample. Negative results may reflect the absence of drug or drug present below the assay's limit of quantitation. The detection of drugs in urine is dependent on several factors including the amount of drug ingested, time since exposure, individual metabolism, kidney and liver function, urine concentration, and the assay's limit of detection. Drug concentrations in urine cannot determine the dose, route of administration, or exact time of exposure.	Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)		Fentanyl: 2.5-5,000.0 ng/mL Norfentanyl: 2.5-5,000.0 ng/mL	Fentanyl: 2.5-25,000.0 ng/mL Norfentanyl: 2.5-25,000.0 ng/mL
Gentamicin Level, Extended Infusion	FENTO GENTEI Garamicin	Peak: 3-15.0 mcg/mL, Trough: ≤1.0 mcg/mL		Enzyme immunoassay		0.3-10.0 mcg/mL	0.3-20.0 mcg/mL
Gentamicin Level, Peak (Post Drug Level)	GENTP Garamicin	Therapeutic Range: 3.0-15.0 mcg/mL		Enzyme immunoassay	OSU Pharmacy	0.3-10.0 mcg/mL	0.3-20.0 mcg/mL
Gentamicin Level, Random	GENT Garamicin	Peak: 3-15.0 mcg/mL, Trough: ≤1.0 mcg/mL		Enzyme immunoassay		0.3-10.0 mcg/mL	0.3-20.0 mcg/mL
Gentamicin, Trough (Pre Drug Level)	GENTR Garamicin	0-365 days: Therapeutic Range: <1.6 mcg/mL 1+ years: Therapeutic Range: ≤1.0 mcg/mL		Enzyme immunoassay	OSU Pharmacy	0.3-10.0 mcg/mL	0.3-20.0 mcg/mL
Lidocaine Level	LIDOU	Therapeutic Range: 1.5-5.0 mcg/mL		Enzyme immunoassay	OSU Pharmacy	0.5-12.0 mcg/mL	0.5-36.0 mcg/mL
Methadone, Urine, Confirmation		None Detected Cutoff concentrations by liquid chromatography-tandem mass spectrometry: Methadone: 25 ng/mL EDDP Methadone: 25 ng/mL	Numeric results (positive) indicate the measured concentration of drug in the urine sample. Negative results may reflect the absence of drug or drug present below the assay's limit of quantitation. The detection of drugs in urine is dependent on several factors including the amount of drug ingested, time since exposure, individual metabolism, kidney and liver function, urine concentration, and the assay's limit of detection. Drug concentrations in urine cannot determine the dose, route of administration, or exact time of exposure.	Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)		Methadone: 25-5,000 ng/mL EDDP Methadone: 25-5,000 ng/mL	Methadone: 25-25,000 ng/mL EDDP Methadone: 25-25,000 ng/mL
	METHIMS						
Nicotine Screen Urine		None Detected Cutoff concentrations by immunoassay detection: 500 ng/mL	Positive results indicate recent exposure to cigarette smoke.	Beckman Coulter DxC700AU: Thermo Scientific DRP [®] Cotinine Assay			None Detected, Positive, Presumptive Positive. Confirmation to Follow.
	NICOTU						
Opioids, Urine, Confirmation		None Detected Cutoff concentrations by liquid chromatography-tandem mass spectrometry: 6-Monoacetylmorphine: 5.0 ng/mL Codeine: 25 ng/mL Hydrocodone: 25 ng/mL Hydromorphone: 25 ng/mL Morphine: 25 ng/mL Tramadol: 25 ng/mL	Numeric results (positive) indicate the measured concentration of drug in the urine sample. Negative results may reflect the absence of drug or drug present below the assay's limit of quantitation. The detection of drugs in urine is dependent on several factors including the amount of drug ingested, time since exposure, individual metabolism, kidney and liver function, urine concentration, and the assay's limit of detection. Drug concentrations in urine cannot determine the dose, route of administration, or exact time of exposure.	Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)		6-Monoacetylmorphine: 5.0-5,000.0 ng/mL Codeine: 25-5,000 ng/mL Hydrocodone: 25-5,000 ng/mL Hydromorphone: 25-5,000 ng/mL Morphine: 25-5,000 ng/mL	6-Monoacetylmorphine: 5.0-25,000.0 ng/mL Codeine: 25-25,000 ng/mL Hydrocodone: 25-25,000 ng/mL Hydromorphone: 25-25,000 ng/mL Morphine: 25-25,000 ng/mL
	OPICON						
Oxycodone, Urine, Confirmation		None Detected Cutoff concentrations by liquid chromatography-tandem mass spectrometry: Oxycodone: 25 ng/mL Oxymorphone: 25 ng/mL Noroxycodone: 25 ng/mL	Numeric results (positive) indicate the measured concentration of drug in the urine sample. Negative results may reflect the absence of drug or drug present below the assay's limit of quantitation. The detection of drugs in urine is dependent on several factors including the amount of drug ingested, time since exposure, individual metabolism, kidney and liver function, urine concentration, and the assay's limit of detection. Drug concentrations in urine cannot determine the dose, route of administration, or exact time of exposure.	Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)		Oxycodone: 25-5,000 ng/mL Oxymorphone: 25-5,000 ng/mL Noroxycodone: 25-5,000 ng/mL	Oxycodone: 25-25,000 ng/mL Oxymorphone: 25-25,000 ng/mL Noroxycodone: 25-25,000 ng/mL
	OXYCON						
Penicillins Level	PENTU	Intracranial Pressure Therapy: 30-40 ng/mL		Gas Chromatography (GC)	OSU Pharmacy	5-50 ng/mL	5-50 ng/mL
Phenoin Free Level	FFPN Diluent, Free	Therapeutic Range: 0.6-2.4 mcg/mL		Chemiluminescent microparticle immunoassay	OSU Pharmacy	0.5-40.0 mcg/mL	0.5-40.0 mcg/mL
Sivolums (Rapamycin) Level, Random	SIBOR	Bone Marrow Transplant: 3-12 Kidney/Pancreas Transplant: 0 to 3 months: 7-10; 3 to 12 months: 5-8; >12 months: 4-8		Chemiluminescent microparticle immunoassay	OSU Pharmacy	2.0-30.0 ng/mL	2.0-60.0 ng/mL
Sivolums (Rapamycin) Level, Trough (Pre Drug Level)	SIBOTR	Bone Marrow Transplant: 3-12 Kidney/Pancreas Transplant: 0 to 3 months: 7-10; 3 to 12 months: 5-8; >12 months: 4-8		Chemiluminescent microparticle immunoassay	OSU Pharmacy	2.0-30.0 ng/mL	2.0-60.0 ng/mL
Tacrolimus Level, Trough (Pre Drug Level)	TACRO Prograf	Bone Marrow Transplant: 5-15 Kidney/Pancreas Transplant: 0 to 3 months: 8-10; 3 to 12 months: 6-8; >12 months: 4-6		Chemiluminescent microparticle immunoassay	OSU Pharmacy	2.0-30.0 ng/mL	2.0-60.0 ng/mL
Tacrolimus, Random	TACROR Prograf	Bone Marrow Transplant: 5-15 Kidney/Pancreas Transplant: 0 to 3 months: 8-10; 3 to 12 months: 6-8; >12 months: 4-6		Chemiluminescent microparticle immunoassay	OSU Pharmacy	2.0-30.0 ng/mL	2.0-60.0 ng/mL
Tramadol, Urine		None Detected Cutoff concentrations by liquid chromatography-tandem mass spectrometry: ≤25 ng/mL	Numeric results (positive) indicate the measured concentration of drug in the urine sample. Negative results may reflect the absence of drug or drug present below the assay's limit of quantitation. The detection of drugs in urine is dependent on several factors including the amount of drug ingested, time since exposure, individual metabolism, kidney and liver function, urine concentration, and the assay's limit of detection. Drug concentrations in urine cannot determine the dose, route of administration, or exact time of exposure.	Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)		25-5,000 ng/mL	25-25,000 ng/mL
	TramU Utram ConZin						
Urine Drug Screen 10		None Detected Cutoff concentrations by immunoassay detection: Amphetamines/Methamphetamine: 500 ng/mL Barbiturates: 200 ng/mL Benzodiazepines: 200 ng/mL Buprenorphine: 5 ng/mL Cannabinoids: 50 ng/mL Cocaine: 150 ng/mL Fentanyl: 1 ng/mL Methadone: 300 ng/mL Opiates: 300 ng/mL Oxycodone: 100 ng/mL	Presumptive positive results indicate the presence of a compound within the specified class of drugs or a structurally related compound in the urine sample. Negative results may reflect the absence of drug or drug present below the assay's limit of detection. The detection of drugs in urine is dependent on several factors including the amount of drug ingested, time since exposure, individual metabolism, kidney and liver function, urine concentration, and the assay's limit of detection.	Beckman Coulter DxC700AU: Emi® II Plus Amphetamines Assay, Barbiturates Assay, Benzodiazepine Assay, Buprenorphine Assay, Cannabinoids Assay, Cocaine Metabolite Assay, Methadone Assay, and Opiates Assay. ARK™ Fentanyl II Assay, DR18 Oxycodone Assay.			Negative, Positive, Presumptive Positive. Confirmation to follow.
	UDRUG						
Urine Drug Screen 10 with Confirmation		None Detected Cutoff concentrations by immunoassay detection: Amphetamines/Methamphetamine: 500 ng/mL Barbiturates: 200 ng/mL Benzodiazepines: 200 ng/mL Buprenorphine: 5 ng/mL Cannabinoids: 50 ng/mL Cocaine: 150 ng/mL Fentanyl: 1 ng/mL Methadone: 300 ng/mL Opiates: 300 ng/mL Oxycodone: 100 ng/mL	Presumptive positive results indicate the presence of a compound within the specified class of drugs or a structurally related compound in the urine sample. Negative results may reflect the absence of drug or drug present below the assay's limit of quantitation. The detection of drugs in urine is dependent on several factors including the amount of drug ingested, time since exposure, individual metabolism, kidney and liver function, urine concentration, and the assay's limit of detection.	Screen: Beckman Coulter DxC700AU: Emi® II Plus Amphetamines Assay, Barbiturates Assay, Benzodiazepine Assay, Buprenorphine Assay, Cannabinoids Assay, Cocaine Metabolite Assay, Methadone Assay, and Opiates Assay. ARK™ Fentanyl II Assay, DR18 Oxycodone Assay. Confirmation: Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)			See individual reflex tests
	UPMSC						
Valproic Acid, Free	FVPA Depakote Divalproex Valproate Depakene	Therapeutic Range: 5-35 mcg/mL		Chemiluminescent microparticle immunoassay	OSU Pharmacy	2-150 mcg/mL	

Valproic Acid, Total	VPA Depakote Divalproex Valproate Depakene	Therapeutic Range: 50-120 mcg/mL		Chemiluminescent microparticle immunoassay		4-150 mcg/mL	4-750 mcg/mL
Volatile Alcohol Panel, Blood	VOLABU Alcohol Drug Methanol Ethanol Isopropanol Acetone Ethyl Alcohol Isopropyl Alcohol	METHANOL <10 mg/dL ETHYL ALCOHOL <10 mg/dL ACETONE <10 mg/dL ISOPROPANOL <10 mg/dL		Gas Chromatography (GC) Flame Ionization Detection (FID)		METHANOL: 10-400 mg/dL ETHYL ALCOHOL: 10-400 mg/dL ACETONE: 10-400 mg/dL ISOPROPANOL: 10-400 mg/dL	METHANOL: 10-400 mg/dL ETHYL ALCOHOL: 10-400 mg/dL ACETONE: 10-400 mg/dL ISOPROPANOL: 10-400 mg/dL
Adrenal Vein Sampling Pre Stimulation	CORT LAV1 CORT RAV1 CORT IVC1 CORT LAV2 CORT RAV2 CORT IVC2 CORT LAV3 CORT RAV3 CORT IVC3 ALDOLAV1 ALDO RAV1 ALDO IVC1 ALDOLAV2 ALDORAV2 ALDO IVC2 ALDOLAV3 ALDO RAV3 ALDO IVC3	CORTISOL LEFT AV1 3.09-22.40 mcg/dL CORTISOL RIGHT AV1 3.09-22.40 mcg/dL CORTISOL IVC1 3.09-22.40 mcg/dL CORTISOL LEFT AV2 3.09-22.40 mcg/dL CORTISOL RIGHT AV2 3.09-22.40 mcg/dL CORTISOL IVC2 3.09-22.40 mcg/dL CORTISOL LEFT AV3 3.09-22.40 mcg/dL CORTISOL RIGHT AV3 3.09-22.40 mcg/dL CORTISOL IVC3 3.09-22.40 mcg/dL ALDOSTERONE LEFT AV1 <23.20 ng/dL ALDOSTERONE RIGHT AV1 <23.20 ng/dL ALDOSTERONE IVC1 <23.20 ng/dL ALDOSTERONE LEFT AV2 <23.20 ng/dL ALDOSTERONE RIGHT AV2 <23.20 ng/dL ALDOSTERONE IVC2 <23.20 ng/dL ALDOSTERONE LEFT AV3 <23.20 ng/dL ALDOSTERONE RIGHT AV3 <23.20 ng/dL ALDOSTERONE IVC3 <23.20 ng/dL	Aldosterone Reference Range Plasma Upright: <35.30 ng/dL Plasma Supine: <23.60 ng/dL Serum Upright: <39.20 ng/dL Serum Supine: <23.20 ng/dL	Chemiluminescent immunoassay	See individual analytes	See individual analytes	See individual analytes
Adrenal Vein Sampling Post Stimulation		ALDOSTERONE 5 MIN LEFT AV No established reference range ALDOSTERONE 5 MIN RIGHT AV <23.20 ng/dL ALDOSTERONE 5 MIN IVC AV <23.20 ng/dL ALDOSTERONE 10 MIN LEFT AV <23.20 ng/dL ALDOSTERONE 10 MIN RIGHT AV <23.20 ng/dL ALDOSTERONE 10 MIN IVC AV <23.20 ng/dL ALDOSTERONE 15 MIN LEFT AV <23.20 ng/dL ALDOSTERONE 15 MIN RIGHT AV <23.20 ng/dL ALDOSTERONE 15 MIN IVC AV <23.20 ng/dL		Chemiluminescent immunoassay	See individual analytes	See individual analytes	See individual analytes

C-Peptide Tolerance (Part of the Insulin Glucose Tolerance Battery): 10 Minutes	CP10M	0.81-3.85 ng/mL			Two-site sandwich immunoassay chemiluminescent	Siemens C-Peptide Package Insert 10997742_EN Rev. 03, 2021-06	0.05-25.00 ng/mL	0.05-5,000.00 ng/mL	
C-Peptide Tolerance (Part of the Insulin Glucose Tolerance Battery): 30 Minutes	CP30M	0.81-3.85 ng/mL			Two-site sandwich immunoassay chemiluminescent	Siemens C-Peptide Package Insert 10997742_EN Rev. 03, 2021-06	0.05-25.00 ng/mL	0.05-5,000.00 ng/mL	
Cryptococcal Antigen	CRAG				Negative	Lateral flow assay	Package Insert	SCREEN: Negative, Positive TITER: 1:2, 1:5, 1:10, 1:20, 1:40, 1:80, 1:160, 1:320, 1:640, 1:1280, ≥12560	
Cryptococcus, Antigen CSF	CRAG				Negative	Lateral flow assay	Package Insert	SCREEN: Negative, Positive TITER: 1:2, 1:5, 1:10, 1:20, 1:40, 1:80, 1:160, 1:320, 1:640, 1:1280, ≥12560	
DHEA-Sulfate	DHEAS	25.90-460.20 ug/dL	34.50-568.90 ug/dL			Chemiluminescent immunoassay	Siemens DHEAS Package Insert 11200382_EN	3.00-1,500.00 ug/dL	3.00-3,000.00 ug/dL
ddDNA Antibody	DSDNAB				Negative	Multiplex flow immunoassay	Package Insert Textbook	QUANTITATIVE: 1-300 IU/mL	QUALITATIVE: Negative, Positive, Indeterminate QUANTITATIVE: 1-30,000 IU/mL
EBV VCA IgG Ab	EBVG				Negative	Multiplex flow immunoassay	Package Insert Textbook	Negative, Positive, Indeterminate	
EBV VCA IgG and IgM	EBV				EBVG: Negative EBVM: Negative	Multiplex flow immunoassay	See individual analytes	See individual analytes	
EBV VCA IgM Ab	EBVM				Negative	Multiplex flow immunoassay	Package Insert Textbook	Negative, Positive, Indeterminate	
ENA Battery (SSA, SSB, SM, RNP)	ENAB				RoIF: Negative SM: Negative SSA: Negative SSB: Negative	Multiplex flow immunoassay	See individual analytes	See individual analytes	
Free Hemoglobin, Plasma	PLASHGB	≤5.0 mg/dL				Photometric	HemoCue Operating Manual	30.0-2,100.0 mg/dL	≥30.0 mg/dL
GGPT, Qualitative	GGPD				Enzyme Activity Present	Visual Fluorescence	Package Insert	Enzyme Activity Present, Enzyme Activity Absent, Enzyme Activity Indeterminate	
Anti-Glomerular Basement Mem Ab	GBMAB				Negative	Multiplex flow immunoassay	Bio-Rad IFU Vasculitis Revision 665-0516f	Negative, Positive	
Growth Hormone	GHIR	<6.88 ng/mL	<1.23 ng/mL			Chemiluminescent immunoassay	Package Insert	0.05-1,600.00 ng/mL	0.05-1,600.00 ng/mL
Hemoglobin A1C	A1CB	4.7-5.6 %				High-Performance Liquid Chromatography (HPLC)	Textbook	3.5-15.0 %	3.5-15.0 %
Hemoglobin, Fetal	HF	<1.0 %				High-Performance Liquid Chromatography (HPLC)	Package Insert Textbook	1.0-40.0 %	1.0-40.0 %
Hemoglobinopathy Eval		<p>HEPB: HGBA ≥95.0%</p> <p>HGBA2 2.1-3.3 %</p> <p>HGBF ≤1.0 %</p> <p>HGBC ≤0.0%</p> <p>HGBS ≤0.0%</p> <p>HGB OTHER ≤0.0%</p> <p>GCBCHE: RBC 0-14 days: 4.12-5.74 Mtd. 15-30 days: 3.32-4.80 Mtd. 31-60 days: 2.93-3.97 Mtd. 61-179 days: 3.45-4.75 Mtd. 180 days-2 years: 3.97-5.01 Mtd. 2-5 years: 3.84-4.92 Mtd. 6-11 years: 3.90-4.96 Mtd. 12-17 years: 3.91-4.90 Mtd. 18+ years: 3.91-5.04 Mtd.</p> <p>HGB 0-7 days: 13.4-20.0 g/dL 8-14 days: 13.4-20.0 g/dL 15-30 days: 10.8-14.6 g/dL 31-60 days: 9.2-11.4 g/dL 61-179 days: 9.9-12.4 g/dL 180 days-2 years: 10.2-12.7 g/dL 2-5 years: 10.2-12.7 g/dL 6-11 years: 10.6-13.2 g/dL 12-17 years: 10.8-13.3 g/dL 18+ years: 11.4-15.2 g/dL</p> <p>MCH 0-14 days: 31.1-35.9 pg 15-30 days: 30.4-35.3 pg 31-60 days: 28.0-32.5 pg 61-179 days: 24.4-28.9 pg 180 days-2 years: 23.2-27.5 pg 2-5 years: 23.7-28.6 pg 6-11 years: 24.8-29.9 pg 12-17 years: 24.8-30.2 pg 18+ years: 25.9-33.9 pg</p> <p>MCV 0-14 days: 92.7-106.4 fL 15-30 days: 90.1-103.0 fL 31-60 days: 83.4-96.4 fL 61-179 days: 74.8-88.3 fL 180 days-2 years: 71.3-82.6 fL 2-5 years: 72.3-85.0 fL 6-11 years: 75.9-87.6 fL 12-17 years: 76.9-90.6 fL 18+ years: 79.6-97.7 fL</p> <p>RDW 0-14 days: 14.6-17.3 % 15-30 days: 14.4-16.2 % 31-60 days: 13.6-15.8 % 61-179 days: 12.2-14.3 % 180 days-2 years: 12.7-15.1 % 2-5 years: 12.4-14.9 % 6-11 years: 12.2-14.4 % 12-17 years: 12.3-14.6 % 18+ years: 10.8-14.9 %</p>	<p>HEPB: HGBA ≥95.0%</p> <p>HGBA2 2.1-3.3 %</p> <p>HGBF ≤1.0 %</p> <p>HGBC ≤0.0%</p> <p>HGBS ≤0.0%</p> <p>HGB OTHER ≤0.0%</p> <p>GCBCHE: RBC 0-14 days: 4.10-5.55 Mtd. 15-30 days: 3.16-4.63 Mtd. 31-60 days: 3.02-4.22 Mtd. 61-179 days: 3.43-4.80 Mtd. 180 days-2 years: 4.03-5.07 Mtd. 2-5 years: 3.89-4.97 Mtd. 6-11 years: 3.96-5.03 Mtd. 12-17 years: 4.03-5.29 Mtd. 18+ years: 4.38-5.83 Mtd.</p> <p>HGB 0-7 days: 13.9-19.1 g/dL 8-14 days: 13.9-19.1 g/dL 15-30 days: 10.0-15.3 g/dL 31-60 days: 8.9-12.7 g/dL 61-179 days: 9.6-12.4 g/dL 180 days-2 years: 10.1-12.5 g/dL 2-5 years: 10.2-12.7 g/dL 6-11 years: 10.3-13.4 g/dL 12-17 years: 11.0-14.5 g/dL 18+ years: 13.4-16.8 g/dL</p> <p>MCH 0-14 days: 31.3-35.6 pg 15-30 days: 29.3-34.1 pg 31-60 days: 27.8-32.0 pg 61-179 days: 24.4-28.9 pg 180 days-2 years: 22.7-27.2 pg 2-5 years: 23.7-28.3 pg 6-11 years: 24.9-29.2 pg 12-17 years: 25.2-30.2 pg 18+ years: 26.1-33.3 pg</p> <p>MCV 0-14 days: 91.3-103.1 fL 15-30 days: 89.4-99.7 fL 31-60 days: 84.2-94.2 fL 61-179 days: 74.1-87.5 fL 180 days-2 years: 69.5-81.7 fL 2-5 years: 71.3-84.0 fL 6-11 years: 74.4-86.1 fL 12-17 years: 76.7-89.2 fL 18+ years: 79.0-94.2 fL</p> <p>RDW 0-14 days: 14.8-17.0 % 15-30 days: 14.3-16.8 % 31-60 days: 13.8-16.1 % 61-179 days: 12.4-15.3 % 180 days-2 years: 12.9-15.6 % 2-5 years: 12.5-14.9 % 6-11 years: 12.3-14.1 % 12-17 years: 12.4-14.5 % 18+ years: 10.9-14.3 %</p>	<p>HEPB High-Performance Liquid Chromatography (HPLC)</p> <p>GCBCHE The Sysmex XN performs hematology analysis according to the hydrodynamic focusing (DF) detection, flow cytometry method (semiconductor laser), and sodium lauryl sulfate (SLS)-hemoglobin method.</p>	<p>HGBA2: 1.0-7.0 % HGBF: 1.0-40.0 % HGBS: See individual analyte HGBC: See individual analyte MCHC: See individual analyte MCV: See individual analyte RDW: See individual analyte</p>	<p>HGBA2: 1.0-7.0 % HGBF: 1.0-40.0 % HGBS: See individual analyte HGBC: See individual analyte MCHC: See individual analyte MCV: See individual analyte RDW: See individual analyte</p>			
Hepatitis A Ab, Total (IgG + IgM)	HAAGM				Negative	Competitive direct chemiluminescent immunoassay	Package Insert	Negative, Positive, Indeterminate	
Hepatitis A IgM Ab	HAABM				Negative	Two-step IgM capture immunoassay	Package Insert	Negative, Positive, Indeterminate	
Hepatitis B Core Ab, Total (IgG + IgM)	HBCGM				Negative	Two-wash antigen sandwich immunoassay	Package Insert	Negative, Positive, Indeterminate	
Hepatitis B Core IgM Ab	HBCBM				Negative	Two-step IgM capture immunoassay	Package Insert	Negative, Positive, Indeterminate	

Hepatitis B Surface Antibody	HBSAB			Negative	Sandwich direct chemiluminescent immunoassay	Package Insert	Negative, Positive
Hepatitis B Surface Antigen	HBSAG			Negative	Sandwich direct chemiluminescent immunoassay	Package Insert	Negative, Positive, Initially reactive, to be confirmed by neutralization
Hepatitis Battery, Acute	HEPB			HBSAG: Negative HBCGM: Negative HBsAb: Negative HCAb: Negative	Chemiluminescent immunoassay	See individual analytes	See individual analytes
Hepatitis Battery, Chronic	HEPB			HBSAG: Negative HBSAB: Negative HBCGM: Negative HCAb: Negative	Chemiluminescent immunoassay	See individual analytes	See individual analytes
Hepatitis Bc Antibody	HBEb			Non Reactive	Chemiluminescent immunoassay	Package Insert	Nonreactive, Reactive
Hepatitis Bc Antigen	HBEc			Negative	Two-step sandwich immunoassay	Package Insert	Negative, Positive
Hepatitis C Antibody	HCAB			Negative	Indirect sandwich immunoassay	Package Insert	Negative, Positive, Indeterminate
Hepatitis Infections	HEPPI			HBSAG: Negative HBCGM: Negative HBSAB: Negative	Chemiluminescent immunoassay	See individual analytes	See individual analytes
HIV 1 and 2 Antibody/24 Antigen	HIV12			Non Reactive	Two-step sandwich immunoassay	Package Insert	Non Reactive, Reactive
HIV-1/HIV-2 Differentiation	HIV12C			HIV-1: Non Reactive HIV-2: Non Reactive	Immunochromatographic assay	Package Insert	HIV-1: Non Reactive, Reactive, Indeterminate HIV-2: Non Reactive, Reactive, Indeterminate
HPV Human Papilloma Virus Genotyping w/o Pap Smear	HPVG			HPVG16: Negative HPVG18: Negative HPVG0: Negative The assay detects HPV 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, 68.	Polymerase chain reaction (PCR)	Package Insert	HPVG16: Negative, Positive HPVG18: Negative, Positive HPVG0: Negative, Positive
HPV Testing, Reflex to Cytology for All Positive Results	HPVGSF HPVCT HPV			HPVG16: Negative HPVG18: Negative HPVG0: Negative The assay detects HPV 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, 68.	Polymerase chain reaction (PCR)	Package Insert	HPVG16: Negative, Positive HPVG18: Negative, Positive HPVG0: Negative, Positive
HPV Testing, Reflex to Cytology for HPV Positive Other Category	HPVGOTH HPVCT HPV			HPVG16: Negative HPVG18: Negative HPVG0: Negative The assay detects HPV 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, 68.	Polymerase chain reaction (PCR)	Package Insert	HPVG16: Negative, Positive HPVG18: Negative, Positive HPVG0: Negative, Positive
HSV 1 and 2 IgG Antibody	HSVG12			HSVG1: Negative HSVG2: Negative	Multiplex flow immunoassay	Package Insert Textbook	HSVG1: Negative, Positive, Indeterminate HSVG2: Negative, Positive, Indeterminate
HSV 2 IgG Antibody	HSVG2			Negative	Multiplex flow immunoassay	Package Insert Textbook	Negative, Positive, Indeterminate
HSV IgM Antibody	HSVIM			Negative	ELISA manual method	Textbooks, HSV-1&2 IgG procedure, Package Insert	Negative, Positive, Equivocal
Immune Status Hepatitis Battery	HEP2			HBCGM: Negative HBSAB: Negative HCAB: Negative	Chemiluminescent immunoassay	See individual analytes	See individual analytes
Immunofixation, Serum	SIMFX SIMFXB Serum Protein Electrophoresis with Immunofixation and Serum Total Protein		MPRO1 ≤0.0 mg/dL MPRO2 ≤0.0 mg/dL MPRO3 ≤0.0 mg/dL MPRO4 ≤0.0 mg/dL		Capillary electrophoresis	Package Insert	MPRO1: ≤0.0 mg/dL MPRO2: ≤0.0 mg/dL MPRO3: ≤0.0 mg/dL MPRO4: ≤0.0 mg/dL
Immunoglobulin Free Chains			KFLC 3.9-26.0 mg/L LFLC 6.4-22.1 mg/L RATHO 0.51-1.72		Turbidimetry	2017 OSU Study	KFLC: 2.9-127.0 mg/L LFLC: 5.2-139.0 mg/L KFLC: 0.6-63,500.0 mg/L LFLC: 1.3-139,000.0 mg/L
Immunoglobulin IgG	IGG IgG, Total		\$165.3 IU/mL		Two-site sandwich direct chemiluminescent immunoassay	Reference Range Study 11.3.2016	2.5-3,000.0 IU/mL 2.5-3,000.0 IU/mL
Insulin	INSI		3.0-25.0 uIU/mL		Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05_2021-06	0.5-300.0 uIU/mL 0.5-1,500.0 uIU/mL
Insulin Glucose Tolerance Minus 5 Min	INSUL1		3.0-25.0 uIU/mL		Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05_2021-06	0.5-300.0 uIU/mL 0.5-1,500.0 uIU/mL
Insulin Glucose Tolerance 0 Minute	INSUL2		3.0-25.0 uIU/mL		Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05_2021-06	0.5-300.0 uIU/mL 0.5-1,500.0 uIU/mL
Insulin Glucose Tolerance 2 Minute	INSUL3		3.0-25.0 uIU/mL		Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05_2021-06	0.5-300.0 uIU/mL 0.5-1,500.0 uIU/mL
Insulin Glucose Tolerance 5 Minute	INSUL4		3.0-25.0 uIU/mL		Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05_2021-06	0.5-300.0 uIU/mL 0.5-1,500.0 uIU/mL
Insulin Glucose Tolerance 10 Minute	INSUL5		3.0-25.0 uIU/mL		Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05_2021-06	0.5-300.0 uIU/mL 0.5-1,500.0 uIU/mL
Insulin Glucose Tolerance 30 Minute	INSUL6		3.0-25.0 uIU/mL		Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05_2021-06	0.5-300.0 uIU/mL 0.5-1,500.0 uIU/mL
Insulin-Like Growth Factor 1	IGF1 Somatomedin C		All Reference Ranges		Chemiluminescent immunoassay	Package Insert	10.0-1,000.0 ng/mL 10.0-1,000.0 ng/mL
Intact PTH (Intraoperative)	RPTH Intact PTH Rapid Parathyroid Hormone		14.0-72.0 pg/mL		Two-site sandwich immunoassay	Atellica product insert (PTH) REV 4, 2020/11	6.3-2,000.0 pg/mL 6.3-160,000.0 pg/mL
JO-1 Antibody	JO1 RJT			Negative	Multiplex flow immunoassay	Package Insert Textbook	Negative, Positive
Lyme Antibody	LYME			Negative	Chemiluminescent immunoassay	Package Insert	Negative, Positive
Legionella Urinary Ag	LEGONU			Negative	Enzyme Immunoassay (EIA)	Package Insert	Negative, Positive
M Tuberculosis By Quantiferon	QFTB QFT M. Tuberculosis Antigen			Negative	Chemiluminescent immunoassay	Package Insert	MTB/BEG/ANT: Negative, Positive, Indeterminate MTB TB1-NIL: 0.00-10.00 IU/mL MTB TB2-NIL: 0.00-10.00 IU/mL MTB MITOGEN-NIL: 0.00-10.00 IU/mL MTB NIL: 0.00-10.00 IU/mL
Maternal Hepatitis B Surface Ag	MHBSAG			Negative	Sandwich direct chemiluminescent immunoassay	Package Insert	Negative, Positive, Initially reactive, to be confirmed by neutralization
Maternal Rubella, IgG/Hep B Ag	MHBSAG RUBAH			MHBSAG: Negative RUBAH: Positive	Multiplex flow immunoassay	See individual analytes	See individual analytes

Monoclonal Prot Infs, Urine 24 Hr			MPROU1 ≥9 mg/24hrs MPROU2 ≥9 mg/24hrs MPROU3 ≥9 mg/24hrs MPROU4 ≥9 mg/24hrs PROU4 40-225 mg/24 hrs		Electrophoresis	Package Insert			
	Urine Immunofixation - 24 Hour UMFXB URFO								
Monoclonal Prot Immuno, Serum		PROTEIN 0-30 days: 4.2-6.2 g/dL 31-182 days: 4.4-6.4 g/dL 183-365 days: 5.6-7.9 g/dL 1-18 years: 5.7-8.0 g/dL 19+ years: 6.4-8.3 g/dL ALBUMIN 3.5-5.0 g/dL ALPHA 1 0.2-0.4 g/dL ALPHA 2 0.5-1.0 g/dL BETA 0.5-1.1 g/dL GAMMA 0.6-1.5 g/dL HGA 0-30 days: 0-10 mg/dL 31-182 days: ≥42 mg/dL 183-365 days: 6-68 mg/dL 1-3 years: 15-111 mg/dL 4-6 years: 33-166 mg/dL 7-9 years: 28-180 mg/dL 10-12 years: 55-193 mg/dL 13-15 years: 62-241 mg/dL 16-18 years: 69-262 mg/dL 19-59 years: 66-413 mg/dL 60+ years: 90-410 mg/dL HGG 0-30 days: 162-872 mg/dL 31-182 days: 311-664 mg/dL 183-365 days: 325-647 mg/dL 1-3 years: 451-1,202 mg/dL 4-6 years: 560-1,319 mg/dL 7-9 years: 485-1,473 mg/dL 10-12 years: 586-1,609 mg/dL 13-15 years: 749-1,640 mg/dL 16-18 years: 804-1,817 mg/dL 19-59 years: 600-1,714 mg/dL 60+ years: 600-1,560 mg/dL HGM 0-29 days: 1-57 mg/dL 30-182 days: <128 mg/dL 183-365 days: <131 mg/dL 1-3 years: 35-184 mg/dL 4-6 years: 42-184 mg/dL 7-9 years: 30-165 mg/dL 10-12 years: 42-211 mg/dL 13-15 years: 34-226 mg/dL 16-18 years: 45-224 mg/dL 19-59 years: 45-281 mg/dL 60+ years: 30-360 mg/dL MPRO1 ≥0.0 mg/dL MPRO2 ≥0.0 mg/dL MPRO3 ≥0.0 mg/dL MPRO4 ≥0.0 mg/dL	PROTEIN 0-30 days: 4.1-6.3 g/dL 31-182 days: 4.7-6.7 g/dL 183-365 days: 5.5-7.0 g/dL 1-18 years: 5.7-8.0 g/dL 19+ years: 6.4-8.3 g/dL ALBUMIN 3.5-5.0 g/dL ALPHA 1 0.2-0.4 g/dL ALPHA 2 0.5-1.0 g/dL BETA 0.5-1.1 g/dL GAMMA 0.6-1.5 g/dL HGA 0-30 days: ≤11 mg/dL 31-182 days: ≤40 mg/dL 183-365 days: 1-82 mg/dL 1-3 years: 9-137 mg/dL 4-6 years: 44-187 mg/dL 7-9 years: 58-204 mg/dL 10-12 years: 46-218 mg/dL 13-15 years: 29-251 mg/dL 16-18 years: 68-259 mg/dL 19-59 years: 66-413 mg/dL 60+ years: 90-410 mg/dL HGG 0-30 days: 197-833 mg/dL 31-182 days: 140-533 mg/dL 183-365 days: 130-623 mg/dL 1-3 years: 413-1,112 mg/dL 4-6 years: 468-1,328 mg/dL 7-9 years: 582-1,441 mg/dL 10-12 years: 685-1,620 mg/dL 13-15 years: 590-1,600 mg/dL 16-18 years: 522-1,703 mg/dL 19-59 years: 600-1,714 mg/dL 60+ years: 600-1,560 mg/dL HGM 0-29 days: 0-65 mg/dL 30-182 days: 6-64 mg/dL 183-365 days: 15-117 mg/dL 1-3 years: 30-146 mg/dL 4-6 years: 31-151 mg/dL 7-9 years: 21-140 mg/dL 10-12 years: 27-151 mg/dL 13-15 years: 26-184 mg/dL 16-18 years: 28-179 mg/dL 19-59 years: 45-281 mg/dL 60+ years: 30-360 mg/dL MPRO1 ≥0.0 mg/dL MPRO2 ≥0.0 mg/dL MPRO3 ≥0.0 mg/dL MPRO4 ≥0.0 mg/dL		Capillary electrophoresis	See individual analytes	See individual analytes	See individual analytes	
Mumps, IgG Ab, Immune Status		SERUM TP PSE QDM SIMFX			Positive	Package Insert		Negative, Positive, Indeterminate	
Myeloperoxidase Antibodies		MUMPSB			Negative	Textbook		Negative, Positive	
		MPO							
Post-Distal Splenic Artery		DISTSPART30 DISTSPART60 DISTSPART90 DISTSPART120 DISTPART180	3.0-25.0 uIU/mL			Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05, 2021-06	0.5-300.0 uIU/mL	0.5-1,500.0 uIU/mL
Post Gastrohepatic Artery		GASTART30 GASTART60 GASTART90 GASTART120 GASTART180	3.0-25.0 uIU/mL			Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05, 2021-06	0.5-300.0 uIU/mL	0.5-1,500.0 uIU/mL
Post Miscellaneous Artery		MISC30 MISC60 MISC90 MISC120 MISC180	3.0-25.0 uIU/mL			Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05, 2021-06	0.5-300.0 uIU/mL	0.5-1,500.0 uIU/mL
Post Proximal Splenic Artery		PROXSPART30 PROXSPART60 PROXSPART90 PROXSPART120 PROXSPART180	3.0-25.0 uIU/mL			Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05, 2021-06	0.5-300.0 uIU/mL	0.5-1,500.0 uIU/mL
Post Superior Mesenteric Artery		SUPMART30 SUPMART60 SUPMART90 SUPMART120 SUPMART180	3.0-25.0 uIU/mL			Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05, 2021-06	0.5-300.0 uIU/mL	0.5-1,500.0 uIU/mL
Pre-Distal Splenic Artery		DISTSPARTB1 DISTSPARTB2	3.0-25.0 uIU/mL			Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05, 2021-06	0.5-300.0 uIU/mL	0.5-1,500.0 uIU/mL

Pre Gastrohepatic Artery	GASTARTB1 GASTARTB2	3.0-25.0 uIU/mL			Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05, 2021-06	0.5-300.0 uIU/mL	0.5-1,500.0 uIU/mL
Pre Miscellaneous Artery	MISC B1 MISC B2	3.0-25.0 uIU/mL			Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05, 2021-06	0.5-300.0 uIU/mL	0.5-1,500.0 uIU/mL
Pre-Proximal Splenic Artery	PROXSPARTB1 PROXSPARTB2	3.0-25.0 uIU/mL			Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05, 2021-06	0.5-300.0 uIU/mL	0.5-1,500.0 uIU/mL
Pre Superior Mesenteric Artery	SUPMARTB1 SUPMARTB2	3.0-25.0 uIU/mL			Two-site sandwich chemiluminescent immunoassay	Siemens Insulin Package Insert 10997752_EN Rev. 05, 2021-06	0.5-300.0 uIU/mL	0.5-1,500.0 uIU/mL
Protein Electrophoresis Serum, with Reflex	SERIM TP SPE Serum Electrophoresis PSE Serum Protein Electrophoresis with Reflex to Immunofixation and serum total protein	PROTEIN 0-30 days: 4.2-6.2 g/dL 31-182 days: 4.4-6.6 g/dL 183-365 days: 5.6-7.9 g/dL 1-18 years: 5.7-8.0 g/dL 19+ years: 6.4-8.3 g/dL ALBUMIN 3.5-5.0 g/dL ALPHA 1 0.2-0.4 g/dL ALPHA 2 0.5-1.0 g/dL BETA 0.5-1.1 g/dL GAMMA 0.6-1.5 g/dL	PROTEIN 0-30 days: 4.1-6.3 g/dL 31-182 days: 4.7-6.7 g/dL 183-365 days: 5.5-7.0 g/dL 1-18 years: 5.7-8.0 g/dL 19+ years: 6.4-8.3 g/dL ALBUMIN 3.5-5.0 g/dL ALPHA 1 0.2-0.4 g/dL ALPHA 2 0.5-1.0 g/dL BETA 0.5-1.1 g/dL GAMMA 0.6-1.5 g/dL		Capillary electrophoresis	Package Insert	PROTEIN: 3.0-12.0 g/dL ALBUMIN: <=0.0 g/dL ALPHA 1: <=0.0 g/dL ALPHA 2: <=0.0 g/dL BETA: <=0.0 g/dL GAMMA: <=0.0 g/dL	PROTEIN: 3.0-24.0 g/dL ALBUMIN: <=0.0 g/dL ALPHA 1: <=0.0 g/dL ALPHA 2: <=0.0 g/dL BETA: <=0.0 g/dL GAMMA: <=0.0 g/dL
PTH Intact	PTH Parathyroid Hormone	14.0-72.0 pg/mL			Two-site sandwich immunoassay	Atellica product insert (PTH) REV 4, 2020/11	6.3-2,000.0 pg/mL	6.3-160,000.0 pg/mL
Renin	RENIN Direct	0.39 years: 4.2-52.2 pg/mL 40+ years: 3.6-81.6 pg/mL			Chemiluminescent immunoassay	Package Insert	2.1-300.0 pg/mL	2.1-3,000.0 pg/mL
Ribonuclease P Antibody	RIBOPT			Negative	Multiplex flow immunoassay	Package Insert Textbook		Negative, Positive
RNP Antibody	RNP			Negative	Multiplex flow immunoassay	Package Insert Textbook		Negative, Positive
RPR	Rapid Plasma Reagin RPRC			Non Reactive	Macroscopic nontreponemal flocculation	Package Insert		SCREEN: Non Reactive, Reactive TITER: 1:1, 1:2, 1:4, 1:8, 1:16, 1:32, 1:64, 1:128, 1:256, 1:512, 1:1024, <1:2048
RPR with Titer	RPR T			Non Reactive	Macroscopic nontreponemal flocculation	Package Insert		SCREEN: Non Reactive, Reactive TITER: 1:1, 1:2, 1:4, 1:8, 1:16, 1:32, 1:64, 1:128, 1:256, 1:512, 1:1024, <1:2048
Rubella IgG With Reflex To IgM	RUBAGR			Positive	Multiplex flow immunoassay	Package Insert Textbook		Negative, Positive, Indeterminate
Rubella Immune Status IgG Antibody	RUBAB			Positive	Multiplex flow immunoassay	Package Insert Textbook		Negative, Positive, Indeterminate
Rubella IgG Ab (Immune Status)	RUBOIB			Positive	Multiplex flow immunoassay	BioPlex 2200 MMRV IgG Procedure March 2010		Negative, Positive, Indeterminate
Sex Hormone Binding Globulin	SHBG	18.00-144.00 nmol/L		10.00-57.00 nmol/L	Sandwich immunoassay	Siemens IMMULITE 2000 SHBG (PIL2KSH-20, 2018-03-15)	1.60-180.00 nmol/L	1.60-360.00 nmol/L
Sm Antibody	SMA SMT			Negative	Multiplex flow immunoassay	Package Insert Textbook		Negative, Positive
Sm/RNP Antibody	SMBNPT			Negative	Multiplex flow immunoassay	Package Insert Textbook		Negative, Positive
SS-A/Ro Antibody	SSAT			Negative	Multiplex flow immunoassay	Package Insert Textbook		Negative, Positive
SS-B/LA Antibody	SSBT			Negative	Multiplex flow immunoassay	Package Insert Textbook		Negative, Positive
Syphilis Ab w/ Reflex RPR	SYPHIT Syphilis IgG/IgM Antibody			Non Reactive	Direct sandwich assay	Package Insert		SYPHIG: Non Reactive, Reactive, Equivocal RPR SCREEN: Non Reactive, Reactive RPR TITER: 1:1, 1:2, 1:4, 1:8, 1:16, 1:32, 1:64, 1:128, 1:256, 1:512, 1:1024, <1:2048
Toxoplasma IgG Antibody	TOXOG			Negative	Sandwich direct chemiluminescent immunoassay	Package Insert		Negative, Positive, Equivocal
Urine Immunofixation, Random	UIMPXR Monoclonal Prot Immfx, Urine - Random UIMPXR				Electrophoresis	Package Insert		
Varicella IgG Ab (Immune Status)	VZVSI			Positive	Multiplex flow immunoassay	Package Insert Textbook		Negative, Positive, Indeterminate
Vitamin D (25-Hydroxy, Total)	D25OH Vitamin D, Total	30.0-100.0 ng/mL			Chemiluminescent immunoassay	Package Insert	4.0-150.0 ng/mL	4.0-150.0 ng/mL
Vitamin D (1,25 Dihydroxy)	D125	20.0-79.0 pg/mL			In vitro chemiluminescent immunoassay	Package Insert	5.0-180.0 pg/mL	5.0-540.0 pg/mL
ABO/RH(D) Typing	ABORH Blood Type		A, B, O, AB Rh Negative, Rh Positive		Agglutination			
ABORH - Not Valid for Transfusion	ABORHID ABOD Blood Type		A, B, O, AB Rh Negative, Rh Positive		Agglutination			
Antibody Titer	ABTTT		Negative		Agglutination	Albimmunization Committee		
Antigen Testing	AGD Antigen typing				Agglutination			
Baby Type and DAT (Direct Antiglobulin Test)	HEELS Feedback Evaluation		ABORH A, B, O, AB Rh Negative, Rh Positive DAT Negative		Agglutination			
Cold Agglutinin Titer	COLD		Negative		Agglutination			
Cord Blood Evaluation	CORDB CORDH		ABORH A, B, O, AB Rh Negative, Rh Positive DAT Negative		Agglutination			

Cordocentesis				Agglutination			
	CORDO						
Direct Antiglobulin Test (DAT)			Negative	Agglutination			
	DATO DAT						
Rhoig Evaluation				Agglutination			
	RHEV Rhoigm Evaluation						
Transfusion Reaction Battery				Agglutination			
	TEEN						
Type and Screen			ABORH A, B, O, AB Rh Negative, Rh Positive ANTIBODY SCREEN Negative	Agglutination			
	XM						
Type and Screen - Not for Transfusion			ABORH A, B, O, AB Rh Negative, Rh Positive ANTIBODY SCREEN Negative	Agglutination			
	TYSC						
Type and Screen - Preadmission			ABORH A, B, O, AB Rh Negative, Rh Positive ANTIBODY SCREEN Negative	Agglutination			
	XMPO						
BMT Panel 1				Flow cytometry immunophenotyping			
	BMT1R						
CD19 CAR T				Flow cytometry immunophenotyping	Known value, circulating CAR T can only be found in infused patients		CD3PERCENTIL: 0.1-100.0 % CD3PERTOT: 0.1-100.0 % CAR19PERTL: 0.1-100.0 % CAR19PME: 0.1-100.0 % CAR19PERCD3: 0.1-100.0 %
	CAR19T CD19 probe CAR 19 CAR T19 Chimeric antigen receptor T detection						
CD3EN				Flow cytometry immunophenotyping			
	CD3EN CD3 Enumeration						
Helper/Suppressor Quant			CD4CD3 DUAL (T HELPER) 0-179 days: 50.0-57.0 % 180-365 days: 49.0-55.0 % 1-2 years: 46.0-51.0 % 2-3 years: 38.0-46.0 % 3+ years: 32.0-62.0 % CD4CD3 DUAL (T HELPER) 266-2,213 ABS/mm3 CD4CD3 DUAL (T SUPPRESSOR) 11.0-40.0 CD4CD3 DUAL (T SUPPRESSOR) 91-1,428 ABS/mm3	Flow cytometry immunophenotyping	OSU Flow Lab established		
	H5FB H5F CD4CD8 T helper T suppressor						

<p>Immuno Recount Panel Flow Cytometry</p> <p>BRP FBRPD FBRP</p>	<p>CD3+ 59.0-92.0% CD19+ 2.0-23.0% CD56+<CD3+ 3.0-25.0% CD13+HLA DR: No established reference range CD13+HLA DR+: No established reference range CD3<CD8+: 0.0-2.9% CD3<CD14+: 0.0-0.5% CD3<HLA ADH+: 0.0-0.4% CD3<CD69+HLA ADH+: 0.0-0.2% CD3<CD14+HLA DR+: 0.0-0.1% CD3<CD69+CD14+: 0.0-0.1% CD3<CD69+CD14+HLA ADH+: 0.0-0.0% CD3<CD80+: 0.0-0.2% CD3<CD86+: 0.0-0.5% CD3<CD88+: 59.8-86.9% CD3<CD86+: 59.7-86.3% CD19<CD80+: 0.0-0.4% CD19<CD86+: 0.0-0.5% CD19<CD88+: 3.6-20.5% CD19<CD86+: 2.9-20.7% CD19<CD88<CD86+: 2.2-19.8% CD19<CD88+CD86+: 0.0-0.0% CD3<CD88<CD86+: 59.3-85.8% CD3<CD88+CD86+: 0.0-0.0% CD4<CD45RA+: 6.4-26.5% CD4<CD45RO+: 12.2-41.8% CD8<CD45RA+: 14.3-45.8% CD8<CD45RO+: 5.2-31.8% CD4<CD45RA+<CD27+: 0.0-0.8% CD4+<CD27+<CD45RA+<CD45RO+: 0.0-12.7% CD4<CD45RA+<CD27+: 4.6-30.2% CD4<CD45RO+<CD27+: 0.8-8.0% CD4<CD45RA+<CD27+: 9.9-36.1% CD8<CD45RA+<CD27+: 0.0-15.9% CD8+<CD27+<CD45RA+<CD45RO+: 0.0-0.1% CD8<CD45RA+<CD27+: 0.8-18.3% CD8<CD45RO+<CD27+: 0.4-7.7% CD8<CD45RO<CD27+: 0.0-12.9% CD4<CD45RA+<CD28+: 2.5-26.1% CD4+<CD28+<CD45RA+<CD45RO+: 0.0-0.7% CD4<CD45RA+<CD29+: 0.3-5.9% CD4<CD45RO+<CD29+: 0.9-8.4% CD4<CD45RO+<CD29+: 8.3-24.6% CD8<CD45RA+<CD29+: 0.0-14.2% CD8+<CD29+<CD45RA+<CD45RO+: 0.0-0.1% CD8<CD45RA+<CD29+: 2.6-18.3% CD8<CD45RO+<CD29+: 0.0-2.0% CD8<CD45RO+<CD29+: 0.0-14.4% CD4<CD31+: No established reference range CD8<CD31+: No established reference range CD3<CD49+: No established reference range CD4<CD49+: No established reference range CD8<CD49+: No established reference range CD4+<CD193+: 0.0-0.1% CD4+<CD294+: 0.0-0.8% CD4+<CD183+: 6.7-30.5% CD4+<CD193+<CD294+: 0.0-0.0% CD4+<CD193+<CD183+: 0.0-0.0% CD4+<CD294+<CD183+: 0.0-0.0% CD4+<CD194+: 0.0-27.8% CD4+<CD193+<CD194+: 0.0-0.0% CD4+<CD294+<CD194+: 0.0-0.2% CD4+<CD193+<CD294+<CD194+: 8.9-32.5% CD4<CD25+: 0.0-3.2% CD8<CD127+: 30.6-53.9% CD8<CD25+: 0.0-0.1% CD8<CD127+: 6.7-24.0% CD8<CD127<CD25+: 6.3-24.3% CD8<CD25<CD127+: 0.0-0.0% CD4<CD25<CD127+: 0.0-2.1% CD4<CD25<CD127+: 0.0-1.4% CD3+<AlphaBeta+: 47.3-80.4% CD3+<GammaDelta+: 0.0-7.3% CD3<CD158b+: 0.0-4.4% CD3<CD56+<CD80+: 0.0-0.2% CD3<CD69+<CD158b+: 0.0-0.1% CD3<CD56+<CD158b+: 0.0-7.1% CD3<CD56+<CD80+: 0.0-1.1% CD3<CD56+<CD69+<CD158b+: 0.0-0.2% CD3<CD159a+: 0.0-6.8% CD3<CD107a+b+: 0.0-0.3% CD3<CD159a+<CD107a+b+: 0.0-0.0% CD3<CD56+<CD107a+b+: 0.0-0.0% CD3<CD56+<CD159a+: 0.0-1.6% CD3<CD56+<CD159a+<CD107a+b+: 0.0-0.0% CD3<CD314+: 10.3-36.2% CD3<CD56+<CD314+: 0.0-2.1% CD3<CD56+<CD314+: 0.0-0.2% CD3<CD63+<CD314+: 0.0-0.5% CD3<CD56+<CD314+: 0.0-14.8% CD3<CD56+<CD63+<CD314+: 0.0-0.2% CD16<CD56<CD3<CD117+: 2.2-23.0% CD16<CD56<CD3<CD117+: 0.8-5.2% CD16<CD56<CD3<CD117+: 0.2-1.1% CD16<CD56<CD3<CD117+: 0.0-17.9% CD16<CD56<CD3<CD117+: 57.2-82.8% CD16<CD56<CD3<CD117+: 0.0-1.6% CD16<CD56<CD3<CD117+: 0.0-7.0% CD16<CD56<CD3+<CD117+: 0.0-2.1% CD16<CD56<CD3<CD117+: 0.0-0.2% CD16<CD56<CD3<CD117+: 0.0-0.0%</p>	<p>CD3+ 590-4416 ABS/mm3 CD19+ 17-750 ABS/mm3 CD56+<CD3+ 25-893 ABS/mm3 CD13+HLA DR: No established reference range CD13+HLA DR+: No established reference range CD3<CD8+: No established reference range CD3<CD14+: 0-95 ABS/mm3 CD3<HLA ADH+: 0-139 ABS/mm3 CD3<CD69+HLA ADH+: 0-4 ABS/mm3 CD3<CD14+HLA DR+: 0-2 ABS/mm3 CD3<CD69+CD14+: 0-2 ABS/mm3 CD3<CD69+CD14+HLA ADH+: 0-0 ABS/mm3 CD3<CD80+: 0-5 ABS/mm3 CD3<CD86+: 0-10 ABS/mm3 CD3<CD88+: 607-2519 ABS/mm3 CD3<CD86+: 607-2517 ABS/mm3 CD19<CD80+: 0-11 ABS/mm3 CD19<CD86+: 0-14 ABS/mm3 CD19<CD88+: 121-557 ABS/mm3 CD19<CD86+: 113-554 ABS/mm3 CD19<CD88<CD86+: 114-554 ABS/mm3 CD19<CD88+CD86+: 0-0 ABS/mm3 CD3<CD88<CD86+: 603-2502 ABS/mm3 CD3<CD88+CD86+: 0-0 ABS/mm3 CD4<CD45RA+: 58-842 ABS/mm3 CD4<CD45RO+: 146-1011 ABS/mm3 CD8<CD45RA+: 187-1078 ABS/mm3 CD8<CD45RO+: 132-816 ABS/mm3 CD4<CD45RA+<CD27+: 0-18 ABS/mm3 CD4+<CD27+<CD45RA+<CD45RO+: 0-25 ABS/mm3 CD4<CD45RA+<CD27+: 0-783 ABS/mm3 CD4<CD45RO+<CD27+: 0-163 ABS/mm3 CD4<CD45RA+<CD27+: 9-991 ABS/mm3 CD8<CD45RA+<CD27+: 0-341 ABS/mm3 CD8+<CD27+<CD45RA+<CD45RO+: 0-1 ABS/mm3 CD8<CD45RA+<CD27+: 0-426 ABS/mm3 CD8<CD45RO+<CD27+: 0-122 ABS/mm3 CD8<CD45RO<CD27+: 0-291 ABS/mm3 CD4<CD45RA+<CD28+: 131-666 ABS/mm3 CD4+<CD28+<CD45RA+<CD45RO+: 0-17 ABS/mm3 CD4<CD45RA+<CD29+: 0-145 ABS/mm3 CD4<CD45RO+<CD29+: 2-201 ABS/mm3 CD4<CD45RO+<CD29+: 8-388 ABS/mm3 CD8<CD45RA+<CD29+: 0-331 ABS/mm3 CD8+<CD29+<CD45RA+<CD45RO+: 0-2 ABS/mm3 CD8<CD45RA+<CD29+: 30-80 ABS/mm3 CD8<CD45RO+<CD29+: 0-56 ABS/mm3 CD8<CD45RO+<CD29+: 0-343 ABS/mm3 CD4<CD31+: No established reference range CD8<CD31+: No established reference range CD3<CD49+: No established reference range CD4<CD49+: No established reference range CD8<CD49+: No established reference range CD4+<CD193+: 0-3 ABS/mm3 CD4+<CD294+: 0-23 ABS/mm3 CD4+<CD183+: 64-496 ABS/mm3 CD4+<CD193+<CD294+: 0-0 ABS/mm3 CD4+<CD193+<CD183+: 0-0 ABS/mm3 CD4+<CD193+<CD294+<CD183+: 0-0 ABS/mm3 CD4+<CD194+: 0-560 ABS/mm3 CD4+<CD193+<CD194+: 0-0 ABS/mm3 CD4+<CD294+<CD194+: 0-4 ABS/mm3 CD4+<CD193+<CD294+<CD194+: 64-836 ABS/mm3 CD4<CD25+: 0-68 ABS/mm3 CD8<CD127+: 316-1467 ABS/mm3 CD8<CD25+: 0-1 ABS/mm3 CD8<CD127+: 55-624 ABS/mm3 CD8<CD127<CD25+: 50-616 ABS/mm3 CD8<CD25<CD127+: 0-0 ABS/mm3 CD4<CD25<CD127+: 0-43 ABS/mm3 CD4<CD25<CD127+: 0-30 ABS/mm3 CD3+<AlphaBeta+: 50-2158 ABS/mm3 CD3+<GammaDelta+: 0-135 ABS/mm3 CD3<CD158b+: 0-89 ABS/mm3 CD3<CD56+<CD80+: 0-5 ABS/mm3 CD3<CD69+<CD158b+: No established reference range CD3<CD56+<CD158b+: 0-178 ABS/mm3 CD3<CD56+<CD80+: 0-24 ABS/mm3 CD3<CD56+<CD69+<CD158b+: 0-6 ABS/mm3 CD3<CD159a+: 0-170 ABS/mm3 CD3<CD107a+b+: 0-6 ABS/mm3 CD3<CD159a+<CD107a+b+: 0-0 ABS/mm3 CD3<CD56+<CD107a+b+: 0-0 ABS/mm3 CD3<CD56+<CD159a+: 0-0 ABS/mm3 CD3<CD56+<CD159a+<CD107a+b+: 0-0 ABS/mm3 CD3<CD314+: 89-929 ABS/mm3 CD3<CD56+<CD314+: 0-41 ABS/mm3 CD3<CD56+<CD314+: 0-6 ABS/mm3 CD3<CD63+<CD314+: 0-11 ABS/mm3 CD3<CD56+<CD314+: 0-356 ABS/mm3 CD3<CD56+<CD63+<CD314+: 0-48 ABS/mm3 CD16<CD56<CD3<CD117+: 1-419 ABS/mm3 CD16<CD56<CD3<CD117+: 0-122 ABS/mm3 CD16<CD56<CD3<CD117+: 3-24 ABS/mm3 CD16<CD56<CD3<CD117+: 0-446 ABS/mm3 CD16<CD56<CD3<CD117+: 579-2-50 ABS/mm3 CD16<CD56<CD3<CD117+: 0-36 ABS/mm3 CD16<CD56<CD3+<CD117+: 0-157 ABS/mm3 CD16<CD56<CD3+<CD117+: 0-49 ABS/mm3 CD16<CD56<CD3<CD117+: 0-0 ABS/mm3 CD16<CD56<CD3<CD117+: 0-3 ABS/mm3 CD16<CD56<CD3<CD117+: 0-0 ABS/mm3</p>	<p>Flow cytometry immunophenotyping</p>	<p>OSU Flow Lab established</p>
--	--	--	---	---------------------------------

Immunodeficiency Batt Plus NK	PBDB2	<p>AlphaBeta 47.0-80.0%</p> <p>CD13 No established reference range</p> <p>CD14 No established reference range</p> <p>CD19 2.0-21.0%</p> <p>CD2 70.0-92.0%</p> <p>CD20 2.0-21.0%</p> <p>CD3 59.0-92.0%</p> <p>CD4 32.0-62.0%</p> <p>CD45RA No established reference range</p> <p>CD45RO No established reference range</p> <p>CD56/16 3.0-25.0%</p> <p>CD8 11.0-40.0%</p> <p>GammaDelta 0.0-7.0%</p> <p>HLA DR No established reference range</p>	<p>AlphaBeta 550-2,158 ABS/mm³</p> <p>CD13 No established reference range</p> <p>CD14 No established reference range</p> <p>CD19 17-750 ABS/mm³</p> <p>CD2 581-3,284 ABS/mm³</p> <p>CD20 17-750 ABS/mm³</p> <p>CD3 490-3,284 ABS/mm³</p> <p>CD4 266-2,213 ABS/mm³</p> <p>CD45RA No established reference range</p> <p>CD45RO No established reference range</p> <p>CD56/16 25-893 ABS/mm³</p> <p>CD8 91-1,425 ABS/mm³</p> <p>GammaDelta 0-135 ABS/mm³</p> <p>HLA DR No established reference range</p>		Flow cytometry immunophenotyping			
Immunophenotyping	GPP	<p>CD10 No established reference range</p> <p>CD13 No established reference range</p> <p>CD14 No established reference range</p> <p>CD19 2.0-21.0%</p> <p>CD2 70.0-92.0%</p> <p>CD20 2.0-21.0%</p> <p>CD23 No established reference range</p> <p>CD3 59.0-92.0%</p> <p>CD4 32.0-62.0%</p> <p>CD5 No established reference range</p> <p>CD56/16 3.0-25.0%</p> <p>CD7 No established reference range</p> <p>CD8 11.0-40.0%</p> <p>HLA DR No established reference range</p> <p>KAPPA No established reference range</p> <p>LAMBDA No established reference range</p>	<p>CD10 No established reference range</p> <p>CD13 No established reference range</p> <p>CD14 No established reference range</p> <p>CD19 17-750 ABS/mm³</p> <p>CD2 581-3,284 ABS/mm³</p> <p>CD20 17-750 ABS/mm³</p> <p>CD23 No established reference range</p> <p>CD3 490-3,284 ABS/mm³</p> <p>CD4 266-2,213 ABS/mm³</p> <p>CD5 No established reference range</p> <p>CD56/16 25-893 ABS/mm³</p> <p>CD7 No established reference range</p> <p>CD8 91-1,425 ABS/mm³</p> <p>HLA DR No established reference range</p> <p>KAPPA No established reference range</p> <p>LAMBDA No established reference range</p>		Flow cytometry immunophenotyping			

TCRV Beta by Flow Cytometry, Blood

TKIT
TCRVB
TCR
PRTCR

CD1314
No established reference range

CD19
2.0-21.0%
17-750 ABS/mm³

CD2
70.0-92.0%
581-3,284 ABS/mm³

CD56
No established reference range

CD3
59.0-92.0%
490-3,284 ABS/mm³

CD4
32.0-62.0%
266-2,213 ABS/mm³

CD5616
3.0-25.0%
25-893 ABS/mm³

CD7
No established reference range

CD8
11.0-40.0%
91-1,425 ABS/mm³

Vb1
1.89-11.70%

Vb11
0.25-5.11%

Vb12
1.00-4.76%

Vb13.1
1.62-8.16%

Vb13.2
0.80-5.28%

Vb13.6
0.84-8.80%

Vb14
1.33-8.03%

Vb16
0.42-1.90%

Vb17
2.28-12.61%

Vb18
0.58-5.23%

Vb2
4.03-23.48%

Vb20
0.00-9.73%

Vb21.3
1.08-5.97%

Vb22
1.99-9.89%

Vb23
0.26-4.76%

Vb3
0.52-15.71%

Vb4
0.79-3.26%

Vb5.1
3.19-14.93%

Vb5.2
0.49-4.98%

Vb5.3
0.37-2.98%

Vb7.1
0.64-20.01%

Vb7.2
0.05-5.45%

Vb8
2.26-29.47%

Vb9
1.10-9.30%

Flow cytometry immunophenotyping

Beckman Coulter-verified

Transplant Battery		<p>CD4/CD3 DUAL (T HELPER)% 0-179 days: 50.0-57.0 % 180-365 days: 49.0-55.0 % 1-2 years: 46.0-51.0 % 2-3 years: 38.0-46.0 % 3+ years: 32.0-62.0 %</p> <p>CD4/CD3 DUAL (T HELPER) 266-2,213 ABS/mm³</p> <p>CD4/CD3 DUAL (T SUPPRESSOR)% 11.0-40.0</p> <p>CD4/CD3 DUAL (T SUPPRESSOR) 91-1,428 ABS/mm³</p>		Flow cytometry immunophenotyping	OSU Flow Lab established	
Transplant Battery Plus	TRBAT	<p>CD19 2.0-21.0% 17-750 ABS/mm³</p> <p>CD2 70.0-92.0% 581-3,284 ABS/mm³</p> <p>CD20 2.0-21.0% 17-750 ABS/mm³</p> <p>CD3 59.0-92.0% 490-3,284 ABS/mm³</p> <p>CD4 32.0-62.0% 266-2,213 ABS/mm³</p> <p>CD8 11.0-40.0% 91-1,428 ABS/mm³</p>		Flow cytometry immunophenotyping	OSU Flow Lab established	
Acid Fast Culture	PHSOT Transplant battery plus CD20	No acid fast bacillus seen.	Organisms growing in pure culture are identified to the species level whenever possible.	Culture Susceptibility (if appropriate)		No acid Fast Bacillus Seen Acid Fast Bacilli, Few Acid Fast Bacilli, Rare Acid Fast Bacilli, Moderate Acid Fast Bacilli, Heavy
Acid Fast Culture, Tissue	AFBCX	No acid fast bacillus seen.	Organisms growing in pure culture are identified to the species level whenever possible.	Culture		No acid Fast Bacillus Seen Acid Fast Bacilli, Few Acid Fast Bacilli, Rare Acid Fast Bacilli, Moderate Acid Fast Bacilli, Heavy
Acinetobacter Culture, MICU Only	BKR AFB TISS	No growth	Organisms growing in pure culture are identified to the species level whenever possible.	Surveillance Culture		
Actinomyces, Screen	ACTNSC			Gram Stain		
Anaerobe Culture	ANACX	No growth	Isolation of anaerobes in significant numbers from well-collected specimens including blood, aspirates, tissues or sterile body fluid, indicates infection with the identified organisms.	Culture Susceptibility (if appropriate)		
Atypical Bacterial Pneumonia, PCR	Atypical Bac	Not Detected (for all targets)	Results should be used in conjunction with other clinical and laboratory findings. This result does not rule out co-infections with pathogens that are not screened for by the Respiratory Panel (RP). This RP assay was performed using a Film Array multiplex nucleic acid assay.	Multiplex polymerase chain reaction (PCR)	Package Insert	MINCP: Not Detected, Detected MINBP: Not Detected, Detected MINBPP: Not Detected, Detected MINMP: Not Detected, Detected
Bacterial Culture and Direct Smear, Lesion, Tissue, Device	General Cult Residue culture and smear	No growth	Isolation of anaerobes in significant numbers from well-collected specimens indicates infection with the identified organisms.	Smear Culture Susceptibility (if appropriate)	Validation	
Bacterial Vaginosis Panel	BVAG BV	Negative	This test was performed using Transcription Mediated Amplification for the detection of ribosomal RNA from bacteria associated with bacterial vaginosis.	Transcription-mediated amplification (TMA)	Package Insert Clinical Data	Negative, Positive
BAL Consult	BAL CONSULT					
Beta Strep, Vaginal Screen	GBSP Group B Streptococcus Testing by PCR	Negative	This test was performed using a real-time PCR assay. Results should be interpreted in conjunction with other clinical and laboratory findings.	Concentration in LIM broth for > 18 hours followed by real-time polymerase chain reaction (PCR) testing for Group B Streptococcus (GBS) DNA sequence	Validation	SCREEN: Negative for Group B Strep, Positive for Group B Strep PCR: Negative, Positive
Beta Strep, Vaginal Screen, Reflex Susceptibility for Penicillin Allergy	GBS REFLEX GBS Streptococcus galactosae	Negative	This test was performed using a real-time PCR assay. Results should be interpreted in conjunction with other clinical and laboratory findings.	Concentration in LIM broth for > 18 hours followed by real-time polymerase chain reaction (PCR) testing for Group B Streptococcus (GBS) DNA sequence	Validation	SCREEN: Negative for Group B Strep, Positive for Group B Strep PCR: Negative, Positive
BK Virus DNA PCR, Quant, Urine	BKBUI BKBP	<500 copies/mL	This test was performed using a real time PCR assay. The dynamic range for this assay is 500-10,000,000,000 copies/mL.	Real-time polymerase chain reaction (RT-PCR)	Validation	BKBUT: 500-10,000,000,000 copies/mL. CAT: >10,000,000 copies/mL.
BK Virus DNA Qn, PCR, Plasma	BKBP	<500 copies/mL	This test was performed using a real time PCR assay. The dynamic range for this assay is 500-10,000,000,000 copies/mL.	Real-time polymerase chain reaction (RT-PCR)	Validation	BKBPT: 500-5,000,000,000 copies/mL. CAT: >5,000,000 copies/mL.
Blood Culture	BLDCX	No growth	Isolation of anaerobes in significant numbers from well-collected specimens indicates infection with the identified organisms.	Culture Susceptibility (if appropriate)		
Blood Culture, AFB, Mycobacteria	AFBBCX Acid fast blood culture	Negative A final negative report will be issued after 42 days of incubation.	A positive result may support the diagnosis of mycobacteremia.	Culture Susceptibility (if appropriate)		
Blood Culture, Pediatric	PEDI	No growth	Isolation of anaerobes in significant numbers from well-collected specimens indicates infection with the identified organisms.	Culture Susceptibility (if appropriate)		
Body Fluid Culture and Direct Smear	BFCX Sterile fluid culture	No growth	Isolation of anaerobes in significant numbers from well-collected specimens including sterile body fluids, indicates infection with the identified organisms.	Culture Susceptibility (if appropriate)		
C. difficile 2 Step	CD2STEP	C DIFFICILE BY PCR Negative		Polymerase chain reaction (PCR)	Package Insert	C DIFFICILE BY PCR: Negative, Positive C DIFFICILE TOXIN A/B: Negative, Positive.
Candida Aairn Screen by PCR	CAALPCR	Not Detected	This test was performed using a real-time PCR assay.	Real-time polymerase chain reaction (RT-PCR)	Package Insert Literature	Not Detected, Detected
Candida/Trichomonas Panel	CXTV	Not Detected (for all targets)	This test was performed using Transcription Mediated Amplification for the detection of ribosomal RNA from microorganisms associated with vulvovaginal candidiasis and trichomoniasis.	Transcription-mediated amplification (TMA)	Package Insert Clinical Data	Not Detected, Detected, Invalid
CAPD Fluid Bacterial Culture	BFCX	No growth	Isolation of anaerobes in significant numbers from well-collected specimens including sterile body fluids, indicates infection with the identified organisms.	Culture Susceptibility (if appropriate)		
Chlamydia & Gonorrhea Amplified Probe	BFCX Chlam and Co Chlamydia trachomatis & Neisseria gonorrhoeae NAAT Testing	Not Detected (for all targets)		Transcription-mediated amplification (TMA)	Package Insert	Not Detected, Detected

Chlamydia Amplified Probe	CT	Not Detected A negative test result for Chlamydia trachomatis does not preclude the possibility of infection. Results should be considered in conjunction with other clinical and laboratory findings.	This test was performed using Transcription Mediated Amplification for the detection of Chlamydia trachomatis and/or Neisseria gonorrhoeae nucleic acid. This assay is not intended for the evaluation of suspected sexual abuse or other medico-legal indications.	Transcription-mediated amplification (TMA)	Package Insert	Not Detected, Detected
CMV by PCR, Quantitative, Blood	CMVPCR CMV Viral Load CMV PCR	<50 IU/mL	This test was performed using a real time CMV PCR assay. The dynamic range for this assay is 50-156,000,000 IU/mL. Results should be interpreted in conjunction with other clinical and laboratory findings.	Real-time polymerase chain reaction (RT-PCR)	Literature History	50-156,000,000 IU/mL 50-156,000,000 IU/mL
EBV by PCR, Quantitative, Blood	EBVPCR EBV Viral Load EBV PCR	<1,000 IU/mL	This test was performed using a real time PCR assay. The dynamic range for this assay is 1,000-5,000,000 IU/mL. A result <1000 IU/mL does not rule out the presence of EBV DNA in quantities below the sensitivity of this assay.	Real-time polymerase chain reaction (RT-PCR)	Validation	1,000-5,000,000 IU/mL 1,000-5,000,000 IU/mL
EBV Rapid PCR, CSF Only	EBVFLD EBV PCR EBV CSF	Not Detected	This test was performed using a real time PCR assay. The limit of detection for this assay is 1,165 IU/mL. This assay is only to be used for patients with a clinical history and symptoms consistent with EBV infections, and should be interpreted in conjunction with other clinical findings. This test should not be used to screen asymptomatic patients.	Real-time polymerase chain reaction (RT-PCR)	Validation	Not Detected, Detected: <10,000 IU/mL, Detected: ≥10,000 IU/mL
Fungus Culture	FUNCX	No growth	Positive cultures of yeast and filamentous fungi are reported with the organism identification.	Culture Susceptibility (if appropriate)		
Fungus Culture (Skin, Hair, Nails)	SINCX	No growth	Positive cultures of yeast and filamentous fungi are reported with the organism identification.	Culture		
Fungus Smear	CALCOFLUOR Calcofluor White fluorescent stain			Smear		
Genital Culture, Bacterial	GENCX Vaginal Culture Cervical Culture Urethral Culture	No growth	Isolation of anaerobes in significant numbers from well-collected specimens indicates infection with the identified organisms.	Smear (if appropriate) Culture Susceptibility (if appropriate)		
Gonorrhea Amplified Probe	NG Chlamydia trachomatis & Neisseria gonorrhoeae NAAT Testing	Not Detected A negative test result for Chlamydia trachomatis does not preclude the possibility of infection. Results should be considered in conjunction with other clinical and laboratory findings.	This test was performed using Transcription Mediated Amplification for the detection of Chlamydia trachomatis and/or Neisseria gonorrhoeae nucleic acid. This assay is not intended for the evaluation of suspected sexual abuse or other medico-legal indications.	Transcription-mediated amplification (TMA)	Package Insert	Not Detected, Detected
H. Pylori Urea Breath Test	UBT UBT for H. pylori			Infrared Spectrophotometry	Package Insert	Negative, Positive
Hepatitis B DNA	HBDNAB HBV Viral Load	HBV QUANT <10 IU/mL HBV QUANT LOG <1.00 IU/mL	This test was performed using a real time HBV PCR assay. The dynamic range for this assay is 10-1,000,000,000 IU/mL (1.00-9.00 Logs). Results should be interpreted in conjunction with other clinical and laboratory findings.	Real-time polymerase chain reaction (RT-PCR)	Package Insert Validation	HBDNA: 10-1,000,000,000 IU/mL HBDNA: ≥10 IU/mL HBDNAL: 1.00-9.00 IU/mL
Hepatitis C by PCR, Quant	HEPCQP HCV Viral Load	HCV QUANT <12 IU/mL HCV QUANT LOG <1.08 IU/mL	This test was performed using a real time PCR assay. The dynamic range for this assay is 12-100,000,000 IU/mL.	Real-time polymerase chain reaction (RT-PCR)	Package Insert Validation	HEPCQ: 12-100,000,000 IU/mL HEPCQL: 1.08-8.00 IU/mL HEPCQ: ≥12 IU/mL HEPCQL: 1.08-8.00 IU/mL
HIV Viral Load RNA PCR Quant	HIVRNAB HIV Viral Load	HIV QUANT <40 copies/mL HIV QUANT LOG <1.60 copies/mL	This test was performed using the Abbott RealTime HIV-1 PCR assay. The dynamic range for this assay is 40-10,000,000 copies/mL (1.60-7.00 Logs). This test is designed to provide information for monitoring disease progression and is not intended for use in the diagnosis of HIV-1 infection or Acquired Immune Deficiency Syndrome (AIDS).	Real-time polymerase chain reaction (RT-PCR)	Package Insert Validation	HIVRNA: 40-10,000,000 copies/mL HIVRNAL: 1.60-7.00 copies/mL HIVRNA: ≥40 copies/mL HIVRNAL: 1.60-7.00 copies/mL
HSV by PCR, Fluid/Lesion	HSVPCR Herpes Simplex Virus 1 and 2 Testing by PCR	Not Detected (for all targets)	This test was performed using a real time PCR assay.	Real-time polymerase chain reaction (RT-PCR)	Package Insert Literature	HSV1: Not Detected, Detected HSV2: Not Detected, Detected
Immunocompromised Respiratory Panel	ICRESP BioFire RP2.1	Not Detected (for all targets)	Results should be used in conjunction with other clinical and laboratory findings. This result does not rule out co-infections with pathogens that are not screened for by the Respiratory Panel (RP). This RP assay was performed using a Film Array multiplex nucleic acid assay.	Multiplex polymerase chain reaction (PCR)	Package Insert Literature	All Targets: Not Detected, Detected, Indeterminate
Influenza AB Rapid Molecular	FLUAIRM Rapid Flu	Not Detected (for all targets)	This test utilizes isothermal nucleic acid amplification technology for the differential qualitative detection of influenza A and influenza B viral nucleic acids.	Isothermal nucleic acid amplification	Akte-1 Influenza A+B package insert	All Targets: Not Detected, Detected, Indeterminate
Influenza AB, RSV By PCR	RVPCR Flu PCR RSV PCR	Not Detected (for all targets)	This test was performed using a multiplex real time PCR assay. This result does not rule out co-infections with pathogens that were not screened for by this test. Results should be used in conjunction with other clinical and laboratory findings.	Polymerase chain reaction (PCR)	Validation	All Targets: Not Detected, Detected, Indeterminate
Lactoferrin, Qualitative, Stool	WBKSTL Fecal Leukocytes Stool for WBK's	Negative		Immunochromatographic	Package Insert	Negative, Positive
Legionella Culture	LEGCX	No growth	Identification of Legionella species from respiratory specimens provides a definitive diagnosis of Legionnaires disease.	Culture		
Lower Respiratory Culture, Bacterial	RESPCX Respiratory Culture	No growth	Isolation of anaerobes in significant numbers from well-collected specimens indicates infection with the identified organisms.	Culture Susceptibility (if appropriate)		
M Tuberculosis Complex by PCR	MTBCOM TB PCR MTB	Not Detected	This test should always be performed in conjunction with AFB (acid fast bacilli) smear and culture. This test was performed using a Multiplex PCR Test.	Real-time polymerase chain reaction (RT-PCR)	Package Insert Reference Materials	Not Detected, Detected

Macroscopic Arthropod		Negative, Artifact (not arthropod)		Macroscopic exam			Negative, Bedbug, Centipede, Flea, Flea, Tunga sp, Flea, Ctenocephalides sp, Flea, Xenopsylla sp, Flea, Pulex sp, Ty, larva (anytime coming), Kissing Bug (Reduviid, Triatomine), Louse, Louse, Nit, Louse, Pediculus humanus, Louse, Phthirus pubis, Mite, Demodex sp, Mite, not Scabies, Mite-Sarcoptes scabiei, Scorpion, Spider, Spider-Black Widow, Spider-Brown Recluse, Hard tick, Hard tick, not Ixodes, Soft Tick, Tick, Tick-bodys sp, Tick-Dermacentor sp, Tick-Amblyomma sp, Tick-Rhipicephalus sp, Aradid (not arthropod), Arthropod, Arthropod or worm, other than a human parasite, Arthropod not known to transmit human pathogens
Maternal High Risk Star HIV	MHRHV			RHHV12: Non Reactive RFDV24: Non Reactive			RHHV12: Non Reactive, Reactive RFDV24: Non Reactive, Presumptive Reactive, Reactive
Meningitis/Encephalitis Panel CSF	CSFMEP Ref/Spec ME Panel	Not Detected (for all targets)		A negative result does not exclude the possibility of CNS infection and should not be used as the sole basis for diagnosis, treatment or other management decisions. Negative results may occur when the concentration of organism(s), virus(es), or yeast in the specimen is below the limit of detection. The ME panel does not distinguish between latent and active herpesvirus infections(CMV, HHV-6). This test was performed using a film array method for the detection of <i>Escherichia coli</i> , <i>K1</i> , <i>Haemophilus influenzae</i> , <i>Listeria monocytogenes</i> , <i>Neisseria meningitidis</i> , <i>Streptococcus agalactiae</i> , <i>Streptococcus pneumoniae</i> , <i>Cytomegalovirus</i> , <i>Enterovirus</i> , <i>Herpes Simplex virus 1 and 2</i> , <i>Human Herpesvirus 6</i> , <i>Human parechovirus</i> , <i>Varicella zoster virus</i> , and <i>Cryptococcus neoformans/gam</i> .	Multiplex polymerase chain reaction (PCR)	Package Insert	All Targets: Not Detected, Detected
Molecular Enteric Panel, Stool	STLB Gastrointestinal Panel	Negative (for all targets)		This test was performed using a real time PCR assay. Results should be interpreted in conjunction with clinical findings. A positive result does not necessarily indicate the presence of viable organisms. Positive results do not rule out co-infection with other organisms that are not detected by this assay. For Shiga toxin, this assay detects Shiga toxin 1 / Shiga toxin 2 genes (found in Shiga toxin-producing <i>E. coli</i> as well as <i>Shigella dysenteriae</i>). This test should not be used as a test of cure.	Real-time polymerase chain reaction (RT-PCR)	Package Insert	All Targets: Negative, Positive
Molecular Stool Parasite Panel	SCPD O&P	Negative (for all targets)		This test was performed using a real time PCR assay. Results should be interpreted in conjunction with clinical findings. A positive result does not necessarily indicate the presence of viable organisms. Positive results do not rule out co-infection with other organisms that are not detected by this assay. This test should not be used as a test of cure.	Real-time polymerase chain reaction (RT-PCR)	Package Insert Reference Materials	All Targets: Negative, Positive
Neisseria Gonorrhea Amplified Probe	NG	Not Detected	A negative test result for Chlamydia trachomatis does not preclude the possibility of infection. Results should be considered in conjunction with other clinical and laboratory findings.	This test was performed using Transcription Mediated Amplification for the detection of Chlamydia trachomatis and/or Neisseria gonorrhoeae nucleic acid. This assay is not intended for the evaluation of suspected sexual abuse or other medico-legal indications.	Transcription-mediated amplification (TMA)		Not Detected, Detected
Neisseria-Gonococcus Screen	SCN	No growth		Isolation of anaerobes in significant numbers from well-collected specimens including endocervix, urethra, rectum, upper respiratory tract (throat, nasopharynx), and conjunctiva indicates infection with the identified organisms.	Culture		
Novel Coronavirus PCR - Semi-Private Surveillance	COVID	Not Detected		Negative results do not preclude SARS-CoV-2 infection and should not be used as the sole basis for treatment or other patient management decisions. Optimum specimen types and timing for peak viral levels during infections caused by SARS-CoV-2 has not been determined. The possibility of a false negative result should especially be considered if the patient's recent exposures or clinical presentation suggest that SARS-CoV-2 infection is probable, and diagnostic tests for other causes of illness (e.g., other respiratory illness) are negative. Collection of a new specimen and re-testing may be necessary if the patient is critically ill or clinically deteriorating.	Real-time polymerase chain reaction (RT-PCR)		Not Detected, Detected
Novel Coronavirus PCR - Nasopharyngeal	COVID	Not Detected		Negative results do not preclude SARS-CoV-2 infection and should not be used as the sole basis for treatment or other patient management decisions. Optimum specimen types and timing for peak viral levels during infections caused by SARS-CoV-2 has not been determined. The possibility of a false negative result should especially be considered if the patient's recent exposures or clinical presentation suggest that SARS-CoV-2 infection is probable, and diagnostic tests for other causes of illness (e.g., other respiratory illness) are negative. Collection of a new specimen and re-testing may be necessary if the patient is critically ill or clinically deteriorating.	Real-time polymerase chain reaction (RT-PCR)		Not Detected, Detected
Novel Coronavirus PCR - Nasopharyngeal (Outreach)	COVID	Not Detected		Negative results do not preclude SARS-CoV-2 infection and should not be used as the sole basis for treatment or other patient management decisions. Optimum specimen types and timing for peak viral levels during infections caused by SARS-CoV-2 has not been determined. The possibility of a false negative result should especially be considered if the patient's recent exposures or clinical presentation suggest that SARS-CoV-2 infection is probable, and diagnostic tests for other causes of illness (e.g., other respiratory illness) are negative. Collection of a new specimen and re-testing may be necessary if the patient is critically ill or clinically deteriorating.	Real-time polymerase chain reaction (RT-PCR)		Not Detected, Detected
Pinworm Exam	Pinworm Exam	Negative			Microscopic exam		Negative, Enterobius vermicularis (Pinworm) Eggs Seen, Enterobius vermicularis (Pinworm) Adult Worm(s) Seen, Enterobius vermicularis (Pinworm) adult worm(s) and eggs seen
Plesiomonas/Aeromonas Screen, Stool	SCF	Negative for Aeromonas species Negative for Plesiomonas species		Isolation of anaerobes in significant numbers from well-collected stool specimens indicates infection with the identified organisms.	Culture		
Quantitative Tissue Culture	BKR QUANTITA	No growth		Isolation of anaerobes in significant numbers from well-collected specimens indicates infection with the identified organisms.	Tissue is weighed, serially diluted, and cultured for exact colony count.		

Rapid HIV-1/HIV-2 Ab With P24 Antigen	RHIV12 Aline Determination HIV 1/2 Ag/Ab Combo		RHIV12: Non Reactive RPDP24: Non Reactive	Qualitative immunoassay / immunochromatographic test for simultaneous and qualitative detection of free HIV-1 p24 antigen and antibodies to HIV-1 and HIV-2.	Package Insert		RHIV12: Non Reactive, Reactive RPDP24: Non Reactive, Presumptive Reactive, Reactive.	
Rapid Malaria	MPBR EVIDMAL Plasmodium STRIPAM Rapid Strip Strip A	Negative for malaria antigens		Immunochromatographic membrane assay	Package Insert		Negative for malaria antigens. Positive rapid malaria test for P. falciparum protein antigen only. Positive rapid malaria test for malaria protein antigens representing P. vivax or P. malariae or P. ovale or a mix of species. Positive rapid malaria test. Positive for P. falciparum protein antigen	
Rapid Strep A, Molecular	STRIPAM Rapid Strip Strip A	Negative		Molecular in vitro diagnostic test utilizing isothermal nucleic acid amplification.			Negative, Positive	
Rectal Screening for Cipro Resistance	BKR CIPROFLO Ciprofloxacin Resistance Screening	Negative for Ciprofloxacin Resistant Enterobacterias		Isolation of anaerobes in significant numbers from well-collected specimens indicates infection with the identified organisms.	Culture			
SARS-COV-2 Rapid	COVID Rapid COVID	Not Detected		Negative results do not preclude SARS-CoV-2 infection and should not be used as the sole basis for treatment or other patient management decisions. Optimum specimen types and timing for peak viral levels during infections caused by SARS-CoV-2 has not been determined. The possibility of a false negative result should especially be considered if the patient's recent exposure or clinical presentation suggest that SARS-CoV-2 infection is probable, and diagnostic tests for other causes of illness (e.g., other respiratory illness) are negative. Collection of a new specimen and re-testing may be necessary if the patient is critically ill or clinically deteriorating.	Isothermal nucleic acid amplification	Abbott ID NOW Rapid Covid package insert	Not Detected, Detected	
Screen: MRSA Only	SCREEN: MRSA MRSA Screen	Negative		Culture			Negative, Positive	
Screen: MRSA/MSSA	SCREEN: MRSA MRSA Screen	PCR Negative CLITHIRK Negative for Staphylococcus aureus Negative for Methicillin Resistant Staphylococcus aureus		This test was performed using a real time PCR assay. Results should be interpreted in conjunction with other clinical and laboratory findings. A positive result does not necessarily indicate the presence of viable organisms. This test should not be used as a test of cure.	Real-time polymerase chain reaction (RT-PCR)	Package Insert	PCR: Negative, Positive MSSA CX: Negative for Staphylococcus aureus Positive for Staphylococcus aureus MRSA CX: Negative for Methicillin Resistant Staphylococcus aureus Positive for Methicillin Resistant Staphylococcus aureus	
Screen, VRE	SCREEN: VRE Vancomycin Resistant Enterococcus Screen	Negative		Culture on select agar			Negative, Positive	
Screen, Yeast	YESTSCR	Negative		Culture				
Strep Pneumoniae Antigen, Urine	PNELMU	Negative		Immunochromatographic membrane assay	Binax NOW Package Insert		Negative, Positive	
Upper Respiratory Culture, Bacterial	THIRCS Throat Culture RESN	No growth		Isolation of anaerobes in significant numbers from well-collected specimens indicates infection with the identified organisms.	Culture			
Urine Culture	URN CX	Negative		Isolation of anaerobes in significant numbers from well-collected urine specimens indicates infection with the identified organisms.	Culture Susceptibility (if appropriate)			
Vaginal Infection Panel	BVAG CV TV CT NG	BVAG: Negative CV: Not Detected TV: Not Detected CT: Not Detected NG: Not Detected		BV: This test was performed using Transcription Mediated Amplification for the detection of ribosomal RNA from bacteria associated with bacterial vaginosis. CV/TV: This test was performed using Transcription Mediated Amplification for the detection of ribosomal RNA from microorganisms associated with vulvovaginal candidiasis and trichomoniasis. CT/NG: This test was performed using Transcription Mediated Amplification for the detection of Chlamydia trachomatis and/or Neisseria gonorrhoeae nucleic acid. This assay is not intended for the evaluation of suspected sexual abuse or other medico-legal indications.		See individual analytes	See individual analytes	See individual analytes
Varicella Zoster By PCR, Skin	VZV/PCR	Not Detected		This test was performed using a real time PCR assay.	Real-time polymerase chain reaction (RT-PCR)	Package Insert Literature	Not Detected, Detected	
ADAMTS13 Activity and IgG Ab w/ Reflex to Inhibitor	AD13A AD13AG was Wilchbrand Factor Cleaving Protease Anti-ADAMTS13 antibodies	ACTIVITY >40 % IQC ≤12.0 U/mL		Technozym enzyme-linked immunosorbent assay (ELISA)	Technozym kit	ACTIVITY: 2-100 % IQC: 6.0-104.0 U/mL	ACTIVITY: 2-100 % IQC: 6.0-104.0 U/mL	
Alternative Activation Pathway	APBHD Bb Complement Alternative complement pathway	695-1,974 ng/mL		Quidel enzyme-linked immunosorbent assay (ELISA)	Biomarker Reference Lab			
Terminal Activation Pathway	TPCSAB SC5b-9 Complement Membrane attack complex (MAC) Terminal complement complex (TCC)	6-598 ng/mL		Quidel enzyme-linked immunosorbent assay (ELISA)	Biomarker Reference Lab			